


This article was downloaded by: [Erciyes University]

On: 08 January 2015, At: 16:50

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

IMPROVED SYNTHESIS OF O-BENZENEDISULFONIMIDE

Franklin A. Davis^a, Gajendran Sundarababu^a & Hongyan Qi^a

^a Department of Chemistry, Temple University, Philadelphia, PA, 19122, USA

Published online: 09 Feb 2009.

To cite this article: Franklin A. Davis, Gajendran Sundarababu & Hongyan Qi (1998) IMPROVED SYNTHESIS OF O-BENZENEDISULFONIMIDE, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 30:1, 107-109, DOI: [10.1080/00304949809355269](https://doi.org/10.1080/00304949809355269)

To link to this article: <http://dx.doi.org/10.1080/00304949809355269>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.


This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

IMPROVED SYNTHESIS OF *o*-BENZENEDISULFONIMIDE

Submitted by Franklin A. Davis*, Gajendran Sundarababu and Hongyan Qi
(08/05/97)

Department of Chemistry, Temple University, Philadelphia, PA 19122, USA

N-Fluoro-*o*-benzenedisulfonimide (NFOBS, **5b**) is an important electrophilic source of fluorine which efficiently monofluorinates carbanions and enolates.¹ This reagent is prepared in >90% yield by treatment of *o*-benzenedisulfonimide (**5a**) with 10% F₂/N₂.^{1b} N-Hydroxy-*o*-benzenedisulfonimide (NHOBS, **5c**) is a selective oxidizing reagent for the oxidation of thiols to disulfides, sulfides to sulfoxides and benzyl alcohols to aldehydes.² It is prepared from *o*-benzenedisulfonyl chloride by reduction with sodium sulfite, followed by nitrosation.³ The key intermediate in both these syntheses is *o*-benzenedisulfonyl chloride (**4**) which can be prepared from the disulfonic acid **3**⁴ or from a 1,2-benzenedithiol derivative.⁵ However, the necessity for preparing **4** from anthranilic acid⁵ or aniline-2-sulfonic acid^{5,6} via diazonium salts is inefficient and limits the scale on which NFOBS **5b** and NHOBS **5c** can be synthesized. This paper describes an efficient four-step synthesis of *o*-benzenedisulfonimide (**5a**) in 50% overall yield, from lithium benzenesulfonate (**1**).⁷



Our synthesis of **5a** relies on the *ortho* lithiation of lithium benzenesulfonate (**1**) with *n*-butyllithium first reported by Figuly and Martin.⁷ Thus, treatment of a suspension of **1** at 0° in THF with 1.1 equivalents of *n*-butyllithium rapidly generated the *ortho* lithio species as indicated by the formation of a yellow colored, homogenous solution. Addition of 2 equivalents of elemental sulfur to the dilithio intermediate afforded, after quenching with 3 N HCl and removal of the solvent, a pale yellow solid (Scheme 1). The ¹H NMR indicated that it was an 88:12 mixture of the disulfide **2** and the starting material **1**. The two sets of protons in **1**, appearing as multiplets at δ 7.5 and 7.75, are clearly distinguishable from the protons in **2**. Since separation of **2** from the lithium sulfonate **1** proved impractical the mixture was used without purification and oxidized at 90° with potassium

permanganate (KMnO_4). Removal of the aqueous solvent gave the crude sulfonic acid salt **3** which was dried under vacuum and heated at 110 – 170° , as previously described, with PCl_5 to give the disulfonyl chloride **4**.^{1b} Treatment of **4** with EtOH/NH_3 afforded **5a** in 49% overall yield, following purification, from the lithium sulfonate **1**. Since none of the intermediates were purified until the end, this represents a highly efficient, safe synthesis of *o*-benzenedisulfonimide (**5a**).

EXPERIMENTAL SECTION

Benzenesulfonic acid and *n*-butyllithium were purchased from Aldrich Chemical Co. Evaporations were carried out on a rotary evaporator using an aspirator.

CAUTION: Phosphorus pentachloride is toxic. All operations were conducted in a good fume hood.

***o*-Benzenesulfonic Acid Disulfide (2).**— To a cold (0°) suspension of dry lithium benzenesulfonate⁷ (**2**; 1.64 g, 10 mmol) in dry THF (100 mL) under an inert atmosphere was added *n*-BuLi (3.7 mL of 3M hexane solution, 11 mmol) dropwise over 5 min. The pale yellow suspension became homogeneous in about 10 min. and after stirring for 5 min. elemental sulfur (0.64 g, 20 mmol) was added *via* a solid addition funnel in one portion. The brown reaction mixture was allowed to warm to rt and stirred for 15 h. The clear brown solution was quenched with 3N HCl (50 mL). The solution was concentrated, H_2O (50 mL) was added to the syrupy residue and the mixture was extracted with ether (3 x 75 mL). The organic phase was washed with 3N HCl (30 mL). The combined aqueous phases were concentrated and dried under vacuum to afford 2.4 g of **2** as a hygroscopic, pale yellow solid contaminated with **1** in a ratio of 88:12. ^1H NMR (500 MHz, D_2O): δ 7.82 (dd, 1H, $J = 7.7$ and 1.5 Hz); 7.43 (dd, 1H, $J = 7.7$ and 1.1 Hz); 7.34 (dt, 1H, $J = 7.7$ and 1.5 Hz); 7.24 (dt, 1H, $J = 7.7$ and 1.1 Hz)

***o*-Benzenedisulfonimide (5a).**— To a hot (90°) aqueous solution (30 mL) of the crude disulfide **2** was added saturated aqueous KMnO_4 (10 mL) until a pale pink color persisted in the reaction mixture. After 10 min. at 90° , the excess permanganate was destroyed by addition of *i*-PrOH (2–3 mL). The reaction mixture was cooled to rt, the brown solid was removed by filtration and rinsed with H_2O (3 x 10 mL). The aqueous solution was concentrated to give a white solid which was dried under vacuum for 48 h. The crude disulfonic acid salt **3** was heated, as previously described,^{1b} with PCl_5 (5.12 g, 25 mmol) first at 110° and then at 170° to give the crude disulfonyl chloride **4** which was treated with ammonia (3.1 g in 50 mL of EtOH) at 0° followed by purification of the aqueous extract by ion exchange (Dowex resin 50 x 8–100) to afford 1.08 g (49%) of **5a** as a white solid. mp. 190 – 192° , lit.^{1b} 192 – 194° .

Acknowledgment.— This work was supported by a grant from the National Science Foundation.

REFERENCES

1. a) F. A. Davis and H. Qi, *Tetrahedron Lett.* **37**, 4345 (1966). b) F. A. Davis, W. Han and C. K. Murphy, *J. Org. Chem.*, **60**, 4730 (1995). c) V. Snieckus, F. Deaubieu, K. Mobri, W. Han, C. K. Murphy and F. A. Davis, *Tetrahedron Lett.* **35**, 3465 (1994). d) F. A. Davis and R. E. Reddy, *Tetrahedron: Asymmetry*, **6**, 955 (1994). e) F. A. Davis and W. Han, *Tetrahedron Lett.*, **33**, 1153 (1992).

2. M. Barbero, I. Degani, R. Fochi and P. Perracino, *J. Org. Chem.*, **61**, 8762 (1996).
3. a) W. R. H. Hurtley and S. Smiles, *J. Chem. Soc.*, 1821 (1926). b) J. B. Hendrickson, S. Okano, and R. K. Bloom, *J. Org. Chem.*, **34**, 3434 (1969). c) J. L. Kice and S-t. Liao, *ibid.*, **46**, 2691 (1981).
4. W. E. Parham, T. M. Roder and W. R. Hasek, *J. Am. Chem. Soc.*, **75**, 1647 (1953).
5. I. Degani and R. Fochi, *Synthesis*, 471 (1976).
6. M. Barbero, I. Degani, R. Fochi and V. Regondi, *Gazz. Chim. Ital.*, **116**, 165 (1986).
7. Figuly, G. D.; Martin, J. C. *J. Org. Chem.*, **45**, 3728 (1980).

1-ALKYLIDENE(ARYLIDENE)AMINO-2-AMINOETHANES AND THEIR TAUTOMERIZATION TO IMIDAZOLIDINES

Submitted by
(11/21/96)

Kirill N. Zelenin*

*Department of Chemistry, Military Medical Academy
St. Petersburg 194175, RUSSIA*

Ilya V. Ukraintzev

St. Petersburg Technological Institute 198013, RUSSIA

In theory, the interaction of ethylenediamine (1) with aldehydes and ketones could lead to the corresponding mono-imines (2). In practice, however, the *bis*-imines are actually obtained with aromatic¹⁻⁴ and aliphatic aldehydes,⁴ aliphatic ketones,^{2,5} acetophenone² and benzophenone.² The mono-imines have been isolated only with cyclohexanone and its homologues. The formation of mono-mines has been claimed with formaldehyde⁹ and acetone⁶ but no products were isolated. The instability of these mono-imines has been ascribed to their tendency toward hydrolysis and polymerization. Chromatographic methods and vacuum distillation are not useful for isolation because of the decomposition of these substances on the adsorbent surface, even at relatively low temperatures.⁵

We carried out several PMR-monitored experiments of the reaction mixtures of diamine 1 with typical carbonyl compounds, such as benzaldehyde and acetone, in order to devise a synthetic path to 1-monoalkylidene derivatives 2. The *bis*-derivative 3a predominates in the reaction mixture of benzaldehyde with ethylenediamine in a 1:1 proportion. Although 1-arylideneamino-2-aminoethane (2e) is present in less than 10%, it becomes the only product when the proportion