Branch, Division of Research Resources, National Institutes of Health. under Grant No. PR-330.

> M. C. Wani,\* H. L. Taylor, Monroe E. Wall Chemistry and Life Sciences Laboratory **Research Triangle Institute** Research Triangle Park, North Carolina 27709

P. Coggon, A. T. McPhail Paul M. Gross Chemical Laboratory Duke University, Durham, North Carolina 27706 Received February 16, 1971

## Photocyclization of Aryl-Substituted Acetylenes; Application of Di- $\pi$ -methane-like Rearrangements to Arylcyclopropene Syntheses<sup>1</sup>

Sir:

In a continuation of previous studies on the photocyclization of aryl-substituted propenes<sup>2</sup> and related functionalized propenyl systems, such as 3-alkoxysubstituted propenes,<sup>3</sup> we have investigated the photochemistry of several aryl-substituted acetylenes with the intent of developing a convenient route to cyclopropenes including functionalized systems which in turn might serve as precursors for cyclopropenium derivatives.<sup>4</sup> Our continuing interest in the photo-chemistry of cyclopropenes<sup>5a-d</sup> provided an additional incentive for this study.



Tetraphenylpropyne (1a) was synthesized by a procedure<sup>6a</sup> which is a modification of the Wieland and

(1) A preliminary report of this work was presented at the 160th National Meeting of the American Chemical Society, Chicago, Ill., Sept 1970.

(2) G. W. Griffin, A. F. Marcantonio, H. Kristinsson, R. C. Petterson, and C. S. Irving, Tetrahedron Lett., 2951 (1965).

(3) J. J. Brophy and G. W. Griffin, ibid., 493 (1970).

(4) Recently, J. W. Wilson and K. L. Huhtanen (Chem. Commun., 454 (1968)) described the photochemical rearrangement of methyl 3,3,3triphenylpropyne carboxylate (i) to the indenophenanthrene (iv). The diradical species ii was proposed as a possible intermediate leading to the indene iii which in turn undergoes subsequent dehydrocyclization to iv. A similar diradical species has been postulated by us as an intermediate in the thermal<sup>5a</sup> and photochemical<sup>6b</sup> rearrangement of tetraphenylcyclopropene (2a) to triphenylindene 4.



(5) (a M. A. Battiste, B. Halton, and R. H. Grubbs, Chem. Com-mun., 907 (1967); (b) R. H. Grubbs, M.S. Thesis, University of Florida, 1965; (c) H. Kristinsson and G. W. Griffin, J. Amer. Chem. Soc., 88, 1579 (1966); (d) H. Kristinsson, Tetrahedron Lett., 2343 (1966); (e) A. S. Monahan, J. D. Freilich, and J. J. Fong, ibid., 1865 (1970); (f) H. Dürr, Justus Liebigs Ann. Chem., 723, 102 (1969). (6) (a) A. W. Herriot, Ph.D. Thesis, University of Florida, 1967;

(b) H. Wieland and H. Closs, Justus Liebigs Ann. Chem., 470, 201 (1929).

Closs method.<sup>6b</sup> Irradiation of **1a** (500 mg) in 500 ml of cyclohexane (0.003 M) at 253.7 nm for relatively short periods (3.4 hr) in a preparative photochemical reactor<sup>7a</sup> while simultaneously sparging with a slow stream of argon provided 230 mg (46%) of 2a, mp 177.5-9° (lit.<sup>5b</sup> mp 176-178°) which may be isolated by elution chromatography on alumina.

For convenience the mechanism for the rearrangement of 1a is formulated as a diradical process as outlined below.4



The formation of 2a indicates that the intermediate, perhaps 3b, formed by rearrangement of a diradical-like intermediate such as 3a cyclizes at least in part in a process analogous to the photochemical rearrangement of propenes to cyclopropanes.<sup>2,3</sup> Alternatively, and perhaps more likely, the reaction proceeds in a concerted manner.8

Additional photoproducts begin to form when the irradiation time is extended as evidenced by the appearance of nmr signals at  $\tau$  4.94 and 4.67 (CDCl<sub>3</sub>), characteristic of the benzyl protons of 4 and 5, respectively.<sup>5a,b</sup> An nmr study<sup>9a</sup> of the variation in product composition as a function of time under standard irradiation conditions in benzene<sup>7b</sup> proved instructive. The quantitative data summarized in Table I were

Table I	l
---------	---

Time, hr	0/0/		
	2a	<b>4</b> ´°	5
24	25	42	19
4	46	35	9
1	70	Trace	Trace

determined utilizing the four-proton aromatic multiplet for 2a centered at  $\tau$  2.30 (CDCl<sub>3</sub>) in addition to the singlets for the benzyl protons of 4 and 5. Although these data confirm that the photolysis of 1a in cyclohexane<sup>7b</sup> proceeds in high conversion ( $\sim 80\%$ ) upon

(7) (a) The preparative irradiations were conducted in a Rayonet RPR 208 Reactor (The Southern New England Ultraviolet Co., Middletown, Conn.) with 8 RUL 253.7-nm lamps; (b) an RPR-100 unit equipped with 16 8-W 253.7-nm lamps; (c) an RPR-100 unit fitted with 16 8-W 310.0- or 350.0-nm lamps

(8) R. B. Woodward and R. Hoffmann, Angew. Chem., Int. Ed. Engl., 8, 781 (1969).

(9) (a) All nmr spectra were recorded on a Varian A-60 instrument in the solvent indicated with tetramethylsilane as an internal standard; (b) all new compounds gave satisfactory combustion and/or mass spectral analyses; (c) mass spectral data were obtained on a Perkin-Elmer Hitachi RMU-6 spectrometer.

prolonged irradiation (24 hr) the separation of the products by column chromatography proved difficult. In general, the fractions contained at least two components; however with care, 2a, 4, and 5 were separated in a state of purity sufficient for definitive characterization albeit in low yields (8, 7, and 4%, respectively).

The presence of tetraphenylcyclopropene (2a) in aliquots of irradiated cyclohexane solutions<sup>7b</sup> of 1a is also evident from the ultraviolet absorption maximum apparent at 334 m $\mu$  (lit.<sup>5b</sup>  $\lambda_{max}$  334 m $\mu$  ( $\epsilon$ 23,700)). The photorearrangement of 1a was monitored at 15-min intervals by ultraviolet spectroscopy and in agreement with the nmr data 2a is the major product and only traces of 4 and 5 are formed during the first hour of irradiation. These results indicate that the indene 4 and indenophenanthrene 5 obtained upon photolysis of 1a are secondary products arising from rearrangement of cyclopropene 2a, a result clearly consistent with the known photolability of this and other cyclopropenes (vide infra).<sup>4,5</sup>

No cyclopropenyl or indenyl products are observed when **1a** is irradiated<sup>7c</sup> in a Pyrex vessel for up to 20 hr in acetone solution or in the presence of such sensitizers as benzophenone or acetophenone. This result is indicative that the rearrangement of tetraphenylpropyne (1a) to tetraphenylcyclopropene (2a) proceeds by a singlet mechanism.

The functionalized acetylene 1b also cyclizes upon irradiation to give a cyclopropene 2b, which is a convenient precursor for cyclopropenium salts.<sup>10</sup> 1,3,3-Triphenylprop-1-yn-3-ol,<sup>11</sup> mp 79-80° (lit.<sup>11</sup> mp 82°), prepared by addition of phenylethynylmagnesium bromide to benzophenone and subsequent hydrolysis. was converted to the methyl ether (mp  $122-123^{\circ}$ ) in high yield by treatment with methyl iodide in dimethyl sulfoxide under basic conditions according to the method of Gillis.<sup>12</sup> Irradiation<sup>7b</sup> of degassed solutions of 1b (0.011 M in benzene) gives methoxycyclopropene (2b) in an amount which also is dependent on the duration of irradiation. The nmr signal at  $\tau$  6.66 arising from the methoxy protons of 2b as well as the benzyl proton signal at  $\tau$  5.30 may be differentiated from the aliphatic signal of the precursor 1b at  $\tau$  6.59 after 10-hr irradiation provided the spectra are determined in benzene, the irradiation solvent. The yield of 2b does not exceed 5% and at this time the appearance of an additional photoproduct (17%) was evident because of the emergence of a second methoxy signal at  $\tau$  6.37. This product was shown to be the indenyl ether 8a by hydrolysis to 2,3-diphenyl-1indanone identical in all respects with an authentic sample of this material.<sup>13</sup> The ratio of 8a to 1b as determined by nmr reaches 1:1.7 after photolysis (53 hr).<sup>7b,14</sup> That **8a** may arise from **2b** in a secondary photoprocess was confirmed by irradiation of 2b in benzene for 16  $hr^{7b}$  whereupon the conversion to 8a reaches 43% The indenyl ether 8c was prepared

(14) We found that tractable quantities of 8a were conveniently prepared in high yield (86.5%) by thermolysis of 2b at 250° (10 min). Additional products included 1-methoxy-2,3-diphenylindene (8c)  $(\sim 3\%)$  and an oxidation product, 2,3-diphenyl-1-indanone (9.5%). The thermal reactions of 2a, 2b, and a variety of other cyclopropenes including the cyano analog of 2b will be the subject of a subsequent communication.



independently from 2,3-diphenyl-1-indenone<sup>15</sup> by reduction with sodium borohydride in methanol followed by methylation with diazomethane and boron trifluoride etherate<sup>16</sup> and the characteristic nmr signals at  $\tau$  4.40 and 7.03 (benzyl and methoxy protons, respectively) were conspicuously absent from the benzene solution obtained upon photolysis of 2b or for that matter 1b.7b This is not unexpected since opening and recyclization of 2b would give the isoindene intermediate 7 which can undergo a thermally allowed 1,5-sigmatropic shift<sup>8</sup> with aromatization leading to 8a.<sup>17</sup> In contrast, formation of 8c from 8a, in the absence of base, requires two additional successive 1,5 shifts involving formation of an o-xylylene intermediate<sup>18</sup> and concomitant loss of aromaticity, an overall process which is energetically unfavorable at room temperature, but does occur to some extent at elevated temperatures.<sup>14</sup> It has been demonstrated previously<sup>10</sup> that 2b is readily converted to triphenylcyclopropenium bromide by treatment with anhydrous hydrogen bromide and thus a new route to these ionic species becomes available if convenient separation techniques can be devised.

Irradiation of **1b** in cyclohexane<sup>7b</sup> leads to a complex mixture of products including 8d and perhaps 2d although further studies are necessary to characterize the reaction products obtained in this solvent fully.

An examination of the photochemical behavior of the arylpropynes  $R_1R_2R_3CC \equiv CC_6H_6$  where  $R_1 = R_2 =$  $R_3 = H$  (9),  $R_1 = R_2 = H$ ,  $R_3 = C_6H_5$  (10), and  $R_1 = R_2 = C_6H_6$ ,  $R_3 = H$  (11) also has been performed; however no products analogous to those obtained with 1a and 1b were detected. In fact 9 is relatively photostable<sup>7b</sup> while 10 and 11 gave products whose structures remain to be determined.19

Acknowledgment. We wish to thank Dr. N. Portnoy, Dr. R. Bost, Dr. D. J. Keyton, and Mrs. J. Thompson for invaluable technical assistance provided during the course of this work. Support for this research by the National Science Foundation (Grants GP 9434 and GP 9412), the Petroleum Research Fund (PRF 2813-

(19) J. J. Brophy and G. W. Griffin, unpublished results.

<sup>(10)</sup> R. Breslow and C. Yuan, J. Amer. Chem. Soc., 80, 5991 (1958).

 <sup>(11)</sup> K. Hess and W. Weltzien, Chem. Ber., 54, 2511 (1921).
 (12) R. G. Gillis, Tetrahedron Lett., 1413 (1968).

<sup>(13)</sup> C. F. Koelsch, J. Amer. Chem. Soc., 56, 1337 (1934).

<sup>(15)</sup> K & K Laboratories, Inc., Plainview, N. Y.
(16) M. C. Caserio, J. D. Roberts, M. Neeman, and W. S. Johnson, J. Amer. Chem. Soc., 80, 2584 (1958); E. Müller and W. Rundel, Angew. Chem., 70, 105 (1958).

<sup>(17)</sup> Monahan and coworkers have shown in independent studies that 2c, the ethoxy analog of 2b, is converted to the ethoxy indene 8b in 70% yield upon irradiation in benzene under nitrogen with 350nm lamps.<sup>5</sup>e (18) W. Roth, *Tetrahedron Lett.*, 1009 (1964).

A1), and the Alfred P. Sloan Foundation is also gratefully acknowledged.

> B. Halton, M. Kulig, M. A. Battiste\* Department of Chemistry, University of Florida Gainesville, Florida 32601

J. Perreten, D. M. Gibson, G. W. Griffin\* Department of Chemistry Louisiana State University in New Orleans Lake Front, New Orleans, Louisiana 70122 Received January 21, 1971

## Stereochemistry of [2.2]Cyclophane Rearrangements<sup>1</sup>

Sir:

We report the acid-catalyzed rearrangement of optically pure (+)-(S)-4-methyl[2.2]paracyclophane ((+)-1) to optically pure (+)-(S)-12-methyl[2.2]metaparacyclophane ((+)-2), the photolytic conversion of [2.2]metaparacyclophane (3) to [2.2]metacyclophane (4), and the photolytic racemization and rearrangement of (-)-2 to a mixture of methyl[2.2]metacyclophanes (5).



Optically pure (+)-(S)-1<sup>2</sup> [mp 150–152°,  $[\alpha]^{25}_{546}$  $+114^{\circ}$  (c 1.0 CHCl<sub>3</sub>)] (the configuration of (-)-(R)-1 is established)<sup>3a</sup> was prepared<sup>3b</sup> and submitted to the acid conditions used to rearrange [2.2]paracyclophane to [2.2] metaparacyclophane.<sup>4</sup> The product, (+)-2, was isolated (52%) by chromatographic and sublimation procedures free of optical fractionation, mp 61-64.5°,  $[\alpha]^{25}_{546} + 27.0^{\circ}, \ [\alpha]^{25}_{436} + 28.0^{\circ} \ (c \ 1.02 \ \text{CHCl}_3).$  Recrystallization gave material, mp 66-67°, rotation unchanged. The enantiomer, (-)-2, was prepared from (+)-12-carboxy[2.2]metaparacyclophane ((+)-6),<sup>5a</sup> mp  $164-168^{\circ}, [\alpha]_{546}^{25}+33.7^{\circ}, [\alpha]_{436}^{25}+115^{\circ} (c \ 1.0 \ \text{CHCl}_3).$ Reduction of this acid with lithium aluminum hydride

(1) The authors warmly thank the National Science Foundation for a grant that supported part of this research. M. H. Delton thanks the Regents of the University of California for a First Year Fellowship, and R. E. Gilman thanks the National Science Foundation for a Science Faculty Fellowship, 1970-1971.

(2) All new compounds gave carbon and hydrogen analyses within

(3) % of theory and nmr spectra expected for the assigned structures.
(3) (a) M. J. Nugent and O. E. Weigang, Jr., J. Amer. Chem. Soc., 91, 4557 (1969);
(b) M. H. Delton and D. J. Cram, *ibid.*, 92, 7623 (1970).
(4) D. J. Cram, R. C. Helgeson, D. Lock, and L. A. Singer, *ibid.*,

88, 1324 (1966). (5) (a) D. T. Hefelfinger and D. J. Cram, *ibid.*, in press; (b) *ibid.*, in

press.

(in THF) gave (64%) (-)-12-hydroxymethylene[2.2]metaparacyclophane<sup>2</sup> ((-)-7), mp 99–100°,  $[\alpha]^{25}_{546}$  $-7.3^{\circ}$ ,  $[\alpha]^{25}_{436} + 0.2^{\circ}$ ,  $[\alpha]^{25}_{365} + 38.2^{\circ}$  (c 1.0 CHCl<sub>3</sub>), which with phosphorus tribromide in methylene dichloride gave (67%) (+)-12-bromomethylene[2.2]metaparacyclophane<sup>2</sup> ((+)-8), mp 92–93°,  $[\alpha]^{25}_{546} \pm 85.7^{\circ}$ ,  $[\alpha]^{25}_{436}$  +193° (c 1.0 CHCl<sub>3</sub>). Reduction of this bromide with lithium aluminum hydride (THF) gave (60%) (-)-2,<sup>2</sup> mp 65-66°,  $[\alpha]^{25}_{546}$  -26.3°,  $[\alpha]^{25}_{436}$  $-27.4^{\circ}$  (c 1.0 CHCl<sub>3</sub>). At each stage, the above compounds were recrystallized to maximum melting points and rotations. Admixture of equal amounts of the above samples of (-)- and (+)-2 gave mp 43-45°, undepressed by admixture with authentic  $(\pm)$ -2.<sup>5b</sup> The fact that the rotations of 2 prepared by rearrangement of (+)-1 and from (+)-6 are of equal magnitude indicates that (+)-1, (+)-2, and (-)-2 were essentially optically pure, and that  $(+)-1 \rightarrow (+)-2$  proceeded stereospecifically.

Of the two most likely general mechanisms formulated, these results eliminate the phenonium ion route  $((+)-1 \rightarrow A \rightarrow D \rightarrow (\pm)-C \rightarrow (\pm)-2)$ , since D possesses a mirror plane, and only  $(\pm)$ -2 would be produced. The route (+)-(S)-1  $\rightarrow$  A  $\rightarrow$  B  $\rightarrow$  C  $\rightarrow$ (+)-2 provides the required fully chiral reaction coordinate if rotation of the methyl-substituted benzene ring with respect to the other benzene ring is sterically inhibited in A, B, and C. Molecular models point to such steric inhibition of rotation, and such rotation in (+)-12-carbomethoxy[2.2]metaparacyclophane does not occur up to 200°.5ª Structure B might be an intermediate or a transition state in this mechanism.

The absence of alternate mechanisms with fully chiral reaction coordinates allows assignment of configuration to (+)-2. The fact that the meta ring of 2 rotates with respect to the para ring at 25° to provide a conformational equilibrium mixture<sup>5a,6</sup> complicates the structure and name given to compound (+)-2. However, the facts that the equilibrium mixture is dominated by the form with the methyl anti to the meta ring and crystallizes only in that form<sup>5a,6</sup> suggest that the structure written for (+)-2 is the more appropriate and this structure possesses the S configuration.<sup>7</sup> This assignment fixes the configurations of (+)-6, (-)-7, and (+)-8 as R.



<sup>(6)</sup> D. T. Hefelfinger and D. J. Cram, *ibid.*, 92, 1073 (1970).
(7) R. S. Cahn, C. K. Ingold, and V. Prelog, *Angew. Chem.*, *Int. Ed.* Engl., 5, 400 (1966).