ChemComm

This article is part of the

Porphyrins & Phthalocyanines web themed issue

Guest editors: Jonathan Sessler, Penny Brothers and Chang-Hee Lee

All articles in this issue will be gathered together online at <u>www.rsc.org/porphyrins</u>



Cite this: Chem. Commun., 2012, 48, 4317-4319

COMMUNICATION

A doubly 2,6-pyridylene-bridged porphyrin–perylene–porphyrin triad[†]‡

Shin Ikeda,^a Naoki Aratani*^{ab} and Atsuhiro Osuka*^a

Received 10th February 2012, Accepted 8th March 2012 DOI: 10.1039/c2cc30966a

A doubly 2,6-pyridylene-bridged porphyrin-perylene-porphyrin triad was synthesized *via* Suzuki-Miyaura cross coupling reaction, which captures a tetrakis(3-pyridyl)porphyrin guest in a 2 : 1 manner to form a supramolecular complex that undergoes photo-induced electron transfer.

Covalently linked multiporphyrin arrays have been extensively explored for their applications in photosynthetic models, sensors, molecular wires, nonlinear optical (NLO) materials, and so on.¹ Among these, we have recently explored iridium(i)-catalysed β -borylation reaction of porphyrins that provides β -borylated porphyrins selectively.² Using these β -borylated porphyrins we have developed various functional oligomeric porphyrin arrays including doubly β -to- β bridged porphyrin arrays **1a**, **1b**, **1c**-**d** and **1e** that have butadiyne-,³ 2,5-thienylene-,⁴ 2,6-pyridylene-⁵ and 5,15-porphyrinylene-spacer,⁶ respectively. Interestingly, these porphyrin arrays (**1a**-**d**) displayed large two-photon absorption (TPA) cross-section values owing to the effective π -conjugation between two porphyrins.⁷



We have also developed three-dimensional porphyrin ladders *via* combined reaction sequences of Ag(i)-promoted *meso-meso* coupling of porphyrins and Suzuki–Miyaura coupling.⁶ Furthermore, a similar strategy allowed us to synthesize doubly

2,6-pyridylene-bridged bent porphyrin belts⁷ and porphyrin barrels,⁸ which exhibited C_{60} -encapsulating ability. As an extension of this strategy, we envisioned that insertion of a large aromatic spacer between a 2-pyridyl spacer would lead to modulation of molecular shape of porphyrin arrays in curvature and ring diameter. With this idea, we tried the synthesis of a doubly 2,6-pyridylene-bridged porphyrin–perylene–porphyrin triad as a prototype. Perylene is an important and potential functional organic molecule that has been used in various applications such as field effect transistors (FETs), electrical conductors, and organic electroluminescence (EL) devices.⁹ Interestingly, we have found that a triad zinc(II) complex can bind a tetrakis(3-pyridyl)porphyrin guest in a 2 : 1 manner, in which the efficient electron transfer takes place from the zinc(II) porphyrin host to the free base porphyrin guest.

The synthetic scheme of porphyrin-perylene-porphyrin triad 5 is shown in Scheme 1. 2,5,8,11-Tetraborylperylene 2^{10} was coupled with an excess amount of 2,6-dibromopyridine under Suzuki-Miyaura cross-coupling conditions to afford 2,5,8,11-tetrakis(6-bromopyridin-2-yl)perylene 3 in 50% yield, which is practically insoluble in common organic solvents. 5,10,15-Triaryl-2,18-diborylporphyrin nickel(II) complex 4^2 was coupled with suspended 3. Separation by gel permeation chromatography (GPC) and silica gel chromatography afforded the triad 5Ni in 14% yield. The high-resolution electrosprayionization time-of-flight (HR-ESI-TOF) mass spectrum of the hybrid porphyrin dimer displayed the parent ion peak at m/z = 1209.0845 (calcd for C₁₆₄H₁₆₀N₁₂Ni₂ = 1209.0868 [M]²⁺). The ¹H-NMR spectrum of **5Ni** in CDCl₃ at room temperature was very broad but became sharpened at 60 °C to exhibit one singlet peak and two doublet peaks for the β -protons at 9.78, 8.92 and 8.69 ppm, a singlet peak at 10.71 ppm for the mesoprotons, two singlet peaks for the perylene protons at 8.69 and 8.41 ppm, and two doublets and one triplet for the pyridine protons at 8.09, 7.98 and 7.78 ppm. The structural assignment of perylene has been proved by a ¹H-¹H NOESY technique (see Fig. S4 in ESI[‡]).

The single crystals of **5Ni** suitable for X-ray diffraction analysis were grown by slow vapour diffusion of acetonitrile into its chloroform solution (Fig. 1).§ The crystallographic asymmetric unit of **5Ni** contains two molecules (triads A and B) of different structures, both of which took belt-like *syn* conformations. The centre-to-centre distances between the nickel ions are 16.68 and 18.98 Å for triads A and B, respectively. The porphyrin moieties are tilted by 50.17°, 59.43°, 68.73° and 74.91° relative to the perylene mean planes. These facts indicate certain

^a Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan.

E-mail: aratani@kuchem.kyoto-u.ac.jp, osuka@kuchem.kyoto-u.ac.jp; Fax: +81 75 753 3970; Tel: +81 75 753 4008

^b PRESTO, Japan Science and Technology Agency, Japan

[†] This article is part of the ChemComm 'Porphyrins and phthalocyanines' web themed issue.

[‡] Electronic supplementary information (ESI) available: Experimental details of the synthesis and spectroscopic analytical data of new compounds. CCDC 865051. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c2cc30966a



Scheme 1 Synthesis of porphyrin-perylene-porphyrin hybrid triads and host-guest complexation of **5Zn** with **mP**.

conformational flexibility for **5Ni**. In the solid state, triads A and B are alternatively stacked with their facing perylene parts closer at a distance of *ca*. $3.6 \text{ Å}.\ddagger$

A nickel complex was demetalated with H_2SO_4/TFA to give free base porphyrin dimer **5H**, which was then converted into the corresponding zinc(II) complex **5Zn** upon treatment with Zn(OAc)₂. The UV/vis absorption spectrum of **5Zn** in CH₂Cl₂ is shown in Fig. 2. The **5Zn** exhibits a blue-shifted Soret band at 429 nm and Q bands at 554 and 596 nm as compared with those of reference 5,10,15-triaryl-2,18-dipyridylporphyrin zinc(II) complex **6Zn** (Soret band; 433 nm, Q bands; 557 and 598 nm).‡ These phenomena suggest the weak but distinct excitonic interaction between the porphyrin and perylene units.¹¹

The bent structure of **5Ni** and its conformational flexibility inspired us to examine the supramolecular interaction with tetrakis(3-pyridyl)porphyrin (**mP**).^{12,13} Photometric titration experiments were conducted to determine the stoichiometry between **5Zn** and **mP** in CH₂Cl₂. Upon the addition of **mP** to the **5Zn** solution, the absorption intensities at 429 and 554 nm were found to decrease and that at 515 nm was seen to increase with isosbestic points at 437 and 562 nm. As shown in Fig. S13 (ESI‡), a plot of the absorbance at 429 nm *versus* [**mP**] has an inflection point at [**mP**]/[**5Zn**] = 0.5. The Job's plot clearly supported a 2 : 1 stoichiometry (Fig. S12, ESI‡). The ion peak of the complex was successfully detected by HR-ESI-MS in a



Fig. 1 X-Ray crystal structures of **5Ni**. (a) Side and (b) top view of triad A and (c) side and (d) top view of triad B. Substituents on phenyl groups are omitted for clarity. The ellipsoids are scaled to the 20% probability.



Fig. 2 UV/vis absorption spectrum of 5Zn in CH₂Cl₂. Inset: fluorescence spectra of 5Zn (black) and $5Zn_2$ -mP (red) excited at 437 nm in CH₂Cl₂.

positive mode at m/z = 2740.2639 for $[5Zn_2-mP + 2H]^{2+}$ (calcd for $[C_{368}H_{346}N_{32}Zn_4]^{2+} = 2740.2713$) (Fig. S11, ESI‡). These phenomena explain that **5Zn** and **mP** form a 2 : 1 complex. The ¹H NMR spectrum of the 2 : 1 mixture of **5Zn** and **mP** shows large complex-induced changes in chemical



Fig. 3 ¹H NMR spectra of (a) **5Zn**, (b) a mixture of **5Zn** and **mP** (2:1), and (c) **mP** in CDCl₃.

shifts for the signals corresponding to the pyridyl protons of mP as shown in Fig. 3. These results clearly demonstrate the guest-binding ability of 5Zn.

The steady-state fluorescence spectra of 5Zn and 5Zn₂-mP are shown in the inset of Fig. 2. Upon photoexcitation of 5Zn₂-mP at 437 nm that corresponded to selective excitation at the zinc(II) porphyrin moieties, the complex exhibited reduced fluorescence from the 5Zn part. Importantly, the fluorescence from the mP was not detected. These data suggested the intracomplex electron transfer from the photoexcited 5Zn to **mP**, while the excitation energy transfer is the most common process for zinc(II) free-base hybrid porphyrin pairs.¹⁴ By cyclic and differential pulse voltammetry methods in CH₂Cl₂, the first reduction potential of mP and the first oxidation potential of **6Zn** were measured to be -1.40 and +0.39 V, respectively, with respect to the Ag/AgClO4. The free energy calculated by the Rehm-Weller equation¹⁵ indicates that the electron transfer from ${}^{1}(5Zn)*$ to mP is exothermic by -0.46 eV, while the free energy change associated with the excitation energy transfer is -0.15 eV.[‡] The coordination interactions between the pyridyl groups with the Zn(II) centres render the zinc(II) porphyrins more electron-donating and the pyridyl-appended free base porphyrin more electron-accepting, which makes the intracomplex electron transfer more feasible.

In conclusion, we have synthesized the doubly 2,6-pyridylenebridged porphyrin–perylene–porphyrin triad by Suzuki–Miyaura cross-coupling reaction. The ¹H-NMR, mass, and UV/vis absorption spectra of this triad revealed the structural characterisation in solution, and the bent structure was observed in the solid state of **5Ni**. The supramolecular complex **5Zn₂–mP** leads to novel electro-photochemical properties. Investigation on the peryleneinserted porphyrin nanobarrel is currently in progress.

This work was partly supported by Grants-in-Aid for Scientific Research (No. 19205006 (A), 23685030 for Young Scientists (A) and 20108006 "pi-Space") from the Ministry of Education, Culture, Sports, Science and Technology, Japan, and by JST PRESTO program. The authors thank Prof. H. Maeda and Dr Y. Haketa (Ritsumeikan University) for MALDI-TOF MS measurement.

Notes and references

§ Crystallographic data for **5Ni**: $C_{164}H_{160}N_{12}Ni_2$, M = 2416.46, triclinic, space group $P\overline{1}$ (#2), a = 20.6104(4), b = 28.4361(5), c = 32.8677(6) Å, $\alpha = 113.2820(8)^\circ$, $\beta = 98.5654(9)^\circ$, $\gamma = 95.3241(9)^\circ$, V = 17251.5(6) Å³, T = 93(2) K, Z = 4, reflections measured 165003, 49 820 unique. The final R_1 was 0.0971 (> $2\sigma(I)$), and the final wR on F^2 was 0.2069 (all data), GOF = 1.040. CCDC 865051. The contributions to the scattering arising from the presence of the disordered solvents in the crystal were removed by use of the utility SQUEEZE in the PLATON software package.¹⁶

- N. Aratani and A. Osuka, in *Handbook of Porphyrin Science*, ed. K. M. Kadish, K. M. Smith and R. Guilard, World Scientific Publishing, Singapore, 2010, vol. 1, ch. 1, pp. 1–132 and references therein.
- 2 H. Hata, H. Shinokubo and A. Osuka, J. Am. Chem. Soc., 2005, 127, 8264.
- 3 I. Hisaki, S. Hiroto, K. S. Kim, S. B. Noh, D. Kim, H. Shinokubo and A. Osuka, *Angew. Chem.*, *Int. Ed.*, 2007, 46, 5125.
- 4 J. Song, S. Y. Jang, S. Yamaguchi, J. Sankar, S. Hiroto, N. Aratani, J.-Y. Shin, S. Easwaramoorthi, K. S. Kim, D. Kim, H. Shinokubo and A. Osuka, *Angew. Chem., Int. Ed.*, 2008, 47, 6004.
- 5 J. Song, P. Kim, N. Aratani, D. Kim, H. Shinokubo and A. Osuka, *Chem.-Eur. J.*, 2010, **16**, 3009; J. Song, N. Aratani, J. H. Heo, D. Kim, H. Shinokubo and A. Osuka, *J. Am. Chem. Soc.*, 2010, **132**, 11868.
- 6 J. Song, N. Aratani, P. Kim, D. Kim, H. Shinokubo and A. Osuka, *Angew. Chem., Int. Ed.*, 2010, **49**, 3617.
- 7 N. Aratani, D. Kim and A. Osuka, Chem.-Asian J., 2009, 4, 1172.
- 8 J. Song, N. Aratani, H. Shinokubo and A. Osuka, J. Am. Chem. Soc., 2010, 132, 16356.
- 9 S. Hayashi, T. T. Wang, S. Matsuoka and S. Saito, *Mol. Cryst. Liq. Cryst.*, 1986, **135**, 355; C. Li, M. Liu, N. G. Pschirer, M. Baumgarten and K. Müllen, *Chem. Rev.*, 2010, **110**, 6817; C. Wang, H. Dong, W. Hu, Y. Liu and D. Zhu, *Chem. Rev.*, 2012, **112**, DOI: 10.1021/cr100380z.
- 10 D. N. Coventry, A. S. Batsanov, A. E. Goeta, J. A. K. Howard, T. B. Marder and R. N. Perutz, *Chem. Commun.*, 2005, 2172.
- 11 Perfect overlapping of the absorption spectra and large deference of the molecular extinction coefficients for perylene and porphyrins make selective photo-excitation of the perylene difficult so that excitation energy and/or electron transfer from the perylene to the porphyrin was not clearly observed.
- 12 R. A. Haycock, A. Yartsev, U. Michelsen, V. Sundström and C. A. Hunter, Angew. Chem., Int. Ed., 2000, 39, 3616.
- 13 H. Shinmori, T. Kajiwara and A. Osuka, *Tetrahedron Lett.*, 2001, 42, 3617.
- 14 F. P. Schwarz, M. Gouterman, Z. Muljiani and D. H. Dolphin, *Bioinorg. Chem.*, 1972, 2, 1; J. A. Anton, P. A. Loach and Govindjee, *Photochem. Photobiol.*, 1978, 28, 235; R. L. Brookfield, H. Ellul, A. Harriman and G. Porter, *J. Chem. Soc., Faraday Trans.*, 1986, 82, 219; A. Osuka, K. Maruyama, I. Yamazaki and N. Tamai, *Chem. Phys. Lett.*, 1990, 165, 392.
- 15 D. Rehm and A. Weller, Isr. J. Chem., 1970, 8, 259.
- 16 Squeeze-Platon: A. L. Spek, *PLATON, A Multipurpose Crystallographic Tool*, Utrecht, The Netherlands, 2005; P. van der Sluis and A. L. Spek, *Acta Crystallogr., Sect. A*, 1990, **46**, 194.