View Article Online View Journal

Green Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: F. Jerome, F. Liu, M. Audemar, K. Vigier, D. Cartigny, J. Clacens, A. A.H. Padua, M. Costa Gomes and F. De Campo, *Green Chem.*, 2013, DOI: 10.1039/C3GC41495G.

Green Chemistry



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI[®]), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard **Terms & Conditions** and the **ethical guidelines** that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

RSCPublishing

www.rsc.org/greenchem Registered Charity Number 207890 Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxx

Selectivity enhancement in the aqueous acid-catalyzed conversion of glucose to 5-hydroxymethylfurfural induced by choline chloride

Fei Liu,^{*a*} Maïté Audemar,^{*a*} Karine De Oliveira Vigier,^{*a*} Damien Cartigny,^{*b*} Jean-Marc Clacens,^{*b*} Margarida F. Costa Gomes,^{*c*} Agilio A. H. Pádua,^{*c*} Floryan De Campo^{*b*} and François Jérôme*^{*a*}

s Received (in XXX, XXX) Xth XXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX DOI: 10.1039/b000000x

In this work we wish to show that choline chloride (ChCl), a cheap and safe quaternary ammonium salt industrially produced at thousand tons per year through a 100 % atom economy process, is capable of enhancing the selectivity of metal chlorides such as AlCl₃, FeCl₃ and CuCl₂ in the aqueous tandem

- ¹⁰ isomerization/dehydration of glucose to HMF. Under optimized conditions, 70% yield of HMF was obtained in a water/methylisobutylketone (MIBK) biphasic system which is a competitive yield to those traditionally obtained in imidazolium-based ionic liquids in the presence of hazardous chromium salts. By means of counter experiments, we show that the selectivity enhancement is optimal for a ChCl content of 50 wt% in water. At higher loading of ChCl, strong molecular interaction occurs between ChCl and HMF
- ¹⁵ making the extraction of HMF from the aqueous phase difficult with MIBK, thus enhancing side reactions of HMF with water and hexoses to unwanted products. Interestingly, this process can be transposed to the direct conversion of cellulose to HMF which is an even more challenging reaction. In this case, we show that combination of FeCl₃ with AlCl₃ allowed cellulose to be converted to HMF with 49% yield in a one pot reaction. From the view point of sustainable chemistry, this work gathers
- ²⁰ noticeable advantages such as use of 1) water as a solvent, 2) ChCl as a cheap and safe additive, 3) cheap and naturally abundant metal (Al, Fe, Cu) and 4) renewable raw materials

Introduction

Published on 03 September 2013. Downloaded by Brown University on 06/09/2013 07:28:37

The development of Chemistry Safety Rules, the proposition of the twelve principles of green chemistry in 1998 by P. Anastas ²⁵ and J. Warner as well as the price and scarcity of oils are now key elements in establishing sustainable reactions/processes.¹ Within this framework, use of renewably-sourced raw materials for the production of innovative products has become a top priority in chemical industry. In this context, much effort has been paid to

- ³⁰ the synthesis 5-hydroxymethylfurfural (HMF) as a chemical platform from which solvents, monomers, fuel additives, among others can be then produced.² HMF is produced by an acid-catalyzed triple dehydration of hexoses. Although a plethora of works have been dedicated to the acid-catalyzed dehydration of
- ³⁵ fructose to HMF, the direct use of glucose (directly available from lignocellulosic biomass) is much less widespread mainly because it requires tackling the isomerization of glucose to fructose prior to the dehydration step to HMF. Isomerization of glucose to fructose can be catalyzed by Lewis acids such as for
- ⁴⁰ instance tin-containing zeolites but yields of fructose remained however rather low.³ Basic catalysts were also proposed but hexoses are relatively instable in alkaline conditions affording fructose in low yield.⁴ Up to now, industrial processes involve enzymes which are capable of producing up to 50% yield of
- 45 fructose from glucose although this process has also clearly some

drawbacks (enzyme deactivation, large quantity of enzymes and low activity of enzymes).⁵ Note that enzymatically produced fructose can be then dehydrated to HMF in a sequential process (fermentation followed by acid catalysis).⁶ In 2007, Zhang and ⁵⁰ co-workers reported that chromium chloride can efficiently catalyze the tandem isomerization/dehydration of glucose to HMF with an unprecedented yield of 70% in ionic liquids.⁷ Recently, variants of this process have been proposed especially from cellulose and lignocellulosic biomass. Although from an ⁵⁵ environmental point of view the sustainability of such process is debatable, this work is however a bench-top methodology that allows a better understanding of the different elementary steps involved in the catalytic isomerization of glucose to fructose.

Choline chloride (ChCl), a cheap, biodegradable and safe quaternary ammonium salt industrially produced at thousand tons per year through a 100 % atom economy process, is known to be capable of producing a Deep Eutectic Solvent (DES) when mixed with metal chlorides or hexoses such as glucose or fructose.⁸ This particular property of ChCl suggests that this moiety has the unique ability to stabilize or destabilize hydrogen bond donors such as hexoses or HMF but also metal-derived catalytic species. In this context, this work aims at identifying fundamental issues on the way how ChCl may impact the catalytic cycles involved in the tandem isomerisation/dehydration of glucose to HMF (*i.e.* 70 role as a ligand, stabilization of HMF, *etc...*). By comparing data from the current literature with our experimental results, we wish

ARTICLE TYPE

8

2-2/2

 $2-2/2^{\circ}$

to show here that use of ChCl as an additive in water is a great tool for assisting Lewis acid catalysts in the direct conversion of glucose to HMF. Under optimized conditions, HMF was produced with up to 70% yield in water which is a very ⁵ competitive yield to those obtained in traditional imidazoliumbased ionic liquids with chromium-based catalysts. Beside the clear economic and environmental advantages stemming from the use of ChCl (< 2€/Kg, biocompatible) and water, we wish to demonstrate here that assistance of ChCl also allows using safe ¹⁰ metal chlorides such as AlCl₃, FeCl₃ or CuCl₂.

Results and Discussion

Catalytic experiments

In a first set of experiments, glucose was mixed with ChCl (Glucose/ChCl weight ratio = 1/1, 2/1, 1/5). As described earlier, ¹⁵ when the solution was heated at 100°C a transparent and viscous liquid DES was formed. Next, 3 mol% of Cu(NTf₂)₂ were added to the solution and heated first at 180°C under microwave (250W). The choice for this Lewis acid was motivated by our recent results collected with metal triflates in the catalytic ²⁰ etherification of glycerol.⁹ Note that catalyst screening is described later in the manuscript. Unfortunately, under these conditions, HMF was only produced as a trace presumably due to the low reactivity of glucose when engaged in the formation of a DES as similarly observed in the current literature by Kazlauskas

- ²⁵ and co-workers in a ChCl/glycerol eutectic mixture (Table 1, entry 1).¹⁰ With the aim of lowering the interaction between glucose and ChCl, water was progressively added to the reaction medium. In all experiments, the glucose content was kept constant to 10 wt%. As water was added, we were pleased to see ³⁰ that formation of HMF occurred and yield of HMF reached 31%
- at a ChCl/H₂O weight ratio of 1 (*i.e.* 7.1 M) which correspond to an HMF selectivity of 42% (Table 1, entry 3). Upon prolonged reaction time, the yield of HMF slightly dropped due to its side degradation, and condensation. Note that, under these conditions, ³⁵ levulinic acid was not detected but formation of soluble black
- materials so-called humins was also observed. When the amount of water was further increased (ChCl/H₂O weight ratio of 1/2), the yield of HMF was also decreased suggesting that an optimal ChCl/H₂O ratio need to be respected (Table 1, entry 4). More 40 information about the role of the ChCl/water ratio is provided
- later in the manuscript. Next, in an attempt to increase the yield of HMF.

methylisobutylketone (MIBK) was added to the system in order to "continuously" extract HMF from the catalytic phase and to 45 limit its acid-catalyzed side reaction with water or hexoses to

- unwanted products. As expected, when 10 wt% of glucose were heated at 180° C in a ChCl-H₂O/MIBK (ratio of 1-1/1) biphasic system for 15 min, the selectivity to HMF was improved and HMF was produced with 46% yield (Table 1, entry 5). Note that
- ⁵⁰ upon prolonged reaction time, the yield of HMF started to decrease and, in this case, we clearly observed the formation of levulinic acid (9% yield after 30 min, Table 1, entry 6). A decrease of the reaction temperature from 180°C to 165°C obviously lowered the reaction rate but unfortunately had also a
- 55 negative impact on the reaction selectivity. At 165°C, the maximum yield of HMF was 30% which correspond to an HMF

Table 1 Acid-catalyzed conversion of glucose to HMF in the presence of $Cu(NTf_2)_2$



^{*a*} molar yield; ^{*b*} along with 9% yield of levulinic acid; ^{*c*} from a glucose ⁶⁰ content of 20 wt%

7

7

89

83

29

38

33

46

200

180

selectivity of 43% vs 64% at 180°C (Table 1, entry 7). Similarly, an increase of the reaction temperature from 180°C to 200°C (Table 1, entry 8) also lowered the HMF selectivity (from 64% to 33%) mainly due to the dominant formation of soluble and insoluble black materials as well as other side products. Hence, one may conclude that 180°C is the optimal temperature when working with Cu(NTf₂)₂ as a catalyst. In a similar trend than previous works,² an increase of the glucose content from 10 wt% to 20 wt% also lowered the selectivity of the reaction and HMF 70 was produced with 38% yield (vs 46% at a glucose content of 10

was produced with 38% yield (vs 46% at a glucose content of wt%, Table 1, entries 5, 9).

Although promising yields of HMF have been obtained in the ChCl-H₂O/MIBK system in the presence of Cu(NTf₂)₂, we then screened other Lewis acid catalysts in order to select the most ⁷⁵ active ones. Otherwise noted, all conditions employed for catalytic experiments are 10 wt% of glucose, a ChCl-H₂O/MIBK (ratio of 1-1/1) biphasic system and 3 mol% of catalyst. Reaction temperature was fixed to 165°C (*vs* 180°C with Cu(NTf₂)₂) in order to reduce the reaction rate and to better select the most ⁸⁰ active catalysts. To compare the activity of all catalysts, the reaction time was first fixed to 15 min in all cases. Results are summarized in Table 2.

Among tested metal triflates and metal triflimidates, Al is the most active one affording a glucose conversion of 99% followed ⁸⁵ by Fe and Cu (Table 2, entries 1-7). In the case of metal

- chlorides, large differences of activity were observed. As previously described in imidazolium-based ionic liquids⁷ or in a water/NaCl mixture,¹¹ AlCl₃, CrCl₂ and CrCl₃ were the most active ones (Table 2, entries 8, 10, 11). It is noteworthy that safe ⁹⁰ metal chlorides such as FeCl₃ and CuCl₂ also exhibited an acceptable activity leading to a glucose conversion of 58% and
- acceptable activity leading to a glucose conversion of 58% and 56%, respectively (Table 2, entries 12, 13). Activity of metal chlorides can be classified as follow AlCl₃, CrCl₂, CrCl₃>FeCl₃, CuCl₂>MgCl₂, FeCl₂>MnCl₂, BiCl₃>ZnCl₂. The great catalytic ⁹⁵ activity of AlCl₃, FeCl₃ and CuCl₂ in an aqueous solution of ChCl is attractive mainly because of their low price and natural abundance. Hence, we selected these metal chlorides and we tried to optimize the yield of HMF. As the yield of HMF passes

Published on 03 September 2013. Downloaded by Brown University on 06/09/2013 07:28:37



^{*a*} nd= not determined; ^{*b*} result collected at 180°C, ^{*c*} result collected at150°C, ^{*d*} BHC = betain hydrochloride, pH=1

through a maximum during the catalytic process, we first ⁵ monitored the HMF yield *versus* time. At 165°C, kinetic studies confirmed the high activity of AlCl₃ and we found that HMF was produced with a maximum yield of 67% after only 3 min of reaction which corresponds to a glucose conversion of 89% and an HMF selectivity of 75% (Table 2, entry 8). AlCl₃ is so active ¹⁰ that we were able to decrease the temperature to 150°C (entry 9). At this temperature, a slight increase of the HMF yield to 70% was observed. Note that this yield value is similar to that previously reported by Zhang in the presence of 6 mol% of CrCl₂ in 1-butyl-3-methylimidazolium chloride (reference work in this ¹⁵ field of chemistry),⁷ thus showing the potential of such catalytic

- system for the sustainable production of HMF. In the case of FeCl₃, a maximum yield of HMF of 43% (entry 12) was obtained after 25 min of reaction, which is quite close to the yields of HMF collected with chromium-based catalysts in our conditions (400) and 570 with the property of C Cl and C Cl
- $_{20}$ (49% and 57% yield in the presence of CrCl₂ and CrCl₃, respectively, entries10-11). Remarkably, whereas Bronsted acid usually provides very low yields of HMF from glucose (also observed here in the absence of ChCl), we were pleased to see here that, in the presence of ChCl, HMF was produced with 34%
- ²⁵ and 22% yield with BHC (betaïn hydrochloride) and HCl, respectively (Table 2, entries 19-20), reinforcing the key role played by ChCl in the conversion of glucose to HMF.

Next, we tried to rationalize the greater selectivity of AlCl₃ as compared to that of FeCl₃ in our conditions. Interestingly, we ³⁰ found that yields of HMF were comparable between FeCl₃ and AlCl₃ (43%) up to 70% conversion of glucose (see supporting information, Fig S1, S2 and S3). However, upon prolonged reaction time significant difference was observed between both metal chlorides. In the presence of AlCl₃ the yield of HMF ³⁵ gradually increased up to 70% (at 90% conversion). Reversely, in the presence of FeCl₃, the yield of HMF dramatically dropped at conversion higher than 70%. This surprising difference of behaviour between both metal chlorides prompted us to more deeply investigate the stability of catalytically active species in ⁴⁰ our system.

Recent works have shown that, in the presence of water, metal chlorides such as CrCl₂ and AlCl₃ are in equilibrium with metal hydroxide species M(OH)_x(H₂O)_v and HCl.¹² Metal hydroxide is suspected to be the catalytically active species in the 45 isomerization of glucose to fructose while released HCl would catalyze the dehydration of fructose to HMF. Assuming that metal hydroxide species are suspected to be the catalytically active sites in the isomerization of glucose to fructose, we first heated FeCl₃ and AlCl₃ in neat water (30 min at 165°C under 50 microwaves) prior to add ChCl and glucose. The pH of the solution was measured before and after heating of the corresponding metal chloride in neat water (Table 3). As expected, in both cases, an important drop of the pH was observed after heating in water which is in accordance with a 55 partial hydrolysis of metal chlorides accompanied by the release of HCl in the medium. Note that measured pH were in the same range between AlCl₃ and FeCl₃ suggesting that the higher activity of AlCl₃ cannot be ascribed to a greater acidity. However, noticeable visual differences were observed between pre-heated 60 aqueous solution of AlCl3 and FeCl3. Whereas the solution remained homogeneous in the case of AlCl₃, we clearly observed the formation of a rust-coloured precipitate in the case of FeCl₃ which was attributed to the formation of hydrated iron oxide as supported by RAMAN and XRD analyses (see supporting 65 information, Fig S4 and S5) (Fig. 1). Hence, we suspected here

Table 3 Stability of catalytic species^a

			MCl _x (M=Al	, Fe)		
HOHOHOHOHOHOHOOHOOHOOH						
Without pre-heating				With pre-heating		
Catalyst	pH ^b	Conv. glucose (%)	HMF molar yield $(\%)^a$	pH ^c	Conv. glucose (%)	HMF mola yield (%) ^a
AlCl ₃	3.54	73	44	1.86	77	42
FeCl ₃	2.10	72	43	1.18	72	22

^a 3 mol % of catalyst, 150°C, reaction time was adjusted in compared to compare data at same conversion; ^b initial pH of the solution after addition of the catalyst; ^c initial pH of the solution after pre-heating metal ⁷⁰ chlorides in water for 30 min at 165°C

This journal is © The Royal Society of Chemistry [year]



Fig. 1 Pictures of the solution of FeCl₃ and AlCl₃ (a) before and (b) after heating in water at $165^{\circ}C$

¹⁰ that the lower efficiency of FeCl₃ might be ascribed to the precipitation of iron oxide species during the reaction resulting in a dramatic decrease of the glucose/fructose isomerization rate to the benefit of side reactions catalyzed by released HCl. To further support this hypothesis, ChCl and glucose were added to the pre-¹⁵ heated aqueous solution of AlCl₃ and FeCl₃. Whereas a similar yield of HMF was recovered using a pre-heated aqueous solution of AlCl₃, the selectivity to HMF was dramatically decreased in the case of FeCl₃ further demonstrating that precipitation of iron oxides species has a detrimental effect on the catalyst activity and ²⁰ selectivity. Note that when ChCl was added during the preheating step of metal chlorides in water, a rust-coloured precipitate was also observed from FeCl₃ while all species remained soluble with AlCl₃ suggesting that ChCl did not prevent the deactivation of FeCl₃.

25 Role of ChCl

5

To collect more information on the synergistic role of ChCl and water in our system, the reaction was carried out without ChCl either in neat 1-butyl-3-imidazolium chloride or in neat water. All reactions were performed at 165°C (250W under microwaves)³⁰ with a glucose loading of 10 wt% and a H₂O/MIBK ratio of 2/2. Although AlCl₃ is more stable than FeCl₃ in our conditions, FeCl₃ has been selected in the following experiments mainly because this metal chloride is generally considered as a poorly active catalyst in such reaction. Hence, FeCl₃ appears as a good ³⁵ candidate to clearly highlight the promoting effect of ChCl on the catalysts selectivity. In the absence of ChCl, HMF was produced with a very low yield of 9% and 1% in [BMIM]Cl and in neat water, respectively, and an important formation of black materials was observed (Table 4, entries 2, 3). Similarly, when the catalytic ⁴⁰ reaction was conducted without water in a mixture ChCl/DMSO,

- HMF was also produced with only 1% yield (Table 4, entry 4) suggesting that combination of ChCl with H₂O is one of the key for explaining the selectivity improvement observed with FeCl₃. ChCl was also replaced by other water soluble quaternary
- ⁴⁵ ammonium salts to confirm its effect on the reaction selectivity. In such case, the amount of salt was adjusted to keep constant the mol% of salt introduced in water. Reactions were all stopped and compared at a conversion range of 50-60% to evaluate the role of choline on the HMF selectivity. When ChCl was replaced by
- ⁵⁰ dimethyldibutylammonium chloride, methyltributylammonium chloride or NaCl, the yield of HMF dramatically dropped to 18%,12% and 15%, respectively (vs 33% with ChCl, Table 4 entry 1). Additionally, when ChCl was replaced by ethanol, the

Table 4 Effect of various salts in the acid-catalyzed dehydration of 55 glucose to HMF^a





selectivity of the reaction was shifted towards the formation of fructose (18 % yield at 63 % conversion) in agreement with the recent work of Riisager.¹³ From these results, one may also conclude that both the -OH and ammonium groups of ChCl play a crucial role in the reaction mechanism.

To get more insight on the role of ChCl in such catalytic reaction, we carried out extra experiments. ChCl is capable of producing eutectic mixtures with hydrogen bond donors such glucose or 65 fructose.¹⁴ In a first approximation, one may suspect that, in water, ChCl interact stronger with fructose than with glucose thus facilitating the isomerization of glucose to fructose. By means of calculations (see ESI), we found that the total energy of the solutions, including terms due to the solutes (sugars), the solvent 70 (ChCl/H₂O) and also solute-solvent interactions, were similar between the glucose/ChCl/H2O and fructose/ChCl/H2O systems. Additionally, decompositions of the energy into different terms, either by nature (electrostatic, van der Waals, intramolecular) or by component (solute, solvent), gave very close values for the 75 two systems. Hence, from these results, one may conclude that the two systems have a similar energy and thus rule out the possible "assisting role" of ChCl in the glucose/fructose isomerization step.

Considering that the yield of HMF is optimal at a ChCl/H₂O ratio ⁸⁰ of 1/1, we next focused our attention on the partition coefficient of HMF in these biphasic systems. Dumesic and co-workers have Published on 03 September 2013. Downloaded by Brown University on 06/09/2013 07:28:37

previously showed that a "continuous" extraction of HMF from the aqueous phase allowed its side reactions with water or hexoses (*i.e.* rehydration to levulinic and formic acids, condensation to humins, *etc...*) to be limited.¹⁵ To this end, we s reproduced the composition of the reaction media obtained at the end of the reaction (see composition entry 9 of Table 2). In a

- typical experiments, HMF (a quantity equivalent to 70% yield) was heated in a biphasic water/MIBK system (each phase = 10 mL) at 100°C for 10 min. The MIBK and aqueous phase were to then separated and analyzed by HPLC. The partition coefficient
- was defined as the mass ratio between the amount of HMF detected in the MIBK and water phases, respectively. Then, same experiments were reproduced by progressively substituting water by ChCl. Result of this study is presented on Fig. 2.
- ¹⁵ Interestingly, up to a ChCl content of nearly 50 wt%, the partition coefficient did not significantly vary and remained around 1.2 (*i.e.* about 53 wt% of HMF was extracted with MIBK). Reversely, at a higher ChCl content, the partition coefficient significantly and continuously decreased up to 0.6 (*i.e.* about 38
- 20 wt% of HMF in the MIBK phase) at a ChCl content of 80 wt%. This result suggests that, at a ChCl content higher than 50 wt%, HMF is strongly retained in the aqueous phase, presumably through hydrogen bond interaction. As a consequence, in this scenario, side reactions of HMF with water (and hexoses) should 25 be more favourable and should result in a drop of the reaction





Fig. 2 Partition coefficient of HMF in biphasic ChCl-H₂O/MIBK systems 40 and HMF yield (see conditions Table 2, entry 9) as a function of the ChCl concentration

Although consideration of the partition coefficients allowed explaining the behaviour of our system at high loading of ChCl, it does not explain why the selectivity to HMF was increased when ⁴⁵ the ChCl content was concomitantly raised from 0 to 50 wt% (in this range of concentration, the partition coefficients are similar, Fig. 2). Recently, K. Binnemans and R. Van Deun have

- investigated the speciation of copper chlorides in ChCl/water mixtures.¹⁶ This work clearly showed by EXAFS, electrochemical
- ⁵⁰ measurements and crystallography that the environment of the cupric ion is modified according to the amount of ChCl in water. At a water concentration of 39 wt% in ChCl, ChCl is coordinated on copper while at a concentration higher than 95%, the cupric ion is fully hydrated. At an intermediate water/ChCl composition,
- 55 mixed chloro-aquo complexes are dominant species. Inspired by data from the current literature and by our experimental results, we propose hereafter a plausible explanation for rationalizing our

results. At a ChCl content lower than 50 wt%, metal hydroxide species are produced in water which is accompanied by the ⁶⁰ release of HCl. The coordination sphere of these metal hydroxide species may be modified by increasing the amount of ChCl in the reaction media as previously shown by K. Binnemans and R. Van Deun. Note that the possible role of ChCl as a ligand is consistent with our above-described experiments summarized in Table 4. In ⁶⁵ particular, by increasing the concentration of ChCl in water, the nature and concentration of catalytic species in the reaction media might change which would be consistent with a change of the selectivity to HMF. Concomitantly, the possible extraction of

- HMF from the aqueous phase with MIBK allowed limiting side ⁷⁰ reactions of HMF with water and hexoses. In this context, best selectivity to HMF was obtained in a biphasic aqueous solution of ChCl (50 wt%)/MIBK. At a ChCl content higher than 50 wt%, strong molecular interaction between ChCl and produced HMF became dominant and HMF was more strongly retained in the ⁷⁵ aqueous phase. As a consequence, HMF is subjected to side reactions with water or hexoses leading to a drop of the reaction selectivity. Note that this hypothesis is supported by our recent findings that suggested the formation of a possible DES between HMF and ChCl.¹⁷ Hence, one may conclude that 50 wt% of ChCl ⁸⁰ in water is an acceptable compromise between the catalyst
- activity/selectivity and the extraction of HMF from the aqueous phase.

One should comment that when all the above-described reactions were performed in conventional heating, HMF was produced 85 with a much lower selectivity (in this case insoluble black materials were produced in a higher amount) and that whatever the reaction conditions (temperature, time, fructose and catalyst loading, etc...). In conventional heating, it has been previously demonstrated that the isomerization rate of glucose to fructose ⁹⁰ was nearly 4 times lower than the dehydration rate of fructose to HMF.¹⁸ Because of the relative instability of HMF in aqueous acidic conditions, it is clear that the production of HMF is in competition with its own degradation, rehydration or condensation. Although we are fully convinced that microwaves 95 also contributed to a better control of the reaction selectivity, all attempts to identify which elementary step was affected by microwaves in a greater extent unfortunately failed mainly due to the complexity of the system. Note that during the course of our investigations, similar observation has been reported by Rinaldi, 100 Dumesic and co-workers.¹⁹

Recyclability of the system

Having all these results in hand, we then evaluated the recyclability of the ChCl-H₂O phase containing the acid catalyst. Recycling experiments were performed at 150°C for 15 min in ¹⁰⁵ the presence of 3 mol % of AlCl₃. In all cases, MIBK was added to continuously extract HMF from the reaction media (ChCl-H₂O/MIBK ratio = 2-2/2). After the first catalytic run, the reaction media was cooled down to room temperature and decanted. The MIBK phase was separated and the aqueous phase ¹¹⁰ containing ChCl and AlCl₃ was washed again with MIBK to ensure the complete removal of HMF from the catalytic phase. Then, glucose was directly added to the catalytic phase and heated for another 15 min at 150°C. One can see from Fig. 3 that the yield of HMF remained constant during at least three ¹¹⁵ consecutive recycling. After the third recycling, the yield of HMF

49



Fig. 3 Recycling experiments (150°C, 15 min, 3 mol% of AlCl₃, ChCl- $H_2O/MIBK$ ratio = 2-2/2.

¹⁵ however continuously dropped mainly due to the accumulation of impurities in the catalytic phase. After the fifth run, we attempted to recover ChCl to reuse it. To this end, the reaction media was diluted with 20 mL of water resulting in the precipitation of black materials which were filtered off. Water was then evaporated and ²⁰ the recovered residue was dried overnight at 150°C prior to be extracted with ethanol. After removal of ethanol under reduced pressure, ChCl was recovered (> 90 wt%) with a sufficient purity to be reused (see Fig. S6 for copies of NMR spectra). Indeed, after extraction and reuse of ChCl, a yield of HMF of 60% was ²⁵ recovered demonstrating the relative stability of ChCl in our working conditions.

Transposition to cellulose

Encouraged by all these results, we then attempted the direct conversion of cellulose to HMF which is an even more ³⁰ challenging reaction. This cascade reaction requires first a hydrolysis of cellulose to glucose followed by the isomerization/dehydration of glucose to HMF (Fig. 4). Reactions were first performed at 200°C (under microwave irradiations) in the presence of 4 wt% of FeCl₃ or AlCl₃ for 5 min. ³⁵ Microcrystalline cellulose (PH AVICEL 101) was used in all experiments. Interestingly, in the absence of ChCl, cellulose was selectively converted to glucose, HMF and levulinic acid in 2%, 8% and 3% yield respectively (along with many other side ⁴⁰ products) (Table 5, entries 1, 2). This difference of selectivity

between FeCl₃ and AlCl₃ is consistent with our previous observations. During the reaction with FeCl₃, iron hydroxide



Fig. 4 Catalytic conversion of cellulose to HMF in an aqueous solution of ChCl containing FeCl₃ and AlCl₃

Table 5 Catalytic conversion of cellulose to HMF in an aqueous solution of ChCl



⁶⁰ ^a microcrystalline cellulose PH AVICEL 101; ^b along with 2% of glucose and 3% of levulinic acid; ^c similar yields was obtained with CrCl₂ (17% yield); ^d reaction was performed in two steps. First 200°C for 5 min in neat water containing 4 wt% (vs cellulose) of FeCl₃ followed by addition of 3 mol% of AlCl₃ (vs in-situ produced glucose) and 50 wt% of ChCl

ves

FeCl₃/AlCl₃^d

Low

65 (heating at 135°C for 90 min); ^e cellulose with low cristallinity index was obtained by a ball-milling or a dissolution/regeneration from 1-butyl-3imidazolium chloride. Note that similar results were obtained whatever the pre-treatment.

precipitated (under the form of hydrated iron oxides) from the reaction media leading to an important release of HCl in the solution, a Bronsted acid known to efficiently catalyze the depolymerization of cellulose to glucose. Reversely, from AlCl₃, all species (HCl and aluminium hydroxides) remained soluble and, without ChCl, the reaction suffered from an important lack of selectivity.

Next, ChCl was added to a solution containing FeCl₃ and cellulose. In this case, the selectivity of the reaction was changed and glucose was produced with 26% yield along with 17% yield of HMF, further demonstrating the beneficial effect of ChCl for ⁸⁰ the production of HMF (Table 5, entry 3). Note that the reaction selectivity was more difficult to control from cellulose than from glucose. In such conditions, HMF was indeed also readily converted to levulinic and formic acids. Similar trend was also observed using AlCl₃ and CrCl₂ (14% and 17% yield of HMF, ⁸⁵ respectively, entry 4).

To increase the selectivity to HMF, cellulose was first heated in the presence of FeCl₃ (hydrolysis of cellulose to glucose) prior to add AlCl₃ (conversion of glucose to HMF). As observed above, glucose was produced with 40% yield in the presence of FeCl₃ 90 only. Then AlCl₃ was directly added to the solution (without intermediate purification) and the resulting mixture was heated at 135°C for 90 min. Interestingly, in this case, the in-situ produced glucose was converted to HMF with similar selectivity (68%) than that described above from glucose leading to HMF with a 95 yield of 27% (Table 5, entry 5). Please note that combination of HCl and AlCl₃ did not provide similar results suggesting that FeCl₃ still participate to some extent to the conversion of glucose to HMF. To further increase the yield of HMF, microcrystalline cellulose was subjected to a pre-treatment process prior to its 100 catalytic conversion to HMF. To this end the crystallinity index of microcrystalline cellulose was decreased either by dissolution/regeneration in 1-butyl-3-methylimidazolium chloride ([BMIM]Cl) or by ball-milling. As expected, after these two pre-

10

Published on 03 September 2013. Downloaded by Brown University on 06/09/2013 07:28:37

treatments, cellulose was found to be much less recalcitrant. During the first step in the presence of FeCl₃, cellulose was deconstructed to glucose with a yield of 70% *versus* 40% without pre-treatment (Table 5, entry 6). Then, after addition of AlCl₃, *ins situ* produced glucose was readily converted to HMF with 49%

- yield (*i.e.* 70% selectivity), (Table 5, entry 7). Although the large energy input required by the pre-treatment of cellulose by ballmilling or by dissolution in ionic liquids nowadays represents a major drawback, the coupling of our methodology with the recent
- ¹⁰ work reported by F. Schüth²⁰ in the field of mechanocatalytic deconstruction of cellulose to water soluble cello-oligomers may open a more eco-efficient route for the production of HMF from cellulose.

Conclusion

- ¹⁵ We report here that ChCl can be considered as a safe and cheap additive capable of assisting the catalytic conversion of glucose to HMF in water. Under optimized conditions, HMF was produced with 70% yield which is a competitive yield to those traditionally obtained in imidazolium-based ionic liquids in the
- ²⁰ presence of hazardous chromium salts. A screening of various Lewis acids revealed that ChCl has the ability to improve the selectivity of various safe and abundant metal chlorides such as AlCl₃, FeCl₃ and CuCl₂, thus increasing the sustainability of this process. It should be noted that the promoting effect of ChCl on
- 25 the HMF selectivity is however closely depending on the ChCl content. Although the exact elucidation of the reaction mechanism involved in the presence of ChCl is difficult to be determined with accuracy at this stage, we suggest that at a ChCl lower than 50 wt% in water, the selectivity is governed by 1) the
- ³⁰ coordinating ability of ChCl that presumably modifies the nature and the concentration of catalytic species in water, and 2) the extraction of HMF from the aqueous phase, thus limiting side reactions of HMF with water or hexoses. At a ChCl content higher than 50 wt%, strong molecular interaction between HMF
- ³⁵ and ChCl became dominant making extraction of HMF from the aqueous phase more difficult which resulted in a decrease of the reaction selectivity.

The scientific strategy of this work can be transposed to the direct production of HMF from cellulose which is a much more

 $_{40}$ challenging reaction. In this case, combination of FeCl₃ and AlCl₃ allowed cellulose to be converted to HMF with 49% yield in a one pot process.

From the view point of sustainable chemistry, this work gathers noticeable advantages such as use of 1) water as a solvent, 2)

⁴⁵ ChCl as a cheap and safe additive, 3) cheap and naturally abundant metal (Al, Fe, Cu) and 4) renewable raw materials. We do believe that this work, together with the recent progress reported in the field of process intensification (including biphasic catalysis) will definitely contribute to increase again the eco-⁵⁰ efficiency and sustainability of this work.

Experimental Section

Chemicals: Fructose, glucose, choline chloride and methyisobutylketone (MIBK) were purchased from Sigma Aldrich as well as anhydrous and hydrated Lewis acids. All

55 chemicals were used as received without further purification. Microcrystalline cellulose PH AVICEL 105 was purchased to FMC BioPolymer (degree of polymerization about 250, glucose content > 99%, water content = 5wt %, particle size < 40μ m). Distilled water was used as in all catalytic reactions.

- ⁶⁰ **Catalytic experiments**: All reactions were performed in an automated multimode microwave oven MicroSynth from Milestone s.r.l. (Italy). This oven operates at 2.45 GHz and is temperature controlled by a fiberoptic sensor. Temperature was monitored by inserting the calibrated fiberoptic sensor in the ⁶⁵ reaction mixture while stirring properly. The maximum power input could be adjusted between 0 and 1000 Watt. Under microwave heating, the internal temperature of the reaction vessel reached the reaction set-points in 20 min, and was maintained at \pm 5°C using a power compensation feedback control.
- ⁷⁰ Catalytic conversion of glucose to HMF: In a typical experiment, 0.4 g of glucose was dissolved in 2 g of water containing 2 g of ChCl and 3 mol% of a Lewis acid. Then 2g of MIBK were added and the biphasic solution was heated under microwave (250W) at the desired temperature for 1-60 min (see ⁷⁵ Tables for conditions).
- **Recycling procedure**: At the end of the reaction (conditions: 10 wt% of glucose, 3 mol% of AlCl₃, T =150°C, 15 min, 6g of ChCl-H₂O/MIBK (ratio 2-2/2)), the reaction was cooled down to room temperature and the MIBK phase containing the HMF was separated by decantation. Next, the catalytic phase was washed two times with 2.5 mL of MIBK to remove the residual HMF. Then, glucose was added to the catalytic phase and the reaction was heated again at 150°C for 15 min.
- **Catalytic conversion of cellulose to HMF**: 200 mg of cellulose ⁸⁵ was first suspended in 2 mL of water containing 18 mg of FeCl₃. The solution was then heated under microwave (250W) for 5 min at 200°C. Then the reaction was cooled down and 16 mg of AlCl₃, 2 g of ChCl and 2 mL of MIBK was directly added to the solution. The mixture was then heated for 30-180 min at 120-⁹⁰ 135°C under microwave (250W).

Analytical: Conversion of glucose was calculated by external calibration using a Shimadu Prominence LC equipped with a degasser DGU-20A3, a pump system (5LC-20AQ), a thermostated autosamplor SIL-20AC and an oven CTO-20AC
⁹⁵ maintained at 40°C. Glucose was separated on a Shodex KS-802 column, using water as an eluent, and quantified by a RID. Yields of HMF were determined by external calibration at 25°C using an HPLC equipped with a Nucleosil 100–5C18 column (250 x 4.6 mm), a Shimadzu LC-20AT pump, a Shimadzu RID-10A
¹⁰⁰ detector, and an acetonitrile/water (10:90) mixture as mobile phase (0.8 mL.min⁻¹).

Acknowledgements

Authors are grateful to the CNRS, the French Ministry of Research and SOLVAY for their financial supports.

105 Notes and references

^a Institut de Chimie des Milieux et Matériaux de Poitiers, CNRS/Université de Poitiers/ENSIP, 1 rue Marcel Doré, 86022 Poitiers/FRANCE Fax: +33 5 49 45 33 49; Tel: +33 5 49 45 40 52; E-mail: francois.jerome@univ-poitiers.fr

¹¹⁰ ^b Eco-Efficient Products and Processes Laboratory, Unité Mixte de Recherche UMI 3464 CNRS/RHODIA, 3966 Jin Du Road, Shanghai 201108, China, Fax: +86 21 54425911; Fax, Tel : +86 21 24089391, Email: floryan.decampo@ap.rhodia.com 75

80

85

90

95

100

105

110

120

125

130

- c^a Institut de Chimie de Clermont-Ferrand, CNRS/Université Blaise Pascal Clermont-Ferrand, 24 avenue des Landais, BP 80026, 63171 Aubiére, France. E-mail : agilio.padua@univ-bpclermont.fr
- ‡ Footnotes should appear here. These might include comments relevant
 s to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.
- P. T. Anastas, J. C. Warner, in Green Chemistry-Theory and Practice, Oxford University Press: New York, 1998.
- M. E Zakrzewska, E. Bogel-Lukasik, R. Bogel-Lukasik, *Chem. Rev.*,
 2001, **111** (2), 397-417; b) J. Lewkowski *ArkiVoc* 2001, **2**, 17-54; c)
 T. Ståhlberg, W. Fu, J. M. Woodley, A. Riisager, ChemSusChem,
 2011, **4**(4), 451-458; d) A.A. Rosatella, S. P. Simeonov, R. F. M.
 Frade, C. A. M. Afonso, *Green Chem.*, 2011, **13**, 754-793.
- 3 As representative selected works see M. Molinier, Y. Roman-5 Leshkov, M. E. Davis, *Proc. Natl. Acad. Sci.*, 2010, **107** (14), 6164-6168.
- 4 As selected works see a) R. O. L Souza, P. D. Fabiano; C. Feche, F. Rataboul, D. Cardoso, N. Essayem, *Catal. Today*, 2012, **195**(1), 114-119J; b) S. Yu, E. Kim, S. Park, I. K. Song, J. C. Jung, *Catal.*
- 20 Commun., 2012, 29, 63-67; c) B. Y. Yang, R. Mongomery, Carbohyd. Res., 1996, 280 (1), 27-45
- 5 As an example see S. Bhosale, M. Rao, V. Deshpande, *Microbiol. Rev.*, 1996, **60**(2), 280-300; b) H. Yu, Y. Guo, D. Wu, W. Zhan, G. Lu, *J. Mol. Catal. B Enzym.*, 2011, **72** (1-2), 73-76; c) A. M.
- 25 Dehkordi, M. S. Tehrany, I. Safari, Ind. Eng. Chem. Res., 2009, 48(7), 3271-3278
- 6 As a recent work see P. M. Grande, C. Bergs, P. Dominguez de Maria, *ChemSusChem*, 2012, **5**(7), 1203-1206 and references cited therein
- ³⁰ 7 H. Zhao, J. E. Holladay, H. Brown, Z. C. Zhang, *Science*, 2006, **316**, 15971599.
- a) Q. Zhang, K. De Oliveira Vigier, S. Royer, F. Jérôme, *Chem. Soc. Rev.*, 2012, 41, 7108-7146; b) A. P. Abbott, J. C. Barron, K. S. Ryder and D. Wilson, *Chem.-Eur. J.*, 2007, 13, 6495–6501; c) D.
- ⁵ Carriazo, M. C. Serrano, M. C. Gutierrez, M. L. Ferrer, F. del Monte, *ChemSocRev*, 2012, **1**(14), 4996-5014; d) P. D. de Maria, Z. Maugeri, *Curr. Opin. Chem. Biol*, 2011, **15**(2), 220-225.
- 9 F. Liu, K. De Oliveira Vigier, M. Pera-Titus, Y. Pouilloux, J.-M. Clacens, F. Decampo, F. Jérôme, *Green Chem*, 2013, **15**, 901-909
- 40 10 J. T. Gorke, F. Srienc and R. J. Kazlauskas, Chem. Commun., 2008, 1235–1237
 - 11 Y. J. Pagán-Torres, T. Wang, J. M. R. Gallo, B. H. Shanks, J. A. Dumesic, ACS Catal., 2012, 2, 930–934
- As a very recent work see V. Choudhary, S. H. Mushrif, C. Ho, A.
 Anderko, V. Nikolakis, N. S. Marinkovic, A. I. Frenkel, S. I. Sandler,
 D. G. Vlachos J. Am. Chem. Soc., 2013, 135, 3997-4006 and
- references cited therein.
 13 S. Saravanamurugan, M. Paniagua, J. A. Melero, A. Riisager, J. Am. Chem. Soc., 2013, 135 (14), 5246–5249.
- 50 14 F. Ilgen, D. Ott, D. Kralish, C. Reil, A. Palmberger, B. König, Green Chem., 2009, 11, 1948–1954.
- 15 as a selected comprehensive review from authors see S. G Wettstein, D. Martin Alonso, E. I Gürbüz, J. A Dumesic, *Curr Opin. Chem. Eng.*, 2012, 1, 218-224 and references cited therein.
- 55 16 P. De Vresse, N. R. Brooks, K. Van Hecke, L. Van Meervelt, E. Matthijs, K. Binnemans, R. Van Deun, *Inorg. Chem.*, 2012, **51**, 4972-4981.
- 17 F. Liu, J. Barrault, K. De Oliveira Vigier, F. Jérôme, *ChemSusChem*, 2012, 5 (7), 1223-1226
- 60 18 A) Y. Yang, C.-W. Hu, M. M. Abu-Omar, *ChemSusChem*, 2012, 5, 405; b) Y. Yang, C.-W. Hu, M. M. Abu-Omar, *Green Chem.*, 2012, 14, 509.
- 19 R. Carrasquillo-Flores, M. Käldstöm, F. Schüth, J. A. Dumesic, R. Rinaldi, ACS Catal., 2013, 3, 993-997
- 65 20 N. Meine, R. Rinaldi, F. Schüth, ChemSusChem, 2012, 5, 1449-1454
- 135

140

0

Published on 03 September 2013. Downloaded by Brown University on 06/09/2013 07:28:37

Graphical abstract

Selectivity enhancement in the aqueous acid-catalyzed conversion of glucose to 5-hydroxymethylfurfural induced by choline chloride

Fei Liu,^a Maïté Audemar,^a Karine De Oliveira Vigier,^a Damien Cartigny,^b Jean-Marc Clacens,^b ¹⁰ Margarida F. Costa Gomes,^c Agilio A. H. Pádua,^c Floryan De Campo,^b and François Jérôme^{*a}



Here, we wish to show that addition of choline chloride to an aqueous solution of glucose containing a catalytic amount of AlCl₃ or FeCl₃ promotes the ³⁰ tandem isomerization/dehydration of glucose to 5-hydroxymethylfurfural with yield up to 70%. Such strategy can be also transposed to cellulose.

35

Published on 03 September 2013. Downloaded by Brown University on 06/09/2013 07:28:37.