

TABLE I. Absorption of Immunoprecipitable ^{131}I *via* Portal Vein and Lymphatics after Intraintestinal Administration of ^{131}I -EWL 2 mg/kg in Rats

Absorption	EWL eq μg	% of dose
Amount <i>via</i> portal vein (A)	10.61 ± 2.40	1.99 ± 0.41
Amount <i>via</i> lymphatics (B)	0.37 ± 0.18	0.07 ± 0.04
Amount <i>via</i> both routes (A + B)	10.98 ± 2.45	2.06 ± 0.43
Percentage of lymphatics $[100 \times (\text{B})/(\text{A} + \text{B})]$	3.24 ± 1.69	

Each value is represented as mean with S.E. of 4 experiments.

Table I indicates that the main route in intestinal absorption of ^{131}I -EWL is the portal vein, not lymphatics. This result seems to be in accordance with the results of heparin^{1d)} and insulin.^{1e)} However, as compared with the result that 36% of the total amount of elastase absorbed was *via* lymphatics,^{1e)} the percentage of lymphatics absorption of ^{131}I -EWL is extremely small. Thus, further studies on the relationship between the physicochemical properties of macromolecules and the absorption rate *via* lymphatics will be necessary. In addition, whether or not the immunoprecipitable and protein-bound ^{131}I in serum and lymph originates from intact ^{131}I -EWL administered remains to be investigated.

Acknowledgement The authors thank Dr. S. Ohtake for his valuable advice, Mr. Y. Hidaka for the gift of purified hen egg white lysozyme and also Mr. T. Naito for his support of this research.

Department of Drug Metabolism,
Section of Experimental
Therapeutics Research,
Research & Development Division,
Eisai Co., Ltd.,
4-6-10 Koishikawa Bunkyo-ku Tokyo

TERUAKI YUZURIHA
KOUICHI KATAYAMA
TAKESHI FUJITA

Received August 31, 1973

[Chem. Pharm. Bull.]
21(12)2809-2811(1973)

UDC 547.677.2.04 : 541.14

Wave Length Dependent Photolysis of Phenanthrene Oxide

Though the chemistry of monocyclic arene oxides has been investigated,¹⁾ studies on polycyclic arene oxides are very poor.²⁾ The evidence, however, is growing that arene oxides are the active carcinogens formed from polycyclic aromatic hydrocarbons.³⁾ The stereochemistry of polycyclic arene oxides is dreamy: the epoxide ring is nearly perpendicular to the aromatic ring(s) from investigation by a Dreiding Model. This may cause an extensive conjugation between the pi-bonds and the C-O sigma-bonds.⁴⁾ These situations prompted us to investigate the chemical behaviour of compounds of this kind. In this communication

- 1) J.W. Daly, D.M. Jerina and B. Witkop, *Experientia*, **15**, 1129 (1972).
- 2) D.M. Jerina and D.R. Boyd, *Tetrahedron letters*, **1970**, 457; M.S. Newman and S. Blum, *J. Am. Chem. Soc.*, **86**, 5559 (1964).
- 3) for example, J.K. Selkirk, H. Huberman and C. Heiderberger, *Biochem. Biophys. Res. Commun.*, **43**, 1010 (1971).
- 4) T.G. Traylor, W. Hanstein, H.J. Berwin, N.A. Clinton and R.S. Brown, *J. Am. Chem. Soc.*, **93**, 5715 (1971).

we wish to report that an interesting wave length dependent photochemistry of 9,10-dihydro-9,10-epoxyphenanthrene (**1**).²⁾

Irradiation of the epoxide (**1**) in methylene chloride with a high pressure mercury lamp or a monochrometer⁵⁾ (300—350 nm) gave 9-phenanthrol (**2**, 50%),⁶⁾ phenanthrene (**3**, 11%) and an oxepine (**4**, 0.5%, *vide infra*.) The very similar reaction as above was observed by irradiation of very short wave ultraviolet (UV) light. Irradiation of **1** in methylene chloride by the monochrometer (207—235 nm) gave 9-phenanthrol (**2**, 56%), phenanthrene (**3**, 0.2%), the oxepine (**4**, 3%) and fluorene (**5**, 0.2%).




Fig. 1

TABLE I. Wave Length Dependence of Products

Wave length nm	2	3	4	5
300—350	50	11	0.5	0
250—290	8	0	20—30	3
207—235	56	0.2	3	0.2

On the other hand, irradiation of **1** in methylene chloride with a low pressure mercury lamp gave dibenzo[*b,d*]oxepine (**4**, 20—30%) as the major monomeric product, which was reported⁷⁾ after the completion of the present result, 9-phenanthrol (**3**, 8%), fluorene (**5**, 3%), and a dimer (about 25%), but no phenanthrene was observed. The compound **4** was isolated by a silica gel column chromatography, and easily crystallised from methanol to give prisms, mp 47—47.5°, UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm (ϵ): 237 (37700), 274—276 (5700). The nuclear magnetic resonance (NMR) spectra of **4** (in CDCl₃, δ =5.98 (J =6), δ =6.85 (J =6) for the olefinic hydrogens) and of a hydrogenation product, **6**, mp 114—115°, (in CDCl₃, two triplets at δ =4.59 and at δ =2.82, J =6.5) support the structure **4**. Such a localized double bond of **4** has been expected by a molecular orbital calculation for the compound.⁸⁾ The oxepine could be isolated by the photolysis in benzene or *n*-hexane, but in methanol a hemiacetal methyl ether (**7**) was formed in 40% yield, instead. The methyl ether was purified by a column chromatography and a

5) Concave Radiating Monochrometer, JASCO-CRM-FA.

6) All the new compounds were analysed and their molecular weight were determined by a mass spectrometer. Yields are dependent on concentration and reaction time. Yields shown were determined by gas chromatography after disappearance of the starting epoxide on thin-layer chromatography (0.1% solution.) **1** is thermally stable under the TLC conditions.

7) N.E. Breghtwell and G.W. Griffin, *Chem. Commun.*, 1973, 37.

8) M.J.S. Dewar, *Tetrahedron*, 26, 4269 (1970).

molecular distillation (120° (bath)/0.1 mmHg), mp 54–55°, NMR (CDCl₃) δ : 5.30 (1H, q, $J=8$ and 5.5), 3.62 (OCH₃), and 2.80 (2H, $J_{AB}=14$, $J_{AX}=8$, $J_{BX}=5.5$).

These experiments which were summarised in the Table demonstrates that the epoxide (**1**) does undergo photochemical rearrangements in three different modes according as the three regions of irradiated wave length. Two different excitations give the same major product, 9-phenanthrol, and the other excitation gives other products in a completely different way.




Fig. 2

The formation of the oxepine is reasonably explained by an electrocyclic ring opening of a rearranged epoxide (**8**),⁹⁾ which is a thermal reaction and gains a big aromatic stability. Two routes to the intervening epoxide (**8**) are possible as discussed by Griffin⁷⁾. If the rearrangement is concerted, a [1,5]sigmatropic reaction with inversion of the reaction center (oxygen) (see Fig. 2, **9**) is photochemically allowed. It is possible to speculate a nonconcerted process involving the photochemical cleavage of the C–O bond which gives a diradical or a dipole (**10**, * shows an unpaired electron or a charge.) They may attack the aromatic ring thermally to form the epoxide (**8**), though such a process sacrifices two aromatic rings. Among these two mechanisms leading to the epoxide (**8**), the concerted mechanism looks more attractive because the overlap shown in **9** is favorable by the σ - π conjugation,⁴⁾ and in turn the photochemical rearrangement to 9-phenanthrol seems to involve the cleavage of the C–O bond (*via* **10**). The formation of fluorene is compared with the photolysis of steroidal epoxyketones,¹⁰⁾ and that of phenanthrene could be discussed in connection with the interesting fate of oxygen atom.¹¹⁾

Faculty of pharmaceutical Sciences
University of Tokyo
Hongo, Bunkyo-ku, Tokyo

KOICHI SHUDO
TOSHIHIKO OKAMOTO

Received May 31, 1973

- 9) E. Vogel and H. Gunster, *Angew. Chem.*, **79**, 684 (1967).
- 10) D. Bauer, T. Iizuka, K. Shaffner and O. Jeger, *Helv. Chim.*, **55**, 852 (1972).
- 11) Methylene instead of oxygen atom, see, D.B. Richardson, L.R. Durett, J.M. Martin, W.E. Putnum, S.C. Slaymaker and I. Davoretcky, *J. Am. Chem. Soc.*, **87**, 2763 (1965).