

Synthetic and Mechanistic Studies of a Versatile Heteroaryl Thioether Directing Group for Pd(II) Catalysis

Andrew M Romine, Kin S. Yang, Malkanthi K. Karunananda, Jason S Chen, and Keary M. Engle

ACS Catal., **Just Accepted Manuscript** • DOI: 10.1021/acscatal.9b01471 • Publication Date (Web): 02 Jul 2019

Downloaded from <http://pubs.acs.org> on July 3, 2019

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Synthetic and Mechanistic Studies of a Versatile Heteroaryl Thioether Directing Group for Pd(II) Catalysis

Andrew M. Romine[†], Kin S. Yang[†], Malkanthi K. Karunananda[†], Jason S. Chen[†], and Keary M. Engle^{*,†}

[†]Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, United States


^{*}Automated Synthesis Facility, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, United States

ABSTRACT: A weakly coordinating monodentate heteroaryl thioether directing group has been developed for use in Pd(II) catalysis to orchestrate key elementary steps in the catalytic cycle that require conformational flexibility in a manner that is difficult to accomplish with traditional strongly coordinating directing groups. This benzothiazole thioether, (BT)S, directing group can be used to promote oxidative Heck reactivity of internal alkenes providing a wide range of products in moderate to high yields. To demonstrate the broad applicability of this directing group, an arene C–H olefination method was also successfully developed. Reaction progress kinetic analysis provides insights into the role of the directing group in each reaction, which is supplemented with computational data for the oxidative Heck reaction. Furthermore, this (BT)S directing group can be transformed into a number of synthetically useful functional groups, including a sulfone for Julia olefination, allowing it to serve as a “masked olefin” directing group in synthetic planning. In order to demonstrate this synthetic utility, natural products (+)-salvianolic acid A and salvianolic acid F are formally synthesized using the (BT)S directed C–H olefination as the key step.

INTRODUCTION

Substrate directivity is a powerful approach for enhancing reactivity and controlling selectivity across numerous types of metal-catalyzed reactions.¹ In substrate-directed reactions, pre-association of the catalyst to one or more Lewis basic sites on the substrate promotes a desired reaction outcome by perturbing the activation barriers for possible divergent pathways along the reaction coordinate. Kinetic reactivity is typically enhanced by virtue of induced intramolecularity as well as enforced proximity between the reactive group on the substrate and a free coordination site on the metal. Regioselectivity is dictated by preferential formation of a thermodynamically and/or kinetically stabilized metalacycle over an energetically disfavored constitutional isomer. Stereoselectivity can also be controlled in key bond-making and bond-breaking steps through stereochemical information on (or in the vicinity of) the coordinating atom(s).

Strategic application of substrate directivity in palladium-catalyzed alkene functionalization is a potent platform for achieving challenging modes of bond construction.^{1b} Palladium-catalyzed Wacker oxidation and Mizoroki–Heck arylation of alkenes constitute the bedrock of modern homogeneous catalysis research, with the power of these reactions stemming from their ability to forge diverse C–C and C–heteroatom bonds from simple alkene starting materials.^{1d,2} In the traditional reaction pathways, upon nucleometalation (or migratory insertion), the resulting alkylpalladium(II) intermediate undergoes facile β -H elimination to deliver the final oxidized alkene product. With substituted alkene starting materials, the regioselectivities of the initial addition step and subsequent β -H elimination step are not straightforward to control in the absence of a strong steric or electronic bias between the two alkenyl carbon atoms. Our laboratory^{1e,3} and others^{1d,4} have studied substrate-directed palladium-catalyzed alkene functionalization reactions that initiate via Wacker- or Heck-type pathways. In these systems, strongly coordinating mono-, bi-, and tri-dentate directing auxiliaries can be used



Scheme 1. Background and Synopsis

to control the regioselectivity of the alkene addition step and stabilize the resulting intermediate such that downstream elementary steps, such as protodemetalation, oxidative addition, or transmetalation, outcompete β -H elimination.

The design of new directing groups with distinct coordination properties constitutes a significant challenge and opportunity in this area of study, given the ability of the directing group to engender control of pathway selectivity during catalysis. In our past work we have used the strongly coordinating 8-aminoquinoline (AQ) and 2-pyridyl-8-aminoquinoline (PAQ) directing groups, which are efficient at suppressing β -H elimination to enable 1,2-difunctionalization and hydrofunctionalization. Based on these precedents, we hypothesized that a more weakly coordinating group may lead to a different scenario in which a β -H elimination step would now be kinetically accessible. Specifically, we sought to identify a directing group that would lie in the “goldilocks” region—

Table 1. Optimization of Directing Group for Oxidative Heck Coupling with 1,2-Dialkyl Alkenes^a

$\text{DG}-\text{CH}_2-\text{CH}=\text{CH}-\text{Me} + \text{Ph-B(OH)}_2 \xrightarrow[\text{DMSO (0.05 M), 45 °C, 6 h [for (E)-I] or 65 °C, 3 h [for (Z)-I]}]{5 \text{ mol\% Pd(OAc)}_2, 1.5 \text{ equiv BQ}}$
 $\text{DG}-\text{CH}_2-\text{CH}(\text{Ph})-\text{CH}_2-\text{Me}$

(E)-I or (Z)-I (1.4 equiv) (E)-II or (Z)-II

Directing Group (DG)	Yield (II)		Directing Group (DG)	Yield (II)		Directing Group (DG)	Yield (II)	
	[from (E)-I]	[from (Z)-I]		[from (E)-I]	[from (Z)-I]		[from (E)-I]	[from (Z)-I]
	<15% ^b	n.r. ^c		45% ^d	30% ^e		59%	84%
	n.r. ^c	n.r. ^c		n.r. ^c	n.r. ^c		65%	82%
	n.r. ^c	<15% ^b		32%	23%		40%	64% ^f
	n.r. ^c	n.r. ^c		86%	80%		<15% ^b	<15% ^b
	n.r. ^c	<15% ^b		85%	91%		n.r. ^c	n.r. ^c
	<15% ^b	<15% ^b		<15% ^b	25%			

^aIsolated yields except where otherwise noted. ^bDetermined by ¹H NMR of the crude reaction using 1,3,5-trimethoxybenzene as internal standard. ^cn.r.=no reaction. ^dInseparable from 22% of two unidentifiable isomers that contain a 1,2-disubstituted internal olefin motif. ^eInseparable from 26% of two unidentifiable isomers that contain a 1,2-disubstituted internal olefin motif. ^fIsolated as a 7:1 mixture of isomers (Z:E).


possessing sufficient coordination strength to facilitate complete selectivity in the nucleopalladation step, even with especially challenging substrates (such as 1,2-dialkyl alkenes), yet still weakly coordinating enough to allow selective β -H elimination. Such a reaction sequence would be synthetically valuable by allowing formal C(alkenyl)-H functionalization,^{2d} enabling conversion of a simple alkene starting material into a regio- and stereodefined tri- or tetra-substituted alkene product (Scheme 1).

To validate this general hypothesis, we elected to focus on the palladium(II)-catalyzed oxidative Heck reaction, anticipating that newly identified directing groups would also find utility in other palladium(II)-catalyzed reactions (*vide infra*). Despite extensive efforts over the past several decades, knowledge gaps persist in Heck-type chemistry. Directed variants of the palladium(II)-catalyzed oxidative Heck reactions, for instance, are relatively rare, with White's seminal study of heteroatom-directed oxidative Heck arylation of terminal alkenes a notable exception.⁵ A closely related body of literature has examined directed variants of classical-polarity palladium(0)-catalyzed Mizoroki-Heck reactions,^{1d} with pioneering work here including contributions by Hallberg,⁵ Carretero,⁶ Yoshida,⁷ and others.⁸ These studies have typically focused on heteroatom-substituted alkene substrates, including vinyl sulfides, ethers, or silanes, and in these designs, the coordinating group is built into the heteroatom linkage. For these systems, competitive binding of the phosphine ligand versus the directing group is an important consideration. Included in these studies are relatively few alkyl-substituted alkenes, with examples mostly limited to terminal allylic alcohols, amines, and their derivatives.

Across all Heck-type chemistry, internal asymmetrically substituted di- and tri-substituted alkenes remain difficult substrates. Sigman has developed elegant catalytic systems for both oxidative and non-oxidative redox relay asymmetric Heck arylation of internal alkenes.⁹ In these reactions, the highly electron-deficient palladium catalyst undergoes rapid β -H elimination/reinsertion (chain-walking) to an appropriate terminating group (e.g., an alcohol). Complementing such chain-walking Heck systems with those that preserve the alkene in a precise geometric relationship to the newly installed aryl group would be desirable in situations where the alkene is required in the resulting product. In parallel to this study, Shenvi reported an intermolecular Heck coupling that can direct arylation of electronically unbiased tri- and tetra-substituted alkenes and establish quaternary carbons while migrating the olefin only one position and not into conjugation with the carboxylic acid directing group as is common in redox-relay Heck reactions.¹⁰

RESULTS AND DISCUSSION


Directing Group and Reaction Optimization. The investigation commenced by testing a series of potential directing groups for their ability to promote a classically challenging oxidative Heck reaction with a 1,2-substituted alkene. We focused on thioether-based directing groups (I) due to on their ease of installation via nucleophilic substitution chemistry, ability to be productively removed in a variety of preparatively useful reactions (e.g., Julia olefination), and established effectiveness as directing groups in palladium complex formation¹¹ and catalysis.¹²



Scheme 2. Scope for (BT)S-Directed Oxidative Heck Reaction^a

^aIsolated yields. ^bInseparable from 8% of an unknown isomer. ^c10:1 isomeric mixture of *E* and *Z*. ^dInseparable from 7% of an unknown isomer. ^eIsolated as a 6:1 mixture of inseparable isomers (*E*:*Z*). ^fIsolated as a 3:2 mixture; isomers separated by LC. ^gInseparable from 10% of an unknown isomer. ^hFrom the *Z*-olefin; 50 mM, 65 °C. ⁱFrom the *E*-olefin; 50 mM, 6 h.

A series of homoallylic thioether-based substrates containing internal (*E*)- or (*Z*)-alkenes were thus prepared. These substrates were subjected to Pd(II)-catalyzed oxidative Heck reaction conditions, which were optimized in parallel using a Design of Experiment workflow with automated reaction setup and sampling (see SI) (Table 1). As a control, a homoallylic alcohol substrate (**I-A**) was tested and did not deliver meaningful quantities of the desired product; similarly, simple aryl thioethers and sulfoxides (**B-E**) were ineffective, despite their previously established utility as directing groups in palladium(II) catalysis.¹³ We then turned our attention to azaheterocyclic thioethers, reasoning that the coordination strength of sulfur would be attenuated in these cases and that such substrates offered additional opportunities for coordination of the palladium catalysts to an N(sp²) atom during catalysis. Tetrazole directing group **G** afforded moderate yield of the desired product with both the *E*- and *Z*-alkene starting material; unfortunately, in this case the remainder of the starting material was completely converted into undesired byproducts. Based on the successful functionalization of the internal alkene with directing group **G**, we tested **F** and **H**, but found minimal to no product. Benzoxazole thioether directing group **I** provided slightly diminished yields compared to directing group **G**. However, in this case the product selectivity was improved, as we did not observe significant byproduct formation. Fortunately, we discovered that two related directing groups, **J** and **K**, the latter of which we refer to as (BT)S throughout the remainder of the manuscript, offered high reactivity and selectivity. Yields in these cases were 86% and 80% for products (*E*)-II-J and (*Z*)-II-J, and 85% and 91% for products (*Z*)-II-K and (*Z*)-II-K. Substrates with similar thiazole-type directing groups, **L-O**, with different steric and electronic properties gave lower yields of **II**. We next tested oxidized versions of (BT)S, **P** and **Q**, as controls, but minimal to no product formation was observed in these cases.



Scheme 3. Results with Tri- and Tetra-substituted Olefins^a

^aReaction conditions: alkene (0.1 mmol), phenylboronic acid (0.14 mmol), BQ (0.15 mmol), Pd(OAc)₂ (5 mol%), DMSO (1 mL), 45 °C, 3 h, unless otherwise noted; isolated yields. ^bRun at 65 °C and in DMSO (2 mL). ^cIsolated as a 5:2:4:3 mixture of isomers; 50 mM at 65 °C. ^dIsomers separated by HPLC. ^eIsolated as a 3:2:1 mixture of isomers; isomers separated by HPLC. ^fObserved by NMR.

Collectively, these results point to a somewhat complex relationship between structure and reactivity/selectivity, which was probed further through a detailed analysis of the possible coordination modes of **K** ((BT)S) (*vide infra*). Based on its effectiveness in this challenging oxidative Heck reaction and its potential utility in downstream chemistry (e.g., oxidation followed by modified Julia olefination), **K** ((BT)S) was selected for further investigation.

Scope of Oxidative Heck Reaction. We proceeded to investigate the substrate and coupling partner scope of this directed oxidative Heck reaction (Scheme 2). Arylboronic acids with a variety of functional groups, including electron-withdrawing and electron-donating moieties, reacted under the standard conditions to give products **2a–k** in high yields. Heteroarylboronic acids also gave the corresponding products **2l–p** in reasonable yields. Notably, (*E*)-(3-phenylprop-1-en-1-yl)boronic acid provided product in reasonable yield; however both (*E*)- and (*Z*)-isomers, **2p**, were formed in a 3:2 ratio. Two representative alkylboronic acids, methylboronic acid and neopentylboronic acid, were tested, but unfortunately these were not competent coupling partners in the reaction.

Next, we moved on to probe the alkene scope (Scheme 2). We found that alkyl branching was tolerated at both the α - and β -positions, as exemplified by products **4a** and **4b**, which were formed in nearly quantitative yields. We next tested a substrate containing an intervening alcohol, which is an established terminating group in Heck chain-walking processes. Nevertheless, in this case we observed that product **4c**, with the alcohol intact, was formed in high yield. The fact that the corresponding ketone product is not formed in this case demonstrates that the alternative *endo*- β -H elimination pathway from the putative palladacycle intermediate is disfavored. Furthermore, 1,1-disubstituted alkenes were found to be tolerated, allowing for formation of **4d** in moderately high yield and **4e**, a derivative of the natural product isopulegol, in high yield. As observed in the directing group optimization table (Table 1), both *E*- and *Z*-configured 1,2-disubstituted alkenes gave high yield of the desired products (*E*)-**II** and (*Z*)-**II**, respectively, and importantly in both cases, the products were obtained as single alkene stereoisomers. With (*E*)-**I** as a representative internal alkene, other aryl boronic acids were tested, and it was found that an electron-rich aryl group led to higher reactivity compared to an electron-poor aryl group (*vide infra*). In the case of the *p*-CF₃-substituted example, a 3:1 ratio of conjugated to non-conjugated products was obtained, suggesting that the identity of the aryl group can influence the energy barriers for competitive β -H elimination pathways following migratory insertion—at least to some extent. A series of other internal alkenes bearing different linear alkyl chains were tested (**4h–4j**) along with examples containing an additional functional group potentially capable of coordinating the catalyst or perturbing the electron properties of the alkene (**4k–4m**), and in all cases moderate to high yield of the desired, conjugated product was observed. Different tether lengths were also examined, and it was found that both γ,δ - and δ,ϵ -unsaturated alkenes (**4n** and **4o**) are competent in this system. No reaction was observed with the corresponding allylic substrate. Across this series of substrates, preference for aryl addition to the alkenyl carbon atom distal from the (BT)S group is consistent with the notion that the (BT)S group is coordinated during migratory insertion. Although some exceptions exist,^{6b,6c,8k} directed Heck-type reactions typically favor formation of the regioisomer arising from a transition state where the migrating aryl or alkenyl group is exo-cyclic to incipient palladacycle,^{1d,5–8} even with internal non-conjugated alkenes,^{3h,4d,10} likely due to minimization of strain compared to the alternative possibility with the aryl group in an *endo*-cyclic orientation.

Tri- and tetra-substituted olefins (**3p–3r**) were also examined (Scheme 3). In these cases complex product mixtures were typically obtained, indicating that these substrate classes are generally incompatible with this method, likely due to increased steric hindrance and due to a greater number of possible side reactions.¹⁰ With tetrasubstituted substrate **3p**, the formation of products **4r** and **4s** suggests that positional isomerization of the alkene takes place under the reaction conditions with this substrate. One interpretation of this result is that in cases where migratory insertion is sluggish due to steric hindrance, other off-





Figure 1. (A) Same-excess experiments with and without product. (B) Different-excess experiments of starting material, phenylboronic acid, and benzoquinone.^aDue to the consumption of 1 equivalent of Benzoquinone per turnover, reactions were set up with Pd(OAc)₂ (2.5 mM) and with (*E*)-**I** (variable), **III**, **BQ**, and (*E*)-**II** (for same-excess 3) corresponding to the expected amounts at time = 80 min for the standard reaction in DMSO (2.0 mL), at 45 °C, and in air. ^bReaction conditions: (*E*)-**I** (variable), **III** (variable), Benzoquinone (variable), Pd(OAc)₂ (variable), DMSO (2.0 mL), 45 °C, air.

cycle pathways can predominate. With substrates **3p** and **3r**, the requirement to undergo β -hydride elimination in an *endo*-fashion (towards the directing group and within the palladacycle) rather than the preferred *exo*-fashion is also likely a contributing factor to low yields.

With alkene **3q**, the product distribution suggests that the migratory insertion step was selective, but that subsequent β -H elimination was unselective. Of the three tri/tetrasubstituted alkenes tested, **3r** offered the highest selectivity. In this case, product **4w** containing a quaternary carbon was formed as the main product along with regioisomer **4x** in a 5:1

ratio and 49% combined yield.

Reaction Progress Kinetic Analysis (RPKA). The high regio- and stereo selectivity of this method with diverse 1,2-disubstituted alkenes prompted us to delve into the reaction mechanism by combining reaction kinetics and computational studies. First, to elucidate the kinetic



	[(E)-I]	[III-1]	[BQ]	[Pd(OAc) ₂]
10 mol%	50 mM	70 mM	75 mM	5.00 mM
7.5 mol%	50 mM	70 mM	75 mM	3.75 mM
5 mol%	50 mM	70 mM	75 mM	2.50 mM
4 mol%	50 mM	70 mM	75 mM	2.00 mM
3 mol%	50 mM	70 mM	75 mM	1.50 mM





Figure 2. Reaction conditions (*E*)-I (50 mM), III-1 (70 mM), Benzoquinone (75 mM), Pd(OAc)₂ (variable), DMSO (2.0 mL), 45 °C, air.






Figure 3. Reaction conditions (*E*)-I (50 mM), III-1-3 (70 mM), Benzoquinone (75 mM), Pd(OAc)₂ (5 mol%), DMSO (2.0 mL), 45 °C, air.


possible monodentate coordination modes




possible bidentate coordination modes



Scheme 4. Potential Binding Modes of Pd^{II} to (BT)S Group



Scheme 5. Crystal Structure of Pd(tfa)₂O₂ Complex



Scheme 6. Relative energies of Possible Intermediates in the Oxidative Heck Reaction^a

^aRelative free energies of the intermediates of the catalytic cycle a) before and b) after migratory insertion.

features of catalytic process, we performed reaction progress kinetic analysis (RPKA)¹⁴ using representative (*E*)- and (*Z*)-alkene substrates. Insights for both substrate types were generally mutually consistent, so the ensuing analysis focuses on the (*E*) substrate, and information regarding the (*Z*)-alkene substrate can be found in the SI. Regarding general kinetic behavior, we noted that the reaction appears to contain two regimes: one that is operative until 35–50% conversion is reached (representing approximately the first 60 min for the reaction with the (*E*)-alkene and 30 min for the reaction with the (*Z*)-alkene under standard conditions), and a second regime from that point forward where the reaction proceeds more slowly. This could arise from several different

root causes, including uncharacterized catalyst deactivation processes. Though both regimes are valuable to understand, we focused our attention on this initial stage of the reaction, as it is more likely to reflect the intrinsic kinetics before other off-cycle processes predominate.

First, we performed a “same-excess” experiment, which simulates entering the catalytic cycle after multiple catalyst turnovers in order to probe for catalyst activation or deactivation pathways (Figure 1A). The time-adjusted same-excess experiment without added product was observed to proceed at a faster rate than the standard reaction, pointing to possible catalyst deactivation or product inhibition. Results were similar




Figure 4. Computed energy profile for migratory insertion and β -hydride elimination steps.




Figure 5. Computed transition states for migratory insertion (left) and β -hydride elimination (right) steps.





Figure 6. Deuterated starting material experiment for the directed oxidative Heck reaction. Reaction conditions (*(E)*-I or (*(E)*-I-*d*₂ (50 mM), *III*-I (70 mM), Benzoquinone (75 mM), Pd(OAc)₂ (5 mol%), DMSO (2.0 mL), 45 °C, air.



Scheme 7. (BT)S-directed Oxidative Heck Catalytic Cycle

with added product, suggesting that the catalyst is being deactivated through off-cycle processes. Catalyst deactivation in this reaction can take place if Pd(0) is not efficiently reoxidized, enabling precipitation of palladium black.

Next, driving forces were determined through a series of “different-excess” experiments (Figure 1B). We first varied catalyst concentration and found that the reaction rates increased at higher catalyst concentration. Visualizing the data using the Burés method (Figure 2)¹⁵ with an x-axis that is normalized by catalyst concentration reveals that the reaction is first-order in [Pd]. We then examined the effect of changing the concentration of others reaction components, namely decreased alkene concentration [*(E)*-I] (different-excess [*(E)*-I]), decreased phenylboronic acid concentration [*III*] (different-excess [*III*]), and decreased benzoquinone concentration [BQ] (different-excess [BQ]). Overlay between the standard reaction and the different-excess BQ experiment throughout the time course is consistent with apparent zero-order

kinetics in [BQ]. The different-excess (*(E)*-I and different-excess [*III*] experiments overlay reasonably well with the standard reaction at <30% conversion (first 30 min) and then deviate as the limiting reagents are consumed. This indicates apparent zero-order kinetics in [*(E)*-I] and [*III*] early in the reaction. First-order kinetics in [Pd] combined with zero-order kinetics in [*(E)*-I], [*III*], and [BQ], rule out substrate binding, transmetalation, and reoxidation, respectively as possible turnover-limiting steps, pointing to either migratory insertion of a substrate-bound [Pd(II)–Ar] species or β-H elimination as possible turnover-limiting steps. As mentioned above, electron-rich arylboronic acids were found to be more reactive (Figure 3). To quantify this trend, we measured initial rates of the three representative arylboronic acids with *p*-CF₃, -H, and -OMe groups and found *k*_{rel} values of 1:3:8 across this series. This result does not necessarily allow one to disambiguate between the two mechanistic possibilities described above, as migratory insertion and β-H elimination could both potentially be facilitated by an electron-rich aryl group due to a more nucleophilic [Pd(II)–Ar] species and a more hydridic C–H bond, respectively. Since the turnover-limiting step cannot be distinguished based on the available rate dependencies, we next turned to complementary techniques to gain insight on these steps and other aspects of the catalytic cycle.

Directing Group Coordination. The results in Table 1 revealed the structural features that are required for a suitable directing group, demonstrating the importance of the embedded benzo-fused methyldiene dithioacetal motif in **J** and **K**. Nevertheless, **K** ((BT)S) also possesses an N(sp²) capable of coordinating the palladium catalyst, complicating the potential coordination chemistry involved, which we sought to elucidate through crystallography and computation studies. Several different mono- and bidentate binding modes can be envisioned (Scheme 4), and to probe this, we first attempted to synthesize various palladium complexes of potential relevance to catalysis. Indeed, we were able to obtain a representative product-bound palladium species by combining **2a** and Pd(tfa)₂. The resulting 2:1 complex, **Pd-I**, was amenable to characterization by X-ray crystallography, and in the structure, the (BT)S directing group is coordinated in a monodentate fashion through the nitrogen atom, consistent with earlier literature precedents (Scheme 5).¹⁶ In this case the alkene was not bound to the palladium center. The same monodentate nitrogen binding mode was also later observed with a representative starting material–palladium complex for a related C–H activation reaction (*vide infra*).

DFT Calculations. Next, we performed density functional theory (DFT) calculations to further probe key aspects of the reaction mechanism. First, the possible monodentate coordination modes were evaluated using **1** as the model substrate. Geometry optimizations were performed on the intermediates of the proposed catalytic cycle at the migratory insertion step. Three coordination modes were investigated, i.e. coordination through N (INT1_N and INT2_N), coordination through S of the heterocycle (INT1_S, INT2_S) and coordination through S outside the heterocycle (INT1_S', INT2_S'). The N-coordinated intermediates were lowest in energy, followed by the S(proximal)-coordinated intermediates (Scheme 6). The S(distal)-coordinated intermediates were highest in energy. These trends also held in the corresponding transition state energies for the migratory insertion step (see Figure S19 in the SI).

One explanation that reconciles these results with the observation that both directing groups **J** and **K** ((BT)S) are similarly effective (Table 1) is that the catalytic reaction can take place with the directing group bound either through the proximal-S atom or through the N-atom. With BT(S), the latter pathway is computationally predicted to be lower-energy, though the former is also energetically accessible. The S(proximal)-coordinated pathway is likely to be operative in cases such as **J**, where the N-atom is absent.


Probing the Turnover-Limiting Step. As experimental mechanistic investigations revealed the rate determining step to be either migratory insertion or β -hydride elimination, DFT calculations were performed to locate the transition states to determine the energetics of these two steps. A truncated (*E*) internal alkene was used as the model substrate to minimize computational time. The energy profile is shown in Figure 4. The *syn*-migratory insertion step (**TS1**) has an activation free energy barrier of 14.3 kcal/mol with respect to the π -alkene complex **INT1**. The subsequent β -hydride elimination step has a higher activation energy barrier of 15.7 kcal/mol (**TS2**) indicating that the β -hydride elimination is the turnover-limiting step. The optimized structures of **TS1** and **TS2** are shown in Figure 5. The weaker coordination of the (BT)S group is evidenced by the longer Pd \cdots N distances of the transition states at both migratory insertion (2.18 Å) and β -hydride elimination (2.22 Å) steps. Shorter Pd \cdots N distances have been reported for transition states with strongly coordinating directing groups such as 8-aminoquinoline and 2-pyridyl-8-aminoquinoline (2.02–2.08 Å).^{3e,17} The energy profile for the migratory insertion and β -hydride elimination was also calculated for the terminal alkene, **1** (see SI; Figure S17 and S18). The calculations revealed that the formation of the *E*-isomer is kinetically favored at the rate- and stereoselectivity-determining step, β -hydride elimination, consistent with experimental selectivity (see SI).

With the computational data suggesting that β -hydride elimination is

containing deuterated ((*E*)-**II-d₂**) and non-deuterated ((*E*)-**II**) alkenyl positions (Figure 6). By monitoring desired product formation over the initial 45 min, we indeed found a primary KIE, $k_H/k_D = 2.0$. Additionally, we found that with the deuterated substrate the product distribution was not as clean, potentially due to other pathways (such as β -H elimination at other positions) being kinetically competitive when the typically favored β -H is replaced with a deuterium. Overall, the experimental and computational results are consistent with β -H elimination as the turnover-limiting step.

By combining the insights gathered from both DFT calculations and the kinetics experiments, we propose the general catalytic cycle as shown in Scheme 7. For simplicity, only the N-bound intermediates are shown, although as mentioned above S(proximal)-bound intermediates are also energetically accessible and could be active participants in the catalytic cycle. The reaction begins with substrate binding and transmetalation steps (in either potential order) to give intermediate **Pd-B**. Following this, a *syn*-migratory insertion gives the palladacycle **Pd-C**. Product is then generated via β -H elimination giving the Pd-hydride species **Pd-D**, which can undergo reductive elimination and reoxidation with BQ to provide the Pd(II) species required for turnover.

C(aryl)-H Olefination Background. Having established the efficacy of the (BT)S directing group in oxidative Heck chemistry, we were interested in exploring its utility in other palladium(II)-catalyzed reac-




Scheme 8. Scope for (BT)S-directed C-H Functionalization^a

^aReaction conditions: **5a–aa** (0.10 mmol), Alkene (1.2 equiv), AgOAc (3 equiv) Benzoquinone (0.5 equiv), Pd(OAc)₂ (10 mol%), tAmylOH (1 mL), Boc-L-phenylalanine (0.2 equiv), DMSO (7 equiv), 4Å molecular sieves, 75 °C, air, 6 h. Percentages refer to isolated yields. ^b110 °C, nBu₄NPF₆ (2 equiv), 2h. ^cAlkene (2.2 equiv). ^dRatio of mono-, mono', and bis-olefinated products, respectively, with the major product depicted. Mono- and mono'-olefinated products isolated as a mixture. ^eRatio of mono- and bis-olefinated product, respectively, with the major product depicted. ^fAlkene (2.2 equiv), 24h. ^g24h. ^hAlkene (2.2 equiv), 24 h.

the turnover-limiting step, we next examined whether there was a kinetic isotope (KIE) in the rates between a representative substrate


tions involving distinct elementary steps. Specifically, we were attracted to directed *ortho*-C–H olefination (Fujiwara–Moritani-type coupling)

since the use of a versatile “masked olefin” directing group could allow rapid access to a number of conjugated natural products and materials.¹⁸ Weakly coordinating directing groups, including those distal to the C–H bond of interest, have provided a valuable solution in many arene C–H functionalization reactions catalyzed by palladium, including



A^d

	[5c]	[7]	[6c]	[6c']
standard	0.100 M	0.200 M	0 M	0 M
same-excess 1	0.075 M	0.160 M	0 M	0 M
same-excess 2	0.075 M	0.160 M	0.020 M	0.003 M



B^b

	[5c]	[7]	[6c]	[6c']
standard	0.100 M	0.120 M	0 M	0 M
diff.-excess 5c	0.060 M	0.120 M	0 M	0 M
diff.-excess V	0.100 M	0.200 M	0 M	0 M





Figure 7. (A) Same-excess experiment. (B) Different-excess experiments of starting material and ethyl acrylate.^aDue to secondary reaction of ethyl acrylate, same-excess 1 and 2 were set up with amounts of **5c**, **7**, **6c**, **6c'**, BQ (0.375 M), and AgOAc (2.250 M) to equal the amounts observed by qNMR at time = 45 min for the standard reaction (see Figure S16 in SI).^bReaction conditions unless otherwise stated: **5c** (variable), **7** (variable), AgOAc (3 equiv), Benzoquinone (0.5 equiv), Pd(OAc)₂ (10 mol%), tAmylOH (1 mL), Boc-L-phenylalanine (0.2 equiv), DMSO (7 equiv), 4Å molecular sieves, 75 °C, air.

amination,¹⁹ formation of lactams,²⁰ construction of dihydrobenzofurans,²¹ iodination,²² and olefination.²³ We were encouraged to see that Zhang has shown the utility of using phenyl-, methyl-, and *p*-tolyl-thioethers²⁴ and the corresponding sulfoxides²⁵ for directed arene olefination, suggesting that the (BT)S directing group may provide a suitable directing group and then be able to serve as a modified Julia olefin precursor for easy transformation to the desired alkene. Furthermore, Maiti has developed an innovative sulfone directing group for *meta*-hydroxylation, which can subsequently serve as a Julia olefination precursor, and we hoped to provide a similar solution for *ortho*-C–H functionalization.²⁶


C–H Functionalization Substrate Scope. After reexamining the directing groups **A–H** and **K–Q** in Table 1 under optimized MPAA-accelerated C–H olefination reaction conditions,²⁷ it was clear that (BT)S again was suitable in its ability to successfully direct the desired reaction (Table S13 in supporting information).

Therefore, we next examined the scope of the (BT)S-directed C–H olefination reaction (Scheme 8). We were pleased to see that the electron-rich starting material **5a** produced a nearly quantitative yield, and electron-rich, electron-poor, and alkyl functional groups were tolerated to yield **6b–h** and **6k–n** in moderate to high yields. The (BT)S directing group can also promote *ortho*-olefination in polyaromatic ring systems, such as naphthalene producing **6i–j** in moderate yields. As expected, a number of acrylates can be tolerated in this reaction to give **6o–q** and



Scheme 9. Unactivated Alkene Reactivity^a

^a (BT)S directed reaction run with the following conditions: starting material (0.10 mmol), Alkene (1.2 equiv), AgOAc (3 equiv) Benzoquinone (0.5 equiv), Pd(OAc)₂ (10 mol%), tAmylOH (1 mL), Boc-L-phenylalanine (0.2 equiv), DMSO (7 equiv), 4Å molecular sieves, 75 °C, air, 6 h; other reactions run both under above conditions and under originally reported conditions.



Scheme 10. Crystal structure of Pd(tfa)₂O **5c** complex

6s–t in high yields. We were excited to see that this (BT)S-directed olefination could also be expanded to other alkene classes such as acrylamide giving **6r** and styrene giving **6u**. Interestingly, methyl methacrylate gave the non-conjugated product **6w**, consistent with a mechanism in which β -H elimination proceeds *exo* to the putative metalacycle in this case. Other competent alkene coupling partners include phenyl vinyl sulfone (**6x**), dodecene (**6y**), ethenesulfonyl fluoride (**6z**)²⁸, and ethyl crotonate (**6aa**).


Notably, **6y** represents a somewhat unique product in directed arene olefination, as the use of non-conjugated terminal alkenes remains rare.²⁹ Notably, the thioether²⁴ and sulfoxide²⁵ directed reactions designed by Zhang were replicated with similar yield as previously reported for ethyl acrylate but yielded no product with dodecene showing the unique characteristics of (BT)S (Scheme 9).

C–H Olefination RPKA. We decided to probe the catalytic cycle of this reaction in an effort to better understand the role of (BT)S in directing C–H olefination. To this end, same excess and different excess experiments were performed using the simple, unsubstituted arene **5c** as a standard substrate.


Due to unique facets of this reaction, several experimental modifications were made compared to the standard same-excess and different-excess protocols that are generally used. For both sets of data, starting material concentration, $[5c]$, was plotted on the y-axis as opposed to product because both **6c** and **6c'** are formed in the reaction, with **6c** presumably being formed from which a second catalytic cycle to install an additional ethyl acrylate yields **6c'**. Moreover, the presence of a

secondary reaction converting **6c** to **6c'** meant that the standard method of assuming a one-to-one coupling (and hence consumption) of starting material **5c** and coupling reagent **7** for the design of the same-excess experiment would not accurately reflect reaction conditions at 45 min. Therefore, values for **5c**, **7**, **6c**, and **6c'** as well as BQ were determined by qNMR at 45 min in the standard reaction, and those values were used when designing the same-excess experiments (see Figure S16 in the Supporting Information). The amount of AgOAc was also lowered to appropriately reflect oxidant consumption for each presumed cycle.

The same-excess data shows excellent overlay for the standard reaction and same-excess 2, which mimics the standard reaction at 45 min (Figure 7A). However, same-excess 1, which does not contain added product, shows an accelerated rate. This qualitatively implies that the decreased reaction rate is due to product inhibition rather than catalyst deactivation. This result, in combination with the result from the different-excess experiment discussed below, suggest a catalytic cycle in which palladium coordinates strongly to the products **6c** and **6c'**, and, due to this strong affinity, produces product **6c'** at a similar rate to the rate of formation of product **6c** at around 0.040M concentration of **6c**.



Scheme 11. Productive Removal and Transformation of the (BT)S Directing Group



Scheme 12. Formal Synthesis of (+)-Salvianolic acid A and Salvianolic acid F

Regarding the different excess experiments, overlay between the standard run and different-excess **V** shows a zero-order dependence in $[V]$ suggesting that ethyl acrylate is not involved in the turnover-limiting step of this catalytic cycle (Figure 7B). Different-excess **5c** has a reduced rate compared to the standard reaction, showing a positive-order dependence in $[5c]$. This result is potentially consistent with either irreversible substrate binding between (BT)S and Pd or reversible substrate binding followed by turnover-limiting C–H functionalization. We favor the latter interpretation because substrate binding and dissociation is expected to be rapid. We were unable to detect the putative cyclopalladated intermediate under a variety of conditions, consistent with the notion that C–H activation may be the slow step in catalysis.

Directing Group Coordination. Similar to our mechanistic studies in the oxidative Heck reaction, we attempted to synthesize various complexes relevant to catalysis, and in this case were able to successfully isolate a starting-material-bound palladium trifluoroacetate complex **Pd-2**, which we could characterize by X-ray crystallography. As with the above crystal for the oxidative Heck reaction, the complex shows coordination of palladium through the nitrogen of the benzothiazole (Scheme 10).


Productive Removal and Transformation. Product **2a** from the oxidative Heck reaction was used as a model substrate for directing group conversion and removal (Scheme 11). Moderate yield was observed in a palladium-mediated isocyanide insertion, the product of which was isolated as **8** after acidic workup.³⁰ Fortunately, by treating **2a** with ammonium molybdate tetrahydrate, the sulfone product **9** could be formed in quantitative yield with no apparent oxidation of the olefin.³¹ Sulfone **9** can be subsequently modified through several methods. Treatment with sodium borohydride affords a nearly quantitative yield of **10** containing the versatile sulfinic acid functional group.³² A modified Julia olefination with veratraldehyde gave product **11** in 54% isolated yield as the (*E*)-alkene.³³ Sulfone **9** was also transformed to the carboxylic ester **12** in high yield via treatment with ethyl cyanofornate in the presence of LiHMDS and subsequent removal of the benzothiazole-sulfone using Zn-dust from the isolated intermediate.³⁴ Oxidation of **2a** with *m*CPBA also gave high yield of sulfoxide **13**. Successive treatment of **13** with ammonium acetate gave the sulfanone product **14** in 61% yield.³⁵ A representative C–H olefination product, **6c**, was also oxidized and subsequently transformed to the (*E*)-alkene **18** with veratraldehyde in 78% isolated yield.

Synthetic Applications. To demonstrate the utility of the (BT)S directing group in complex molecule synthesis, two natural products, salvianolic acid F and (+)-salvianolic acid A were formally synthesized using the (BT)S-directed C–H olefination reaction as the key step following a strategy inspired by the carbonyl directed synthesis by Xuan (Scheme 12).³⁶ Through this approach, salvianolic acid F was

synthesized in 5 steps with a projected 19.2% overall yield from mercaptobenzothiazole, which represents the highest overall yield and lowest step count synthesis yet published.³⁷ (+)-Salvianolic acid A was also formally synthesized in 9 steps overall in 14.1% projected yield with the longest linear route being from the mercaptobenzothiazole representing 7 steps with a projected 16.6% yield. This synthesis of (+)-salvianolic acid A has a similar step-count and slightly elevated yield compared to the most recent, highest yielding synthesis yet published.³⁶

CONCLUSION

In conclusion, we have demonstrated the utility of a new weakly-coordinating heteroaryl thioether directing group, (BT)S, that can facilitate catalytic transformations that require conformational flexibility for integral steps, such as β -hydride elimination (Scheme 13). This new directing group, which can be readily introduced by nucleophilic substitution of a hydroxyl or bromo functional group, serves as a versatile functional group for downstream manipulation. We probed each of the two reactions by performing RPKA and DFT, which helped to elucidate aspects of the mechanism for both catalytic cycles. We also demonstrated the utility of the (BT)S directing group by achieving high overall yields in the synthesis of two natural products.



Scheme 13. Illustration of the Selectivity of (BT)S

ASSOCIATED CONTENT

AUTHOR INFORMATION

Corresponding Author

* Email: Keary@Scripps.edu

Present Addresses

†If an author's address is different than the one given in the affiliation line, this information may be included here.

Author Contributions

The manuscript was written through contributions of all authors. / All authors have given approval to the final version of the manuscript. /

‡These authors contributed equally. (match statement to author names with a symbol)

Notes

The authors declare no competing financial interest.

Supporting Information

Experiment details, spectra data, copies of NMR spectra, X-ray crystallographic data, and computational details. These materials are available free of charge via the Internet at <http://pubs.acs.org>.

ACKNOWLEDGMENT

This work was financially supported by The Scripps Research Institute (TSRI) and the National Institutes of Health (SR35GM125052-02). We further acknowledge the NSF for a Graduate Research Fellowship (NSF/DGE-1842471, A.M.R.). We thank Prof. Ryan Shenvi (TSRI) and Tucker Huffman (Shenvi Lab, TSRI) for helpful discussions and Prof. Donna G. Blackmond (TSRI) for guidance with RPKA experiments. We also thank Prof. Arnold L. Rheingold, Dr. Milan Gembicky, and Dr. Curtis E. Moore for X-ray crystallographic analysis (UCSD). Brittany Sanchez (ASF, TSRI) is acknowledged for HPLC and HRMS analysis. We thank John A. Gurak, Jr. (Engle Lab, TSRI) and Zhen Liu (Engle Lab, TSRI) for their assistance of proofreading the manuscript. We also thank Zichen Wang (Engle Lab, TSRI) for his preliminary exploration of directed oxidative Heck reactions. We further thank Umicore for their generous donation of the Z-selective olefin metathesis catalyst Grubbs C675.

ABBREVIATIONS

(BT)S, Benzothiazole thioether; RPKA, Reaction Progress Kinetic Analysis; BQ, benzoquinone; PTFE, Polytetrafluoroethylene; LCMS, Liquid Chromatography—Mass Spectrometry; HPLC, High Pressure Liquid Chromatography.

REFERENCES

- (a) Brown, J. M.; Directed Homogeneous Hydrogenation [New Synthetic Methods (65)]. *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 190–203. (b) Hoveyda, A. H.; Evans, D. A.; Fu, G. C.; Substrate-Directable Chemical Reactions. *Chem. Rev.* **1993**, *93*, 1307–1370. (c) Breit, B.; Controlling Stereoselectivity with the Aid of a Reagent-Directing Group: Hydroformylation, Cuprate Addition, and Domino Reaction Sequences. *Chem. Eur. J.* **2000**, *6*, 1519–1524. (d) Oestreich, M.; Neighbouring-Group Effects in Heck Reactions. *Eur. J. Org. Chem.* **2005**, *2005*, 783–792. (e) Gurak, J. A.; Engle, K. M.; Regioselective Hydroamination Using a Directed Nucleopalladation/Protodepalladation Strategy. *Synlett* **2017**, *28*, 2057–2065.
- (a) McDonald, R. I.; Liu, G.; Stahl, S. S.; Palladium(II)-Catalyzed Alkene Functionalization via Nucleopalladation: Stereochemical Pathways and Enantioselective Catalytic Applications. *Chem. Rev.* **2011**, *111*, 2981–3019. (b) Jensen, K. H.; Sigman, M. S.; Mechanistic Approaches to Palladium-Catalyzed Alkene Difunctionalization Reactions. *Org. Biomol. Chem.* **2008**, *6*, 4083–4088. (c) Lee, A. L.; Enantioselective Oxidative Boron Heck Reactions. *Org. Biomol. Chem.* **2016**, *14*, 5357–5366. (d) Sigman, M. S.; Werner, E. W.; Imparting Catalyst Control upon Classical Palladium-Catalyzed Alkenyl C–H Bond Functionalization Reactions. *Acc. Chem. Res.* **2012**, *45*, 874–884. (e) Heck, R. F.; Palladium-Catalyzed Reactions of Organic Halides with Olefins. *Acc. Chem. Res.* **1979**, *12*, 146–151. (f) Dounay, A. B.; Overman, L. E.; The Asymmetric Intramolecular Heck Reaction in Natural Product Total Synthesis. *Chem. Rev.* **2003**, *103*, 2945–2964.
- (a) Gurak, J. A.; Yang, K. S.; Liu, Z.; Engle, K. M.; Directed, Regiocontrolled Hydroamination of Unactivated Alkenes via Protodepalladation. *J. Am. Chem. Soc.* **2016**, *138*, 5805–5808. (b) Yang, K. S.; Gurak, J. A.; Liu, Z.; Engle, K. M.; Catalytic, Regioselective Hydrocarbofunctionalization of Unactivated Alkenes with Diverse C–H Nucleophiles. *J. Am. Chem. Soc.* **2016**, *138*, 14705–14712. (c) Liu, Z.; Zeng, T.; Yang, K. S.; Engle, K. M.; β,γ -Vicinal Dicarbofunctionalization of Alkenyl Carbonyl Compounds via Directed Nucleopalladation. *J. Am. Chem. Soc.* **2016**, *138*, 15122–15125. (d) Liu, Z.; Wang, Y.; Wang, Z.; Zeng, T.; Liu, P.; Engle, K. M.; Catalytic Intermolecular Carboamination of Unactivated Alkenes via Directed Aminopalladation. *J. Am. Chem. Soc.* **2017**, *139*, 11261–11270. (e) O'Duill, M. L.; Matsuura, R.; Wang, Y.; Turnbull, J. L.; Gurak, J. A.; Gao, D.-W.; Lu, G.; Liu, P.; Engle, K. M.; Tridentate Directing Groups Stabilize 6-Membered Palladacycles in Catalytic Alkene Hydrofunctionalization. *J. Am. Chem. Soc.* **2017**, *139*, 15576–15579. (f) Liu, Z.; Ni, H.-Q.; Zeng, T.; Engle, K. M.; Catalytic Carbo- and Aminoboration of Alkenyl Carbonyl Compounds via Five- and Six-Membered Palladacycles. *J. Am. Chem. Soc.* **2018**, *140*, 3223–3227. (g) Zeng, T.; Liu, Z.; Schmidt, M. A.; Eastgate, M. D.; Engle, K. M.; Directed, Palladium(II)-Catalyzed Intermolecular Aminohydroxylation of Alkenes Using a Mild Oxidation System. *Org. Lett.* **2018**, *20*, 3853–3857. (h) Matsuura, R.; Jenkins, T. C.; Hill, D. E.; Yang, K. S.; Gallego, G. M.; Yang, S.; He, M.; Wang, F.; Marsters, R. P.; McAlpine, I.; Engle, K. M.; Palladium(II)-Catalyzed γ -Selective Hydroarylation of Alkenyl Carbonyl Compounds with Arylboronic Acids. *Chem. Sci.* **2018**, *9*, 8363–8368.
- (a) Neufeldt, S. R.; Sanford, M. S.; Asymmetric Chiral Ligand-Directed Alkene Dioxygenation. *Org. Lett.* **2013**, *15*, 46–49. (b) Talbot, E. P. A.; Fernandes, T. de A.; McKenna, J. M.; Toste, F. D.; Asymmetric Palladium-Catalyzed Directed Intermolecular Fluoroarylation of Styrenes. *J. Am. Chem. Soc.* **2014**, *136*, 4101–4104. (c) Wang, H.; Bai, Z.; Jiao, T.; Deng, Z.; Tong, H.; He, G.; Peng, Q.; Chen, G.; Palladium-Catalyzed Amide-Directed Enantioselective Hydrocarbofunctionalization of Unactivated Alkenes Using a Chiral Monodentate Oxazoline Ligand. *J. Am. Chem. Soc.* **2018**, *140*, 3542–3546. (d) Wang, C.; Xiao, G.; Guo, T.; Ding, Y.; Wu, X.; Loh, T.-P.; Palladium-Catalyzed Regiocontrollable Reductive Heck Reaction of Unactivated Aliphatic Alkenes. *J. Am. Chem. Soc.* **2018**, *140*, 9332–9336.
- Delcamp, J. H.; Brucks, A. P.; White, M. C.; A General and Highly Selective Chelate-Controlled Intermolecular Oxidative Heck Reaction. *J. Am. Chem. Soc.* **2008**, *130*, 11270–11271.
- (a) Larhed, M.; Carl-Magnus, A.; Hallberg, A.; Chelation-Controlled, Palladium-Catalyzed Arylation of Vinyl Ethers. *Acta Chem. Scand.* **1993**, *47*, 212–217. (b) Larhed, M.; Andersson, C.-M.; Hallberg, A.; Chelation-Controlled, Palladium-Catalyzed Arylation of Enol Ethers with Aryl Triflates. Ligand Control of Selection for α - or β -Arylation of [2-(Dimethylamino)ethoxy]ethene. *Tetrahedron* **1994**, *50*, 285–304. (c) Olofsson, K.; Sahlin, H.; Larhed, M.; Hallberg, A.; Regioselective Palladium-Catalyzed Synthesis of β -Arylated Primary Allylamine Equivalents by an Efficient Pd–N Coordination. *J. Org. Chem.* **2001**, *66*, 544–549. (d) Nilsson, P.; Larhed, M.; Hallberg, A.; A

New Highly Asymmetric Chelation-Controlled Heck Arylation. *J. Am. Chem. Soc.* **2003**, *125*, 3430–3431.

7. (a) Buezo, N. D.; Alonso, I.; Carretero, J. C.; Sulfinyl Group as a Novel Chiral Auxiliary in Asymmetric Heck Reactions. *J. Am. Chem. Soc.* **1998**, *120*, 7129–7130. (b) Buezo, N. D.; Mancheño, O. G.; Carretero, J. C.; The 2-(N,N-Dimethylamino)phenylsulfinyl Group as an Efficient Chiral Auxiliary in Intramolecular Heck Reactions. *Org. Lett.* **2000**, *2*, 1451–1454. (c) de la Rosa, J. C.; Díaz, N.; Carretero, J. C.; Asymmetric Synthesis of 1-Aryl and 1,3-Diarylcyclopentenes by the Heck Reaction of 1-Sulfinylcyclopentenes with Iodoarenes. *Tetrahedron Lett.* **2000**, *41*, 4107–4111. (d) Díaz Buezo, N.; de la Rosa, J. C.; Priego, J.; Alonso, I.; Carretero, J. C.; Sulfoxides as Stereochemical Controllers in Intermolecular Heck Reactions. *Chem. Eur. J.* **2001**, *7*, 3890–3900. (e) Alonso, I.; Carretero, J. C.; Highly Stereoselective Synthesis of Trisubstituted α,β -Unsaturated Sulfoxides by Heck Reaction. *J. Org. Chem.* **2001**, *66*, 4453–4456.

8. (a) Itami, K.; Mitsudo, K.; Kamei, T.; Koike, T.; Nokami, T.; Yoshida, J.; Highly Efficient Carbopalladation Across Vinylsilane: Dual Role of the 2-PyMe₂Si Group as a Directing Group and as a Phase Tag. *J. Am. Chem. Soc.* **2000**, *122*, 12013–12014. (b) Itami, K.; Nokami, T.; Ishimura, Y.; Mitsudo, K.; Kamei, T.; Yoshida, J.; Diversity-Oriented Synthesis of Multisubstituted Olefins through the Sequential Integration of Palladium-Catalyzed Cross-Coupling Reactions. 2-Pyridyldimethyl(vinyl)silane as a Versatile Platform for Olefin Synthesis. *J. Am. Chem. Soc.* **2001**, *123*, 11577–11585. (c) Itami, K.; Ushioji, Y.; Nokami, T.; Ohashi, Y.; Yoshida, J.; Stereoselective Synthesis of Multisubstituted Butadienes through Directed Mizoroki–Heck Reaction and Homocoupling Reaction of Vinyl(2-pyridyl)silane. *Org. Lett.* **2004**, *6*, 3695–3698. (d) Itami, K.; Mineno, M.; Muraoka, N.; Yoshida, J.; Sequential Assembly Strategy for Tetrasubstituted Olefin Synthesis Using Vinyl 2-Pyrimidyl Sulfide as a Platform. *J. Am. Chem. Soc.* **2004**, *126*, 11778–11779. (e) Bernocchi, E.; Cacchi, S.; Ciattini, P. G.; Morera, E.; Ortá, G.; Palladium-Catalyzed Vinylation of Allylic Alcohols with Enol Triflates. A Convenient Synthesis of Conjugated Dienols. *Tetrahedron Lett.* **1992**, *33*, 3073–3076. (f) Filippini, L.; Gusmeroli, M.; Riva, R.; Palladium-Catalyzed Coupling of Organic Halides and Tertiary Allylic Amines. *Tetrahedron Lett.* **1993**, *34*, 1643–1646. (g) Badone, D.; Guzzi, U.; Palladium-Catalyzed β -Arylation of Modified Vinyl Ethers with Aryl Triflates. *Tetrahedron Lett.* **1993**, *34*, 3603–3606. (h) Ono, K.; Fugami, K.; Tanaka, S.; Tamaru, Y.; Palladium Catalyzed Arylation of N-alkyl O-allyl Carbamates: Synthesis of Cinnamyl Alcohols via Heck Arylation. *Tetrahedron Lett.* **1994**, *35*, 4133–4136. (i) Kang, S.-K.; Jung, K.-Y.; Park, C.-H.; Namkoong, E.-Y.; Kim, T.-H.; Palladium-Catalyzed Arylation of Allylic Diols: Highly Selective Synthesis of Phenyl-Substituted Allylic Diols. *Tetrahedron Lett.* **1995**, *36*, 6287–6290. (j) Kang, S.-K.; Lee, H.-W.; Jang, S.-B.; Kim, T.-H.; Pyun, S.-J.; Complete Regioselection in Palladium-Catalyzed Arylation and Alkenylation of Allylic Alcohols with Hypervalent Iodonium Salts. *J. Org. Chem.* **1996**, *61*, 2604–2605. (k) Oestreich, M.; Sempere-Culler, F.; Machotta, A. B.; Catalytic Desymmetrizing Intramolecular Heck Reaction: Evidence for an Unusual Hydroxy-Directed Migratory Insertion. *Angew. Chem. Int. Ed.* **2004**, *44*, 149–152.

9. (a) Werner, E. W.; Mei, T.-S.; Burckle, A. J.; Sigman, M. S.; Enantioselective Heck Arylations of Acyclic Alkenyl Alcohols Using a Redox-Relay Strategy. *Science* **2012**, *338*, 1455–1458. (b) Mei, T.-S.; Werner, E. W.; Burckle, A. J.; Sigman, M. S.; Enantioselective Redox-Relay Oxidative Heck Arylations of Acyclic Alkenyl Alcohols using Boronic Acids. *J. Am. Chem. Soc.* **2013**, *135*, 6830–6833. (c) Mei, T.-S.; Patel, H. H.; Sigman, M. S.; Enantioselective Construction of Remote Quaternary Stereocentres. *Nature* **2014**, *508*, 340–344. (d) Xu, L.;

Hilton, M. J.; Zhang, X.; Norrby, P.-O.; Wu, Y.-D.; Sigman, M. S.; Wiest, O.; Mechanism, Reactivity, and Selectivity in Palladium-Catalyzed Redox-Relay Heck Arylations of Alkenyl Alcohols. *J. Am. Chem. Soc.* **2014**, *136*, 1960–1967. (e) Zhang, C.; Santiago, C. B.; Kou, L.; Sigman, M. S.; Alkenyl Carbonyl Derivatives in Enantioselective Redox Relay Heck Reactions: Accessing α,β -Unsaturated Systems. *J. Am. Chem. Soc.* **2015**, *137*, 7290–7293. (f) Larock, R. C.; Leung, W.-Y.; Stolz-Dunn, S.; Synthesis of Aryl-Substituted Aldehydes and Ketones via Palladium-Catalyzed Coupling of Aryl Halides and Non-Allylic Unsaturated Alcohols. *Tetrahedron Lett.* **1989**, *30*, 6629–6632. (g) Berthiol, F.; Doucet, H.; Santelli, M.; Heck Reaction of Aryl Bromides with Pent-4-en-2-ol, 2-Phenylpent-4-en-2-ol, or Hept-6-en-3-ol Catalysed by a Palladium-Tetraphosphine Complex. *Synthesis* **2005**, *37*, 3589–3602.

10. Huffman, T. R.; Wu, Y.; Emmerich, A.; Shenvi, R. A.; Intermolecular Heck Coupling with Hindered Alkenes Directed by Potassium Carboxylates. *Angew. Chem. Int. Ed.* **2019**, *58*, 2371–2376.

11. For representative examples, see: (a) Perales, J. B.; Van Vranken, D. L.; Thioether-Directed Platinum-Catalyzed Hydrosilylation of Olefins. *J. Org. Chem.* **2001**, *66*, 7270–7274. (b) Dupont, J.; Basso, N. R.; Meneghetti, M. R.; Chloropalladation of Propargyl Thioethers: A Facile Synthesis of Cyclopalladated Compounds. *Polyhedron* **1996**, *15*, 2299–2302. (c) Rao, G. K.; Kumar, A.; Kumar, S.; Dupare, U. B.; Singh, A. K.; Palladacycles of Thioethers Catalyzing Suzuki–Miyaura C–C Coupling: Generation and Catalytic Activity of Nanoparticles. *Organometallics* **2013**, *32*, 2452–2458. (d) Mann, S. E.; Aliev, A. E.; Tizzard, G. J.; Sheppard, T. D.; Sulfur-Directed Olefin Oxidations: Observation of Divergent Reaction Mechanisms in the Palladium-Mediated Acetoxylation of Unsaturated Thioacetals. *Organometallics* **2011**, *30*, 1772–1775.

12. For representative examples, see: (a) Yao, J.; Yu, M.; Zhang, Y.; Thioethers as Directing Group for the Palladium-Catalyzed Direct Arylation of Arenes. *Adv. Synth. Catal.* **2012**, *354*, 3205–3210. (b) Wang, B.; Lin, C.; Liu, Y.; Fan, Z.; Liu, Z.; Zhang, Y.; Thioether-Directed Acetoxylation of C(sp²)–H Bonds of Arenes by Palladium Catalysis. *Org. Chem. Front.* **2015**, *2*, 973–977. (c) Likun, J.; Wang, J.; Dong, G.; Palladium-Catalyzed γ -C(sp³)–H Arylation of Thiols via a Detachable Protecting/Directing Group. *Angew. Chem. Int. Ed.* **2018**, *57*, 12352–12355; for related example with platinum, see: (d) Perales, J. B.; Van Vranken, D. L.; Thioether-Directed Platinum-Catalyzed Hydrosilylation of Olefins. *J. Org. Chem.* **2001**, *66*, 7270–7274.

13. (a) Zheng, Y.; Song, W.; Zhu, Y.; Wei, B.; Xuan, L.; Pd-Catalyzed Acetoxylation of γ -C(sp³)–H Bonds of Amines Directed by a Removable Bts-Protecting Group. *J. Org. Chem.* **2018**, *83*, 2448–2454. (b) Petrova, E.; Rasina, D.; Jirgensons, A.; N-Sulfonylcarboxamide as an Oxidizing Directing Group for Ruthenium-Catalyzed C–H Activation/Annulation. *Eur. J. Org. Chem.* **2017**, *2017*, 1773–1779.

14. (a) Blackmond, D. G.; Reaction Progress Kinetic Analysis: A Powerful Methodology for Mechanistic Studies of Complex Catalytic Reactions. *Angew. Chem. Int. Ed.* **2005**, *44*, 4302–4320. (b) Blackmond, D. G.; Kinetic Profiling of Catalytic Organic Reactions as a Mechanistic Tool. *J. Am. Chem. Soc.* **2015**, *137*, 10852–10866.

15. Burés, J.; A Simple Graphical Method to Determine the Order in Catalyst. *Angew. Chem. Int. Ed. Engl.* **2016**, *55*, 2028–2031.

16. (a) Al-Hayaly, L. A. J.; Buttrus, N. H.; Al-Allaf, T. A. K.; Homobimetallic Complexes of Platinum(II) Group Metals with Sulphur Containing Ligands R₂C = CR₂ (R = 5-Phenyl-1,3,4-oxadiazole-2-thiol; 4,5-Diphenyl-1,2,4-triazole-3-thiol or Mercaptobenzothiazole). *Asian J. Chem.* **2002**, *14*, 1421–1426. (b) Song, R.-F.; Liu, J.-H.; Qian, H.; Zhao, K.-Y.; Dichloro[8-(1,3-benzothiazol-2-ylsulfanylmethyl)-quinoline-jN,N00]palladium(II). *Acta Crystallogr. Sect. E: Struct. Rep. Online* **2005**, *61*, 2142–2144.

(c) Pornsuriyasak, P.; Gangadharmath, U. B.; Rath, N. P.; Demchenko, A. V.; A Novel Strategy for Oligosaccharide Synthesis via Temporarily Deactivated S-Thiazolyl Glycosides as Glycosyl Acceptors. *Org. Lett.* **2004**, *6*, 4515–4518. (d) Li, Y.-W.; Gu, G.-B.; Liu, H.-Y.; Sung, H. H. Y.; Williams, I. D.; Chang, C.-K.; A New iso-Amyl Benzothiazolyl Sulfoxide as an Extractant for Palladium and the Crystal Structure of Its Palladium (II) Complex. *Molecules* **2005**, *10*, 912–921. (e) Hughes, M. N.; Rutt, K. J.; The Far Infrared Spectra of Some Complexes of Palladium (II) with Thiazoles. *Spectrochim. Acta, Pt. A: Mol. Spectrosc.* **1971**, *27*, 924–926. (f) Fuchita, Y.; Yoshinaga, K.; Kusaba, H.; Mori, M.; Hiraki, K.; Takehara, K.; Synthesis and Characterisation of the Six-Membered Cyclopalladated Complexes of 2-Benzylbenzothiazole. *Inorg. Chim. Acta* **1995**, *239*, 125–132.

17. For coordinates of the transition states, see SI in: (a) Liu, M.; Yang, P.; Karunananda, M. K.; Wang, Y.; Liu, P.; Engle, K. M.; C(alkenyl)-H Activation via Six-Membered Palladacycles: Catalytic 1,3-Diene Synthesis. *J. Am. Chem. Soc.* **2018**, *140*, 5805–5815.

18. Carbon-carbon unsaturated bonds have also been utilized as directing groups for arene C–H olefination, but the scope and utility of these transformations tend to be limited due to the difficulty of installing the directing group, reduction of the directing group during the course of the reaction, and the inability to remove or controllably transform after the C–H functionalization reaction. For a review, see: Minami, Y.; Hiyama, T.; Collaborative Interaction of Carbon–Carbon Unsaturated Bond Groups with Transition-Metal Catalysts for C–H Bond Functionalization. *Chem. Lett.* **2018**, *47*, 1–8.

19. Mei, T.-S.; Wang, X.; Yu, J.-Q.; Pd(II)-Catalyzed Amination of C–H Bonds Using Single-Electron or Two-Electron Oxidants. *J. Am. Chem. Soc.* **2009**, *131*, 10806–10807.

20. Wasa, M.; Yu, J.-Q.; Synthesis of β -, γ -, and δ -Lactams via Pd(II)-Catalyzed C–H Activation Reactions. *J. Am. Chem. Soc.* **2008**, *130*, 14058–14059.

21. Wang, X.; Lu, Y.; Dai, H.-X.; Yu, J.-Q.; Pd(II)-Catalyzed Hydroxyl-Directed C–H Activation/C–O Cyclization: Expedient Construction of Dihydrobenzofurans. *J. Am. Chem. Soc.* **2010**, *132*, 12203–12205.

22. Mei, T.-S.; Wang, D.-H.; Yu, J.-Q.; Expedient Drug Synthesis and Diversification via ortho-C–H Iodination using Recyclable Pd₂ as the Precatalyst. *Org. Lett.* **2010**, *12*, 3140–3143.

23. (a) Li, G.; Leow, D.; Wan, L.; Yu, J.-Q.; Ether-Directed ortho-C–H Olefination with a Palladium(II)/Monoprotected Amino Acid Catalyst. *Angew. Chem. Int. Ed.* **2013**, *52*, 1245–1247. (b) Li, G.; Wan, L.; Zhang, G.; Leow, D.; Spangler, J.; Yu, J.-Q.; Pd(II)-Catalyzed C–H Functionalizations Directed by Distal Weakly Coordinating Functional Groups. *J. Am. Chem. Soc.* **2015**, *137*, 4391–4397.

24. Yu, M.; Xie, Y.; Xie, C.; Zhang, Y.; Palladium-Catalyzed C–H Alkenylation of Arenes Using Thioethers as Directing Groups. *Org. Lett.* **2012**, *14*, 2164–2167.

25. Wang, B.; Shen, C.; Yao, J.; Yin, H.; Zhang, Y.; Palladium(II)-Catalyzed ortho-Olefination of Arenes Applying Sulfoxides as Remote Directing Groups. *Org. Lett.* **2014**, *16*, 46–49.

26. Maji, A.; Bhaskararao, B.; Singha, S.; Sunoj, R. B.; Maiti, D.; Directing Group Assisted Meta-Hydroxylation by C–H Activation. *Chem. Sci.* **2016**, *7*, 3147–3153.

27. Engle, K. M.; The Mechanism of Palladium(II)-Mediated C–H Cleavage with Mono-N-Protected Amino Acid (MPAA) Ligands: Origins of Rate Acceleration. *Pure Appl. Chem.* **2016**, *88*, 119–138.

28. For previous examples of ethenesulfonyl fluoride (ESF) in C–H olefination, see: (a) Wang, S.-M.; Li, C.; Leng, J.; Bukhari, S. N.

A.; Qin, H.-L.; Rhodium(III)-Catalyzed Oxidative Coupling of N-Methoxybenzamides and Ethenesulfonyl fluoride: a C–H Bond Activation Strategy for the Preparation of 2-Aryl Ethenesulfonyl Fluorides and Sulfonyl Fluoride Substituted γ -Lactams. *Org. Chem. Front.* **2018**, *5*, 1411–1415. (b) Liu, M.; Yang, P.; Karunananda, M. K.; Wang, Y.; Liu, P.; Engle, K. M.; C(alkenyl)-H Activation via Six-Membered Palladacycles: Catalytic 1,3-Diene Synthesis. *J. Am. Chem. Soc.* **2018**, *140*, 5805–5813. (c) Li, C.; Wang, S.-M.; Qin, H.-L.; A Rh-Catalyzed Air and Moisture Tolerable Aldehyde (Ketone)-Directed Fluorosulfonylvinylolation of Aryl C(sp²)-H Bonds. *Org. Lett.* **2018**, *20*, 4699–4703. (d) Wang, S.-M.; Moku, B.; Leng, J.; Qin, H.-L.; Rh-Catalyzed Carboxylates Directed C–H Activation for the Synthesis of ortho-Carboxylic 2-Arylethenesulfonyl Fluorides: Access to Unique Electrophiles for SuFEx Click Chemistry. *Eur. J. Org. Chem.* **2018**, *2018*, 4407–4410. (e) Ncube, G.; Huestis, M. P.; Directed Cp*Rh^{III}-Catalyzed Fluorosulfonylvinylolation of Arenes. *Organometallics* **2019**, *38*, 76–80. (f) Chen, X.-Y.; Wu, Y.; Zhou, J.; Wang, P.; Yu, J.-Q.; Synthesis of β -Arylethenesulfonyl Fluoride via Pd-Catalyzed Nondirected C–H Alkenylation. *Org. Lett.* **2019**, *21*, 1426–1429.

29. (a) Engle, K. M.; Wang, D.-H.; Yu, J.-Q.; Ligand-Accelerated C–H Activation Reactions: Evidence for a Switch of Mechanism. *J. Am. Chem. Soc.* **2010**, *132*, 14137–14151. (b) Deb, A.; Bag, S.; Kancherla, R.; Maiti, D.; Palladium-Catalyzed Aryl C–H Olefination with Unactivated, Aliphatic Alkenes. *J. Am. Chem. Soc.* **2014**, *136*, 13602–13605.

30. Otsuka, S.; Nogi, K.; Yorimitsu, H.; Palladium-Catalyzed Insertion of Isocyanides into the C–S Bonds of Heteroaryl Sulfides. *Angew. Chem. Int. Ed.* **2018**, *57*, 6653–6657.

31. Baudin J. B.; Stereochemistry of Direct Olefin Formation from Carbonyl Compounds and Lithiated Heterocyclic Sulfones. *Bull. Soc. Chim. Fr.* **1993**, *130*, 856–878.

32. Day, J. J.; Neill, D. L.; Xu, S.; Xian, M.; Benzothiazole Sulfinic Acid: A Sulfinic Acid Transfer Reagent under Oxidation-Free Conditions. *Org. Lett.* **2017**, *19*, 3819–3822.

33. Modified Julia olefinations are known to exhibit variable *E/Z* selectivity as a function of reaction solvent, base, and coupling partner, and the high *E*-selectivity in this case is a result of the reaction conditions and the identity of the arylaldehyde coupling partner, as has been described previously. Blakemore, P. R.; The Modified Julia Olefination: Alkene Synthesis via the Condensation of Metallated Heteroarylalkylsulfones with Carbonyl Compounds. *J. Chem. Soc.; Perkin Trans. 1.* **2002**, 2563–2585.

34. Bon, D. J. Y. D.; Kováč, O.; Ferugová, V.; Zálešák, F.; Pospíšil, J.; One and Two-Carbon Homologation of Primary and Secondary Alcohols to Corresponding Carboxylic Esters Using β -Carbonyl BT Sulfones as a Common Intermediate. *J. Org. Chem.* **2018**, *83*, 4990–5001.

35. Zenzola, M.; Doran, R.; Degennaro, L.; Luisi, R.; Bull, J. A.; Transfer of Electrophilic NH Using Convenient Sources of Ammonia: Direct Synthesis of NH Sulfoximines from Sulfoxides. *Angew. Chem. Int. Ed.* **2016**, *55*, 7203–7207.

36. Zheng, Y.; Song, W.-B.; Xuan, L.-J.; The Asymmetric Total Synthesis of (+)-Salvianolic Acid A. *Tetrahedron* **2016**, *72*, 5047–5050.

37. Dalla, V.; Cotellet, P.; The Total Synthesis of Salvianolic Acid F. *Tetrahedron* **1999**, *55*, 6923–6930.

