

Reactions of *trans*-[IrCl(CO)(PPh₃)₂] and *trans*-[RhCl(CO)(PPh₃)₂] with Organic Tetrahalides – Oxidative Addition and Formation of Dihaloalkenes

IBRAHIM M. AL-NAJJAR

Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh-11451, Saudi Arabia


(Received October 7, 1986)

Oxidative addition reactions of the title compounds with 1,1,2,2-tetrachloroethane yield *trans*-[MCl₃(CO)(PPh₃)₂], (M = Rh(III), Ir(III)) and *cis*-1,2-dichloroethene. The stereochemistry of these complexes was assigned using IR and ³¹P{¹H} NMR spectroscopy, and the addition is stereospecific and *trans* in this particular case. Tetrachloroethane is used as a solvent in many reactions. In view of the finding now that it takes part in the reaction, care must be exercised in the use of such a solvent in similar reactions. Addition of MeI to the title compounds in tetrachloroethane gives a *trans* addition product with the Ir complex, and *cis*, *trans*-addition products with the Rh complex.

The reduction of organic halides by metals is a well-known route to olefins and cycloalkanes [1–3]. The reduction of organic tetrahalides by a metal is still not known.

In view of our interest in biheterometallic and platinum complexes [4–6], we have investigated the reaction between *trans*-[RhCl(CO)(PPh₃)₂] and *cis*-[PtCl₂(PPh₃)₂] in the presence of 1,1,2,2-tetrachloroethane as a good solvent recommended for the preparation of binuclear platinum complexes of the type [Pt₂X₄(PR₃)₂] [7]. By examining the resulting solution of the reaction by IR and ³¹P NMR spectroscopy, we found that, there is no change in the platinum complex spectrum but a dramatic change in the rhodium spectrum. However, the above reaction was repeated in the presence of RhCl(CO)(PPh₃)₂ and tetrachloroethane only, under N₂ and refluxing. The resulting solution was examined by IR, ¹H and ³¹P{¹H} NMR spectroscopy. The results confirm the formation of *mer,trans*-[RhCl₃(CO)(PPh₃)₂] and *cis*-dichloroethene, Scheme 1.

The complex *trans*-[RhCl(CO)(PPh₃)₂] (500 mg) was dissolved in tetrachloroethane (10 ml). The solution was stirred under nitrogen and refluxed for 1–2 h, and the solvent removed *in vacuo*. The yellow solids were dissolved in CDCl₃ and Et₂O was added to precipitate fine, yellow crystals (Et₂O may be added to the original solution to obtain yellow crystals). The vapours from the reaction were cooled



Scheme 1.

through a condenser and then passed through cold CDCl₃ solution. The IR spectra of the yellow solid exhibit a strong band at 2100 cm⁻¹, assigned for the new ν (CO) group of the Rh(III) complex, and a weak band at 1960 cm⁻¹, assigned for ν (CO) of the starting Rh(I) complex (compare with the authentic sample). The IR band at 331 cm⁻¹ assigned for ν (Rh–Cl) (*trans* to CO) and a strong band at 345 cm⁻¹ assigned for RhCl₂ (*trans* Cl), asymmetric stretching vibration. The IR spectra of the new complex are in good agreement with the analogous rhodium complex *trans*-[RhCl₃(CO)(PMe₃)₂] [8]. Further strong evidence was obtained from the CDCl₃ solution of the new complex. The ³¹P{¹H} NMR spectrum consists of a doublet at δ +8.9 and ¹J(¹⁰³Rh–³¹P), 78.1 Hz and small broad doublets appear at δ +29.0 and ¹J(¹⁰³Rh–³¹P), 127.0 Hz assigned for the starting complex (compare with authentic sample). Coupling constant values are comparable to the values of ¹J(¹⁰³Rh–³¹P) obtained from *trans*-[RhCl₃(CO)(PMe₃)₂], ¹J, 72 Hz and for *trans*-[RhCl(CO)(PMe₃)₂], ¹J, 114 Hz [8]. The ¹H and ¹³C NMR spectra of the CDCl₃ solution of the resulting condensed vapour showed a significant quantity of dichloroethene. The ¹H NMR spectrum showed a single peak which appears at δ 6.45 ppm and ¹³C NMR spectra showed a single peak at δ 116.6 ppm. These results are comparable to the authentic sample of *cis*- and *trans*-dichloroethene (¹H, δ 6.47, 6.36 and ¹³C δ , 117.0, 120.0 for *cis* and *trans* respectively in CDCl₃ and also fit well with the reported values [9, 10]).

Accordingly, the olefinic product is more likely to be *cis*-1,2-dichloroethene. This probably is formed as a result of decomposition of Rh(III)-alkyl intermediates; similar intermediates were obtained in the Rh(I) macrocycle reaction with organic dihalides [11]. The above reaction does not work at room

temperature. The case is different when Vaska's complex, *trans*-[IrCl(CO)(PPh₃)₂] (100 mg), is treated with tetrachloroethane (10 ml) at room temperature (the reaction was completed in 24 h) or refluxed under nitrogen for a few minutes. The fine yellow crystals, obtained as before, showed IR and ³¹P NMR spectra for a new Ir(III) complex, *trans*-[IrCl₃(CO)(PPh₃)₂], and the formation of *cis*-1,2-dichloroethene. The ³¹P{¹H} NMR spectrum showed a sharp single peak at δ -17.5 (no traces of the starting Ir(I) complex were observed). The ³¹P NMR spectrum of the starting Ir(I) complex shows a broad peak at δ +23.8 (obtained separately). The IR spectrum of the newly Ir(III) complex, showed a strong peak at 2095 cm⁻¹, assigned for ν (CO), and a characteristic peak near 301 cm⁻¹ assigned for Ir-Cl (*trans* to CO). As a result, reaction of either complex, Ir(I) or Rh(I), with tetrachloroethane gives only one isomer, thus the addition is stereospecific and *trans* in this particular case. The above reactions provide the first examples of the formation of Rh(III) and Ir(III) complexes and elimination of *cis*-1,2-dichloroethene.

We also investigated the reactions of MeI with the title compounds in tetrachloroethane, at room temperature. Examination of the ³¹P{¹H} NMR spectrum of the new Rh(III) products (completed in 48 h in the presence of excess tetrachloroethane) clearly indicates the formation of two new complexes which appear as two doublets, at δ +23.6, ¹J(¹⁰³Rh-³¹P), 107.4 Hz, and at δ +12.3, ¹J(¹⁰³Rh-³¹P), 90.3 Hz, assigned for the *trans* and *cis* addition of MeI (B and C, Scheme 1). In the iridium(I) complex, only one species is formed, assigned for the *trans* addition species (D, Scheme 1); ³¹P{¹H} NMR shows only one single peak at δ -19.1. This behaviour is in contrast to the addition of MeI to the analogous *trans*-[IrCl(CO)(PPh₂Me)₂] complex in methanol which gives both *cis* and *trans* addition products (similar to B and C, Scheme 1) [12, 13].

All NMR spectra were measured on a JEOL JNM FX-100 spectrometer operating in the Fourier transform mode. Infrared spectra were measured on a Beckman 4240 spectrophotometer as KBr pellets.

Acknowledgements

This research (Chem/1406/41) was supported by the Research Centre, College of Science, King Saud University, Riyadh, Saudi Arabia. I would like to thank Prof. G. A. Miana for reading the manuscript. I wish also to thank Mr. H. M. Abdullah for assistance with the synthesis and Mr. M. Gopal for running the NMR spectra.

References

- 1 H. O. House, 'Modern Synthetic Reactions', 2nd edn., W. A. Benjamin, Menlo Park, Calif., 1972, p. 220.
- 2 D. M. Singleton and J. K. Kochi, *J. Am. Chem. Soc.*, **89**, 6547 (1967) and refs. therein.
- 3 J. K. Kochi and D. M. Singleton, *J. Org. Chem.*, **33**, 1027 (1968) and refs. therein.
- 4 J. K. Sarhan, M. Green and I. M. Al-Najjar, *J. Chem. Soc., Dalton Trans.*, 771 (1984).
- 5 J. K. Sarhan, M. Green and I. M. Al-Najjar, *Organometallics*, **3**, 520 (1984).
- 6 I. M. Al-Najjar, *Inorg. Chim. Acta*, (1986) submitted for publication.
- 7 A. C. Smithies, M. Rycheck and M. Orchin, *J. Organomet. Chem.*, **12**, 199 (1968).
- 8 J. Browning, P. L. Goggin, R. J. Goodfellow, M. G. Norton, A. J. M. Rattray and B. F. Taylor, *J. Chem. Soc., Dalton Trans.*, 2061 (1977).
- 9 C. Pascual, J. Meier and W. Simson, *Helv. Chim. Acta*, **49**, 164 (1966).
- 10 G. Miyazima, Y. Utsumi and K. Takahashi, *J. Phys. Chem.*, **73**, 1370 (1969).
- 11 J. P. Collman, J. I. Brauman and A. M. Madonik, *Organometallics*, **5**, 218 (1986).
- 12 A. J. Deming and B. L. Shaw, *J. Chem., Soc. A*, 1128 (1969).
- 13 J. P. Collman and C. T. Sears, Jr., *Inorg. Chem.*, **7**, 27 (1968).