

# Base-Induced Radical Carboamination of Nonactivated Alkenes with Aryldiazonium Salts

Stephanie Kindt, Karina Wicht, and Markus R. Heinrich\*

Department of Chemistry and Pharmacy, Pharmaceutical Chemistry, Friedrich-Alexander-Universität Erlangen-Nürnberg, Schuhstraße 19, 91052 Erlangen, Germany

**Supporting Information** 

**ABSTRACT:** A new transition-metal-free version of the Meerwein arylation has been developed. The key feature of this carboamination-type reaction is the slow base-controlled generation of aryl radicals from aryldiazonium tetrafluoroborates, so that a sufficient quantity of diazonium ions remains to enable efficient trapping of the alkyl radical adduct resulting from aryl radical addition to the alkene. Under strongly basic conditions, diazonhydrides are likely to take over the role of the nitrogen-centered radical scavengers.



The Meerwein arylation<sup>1</sup> has recently become a highly versatile tool for the radical functionalization of alkenes (Scheme 1, A). Besides the traditional diazonium salts ( $X = N_2^+$ ),



bromo- and iodobenzenes (X = Br, I) as well as arylhydrazines (X = NHNH<sub>2</sub>) have been successfully applied as aryl radical sources, and the group of alkenes, which was at first practically limited to activated alkenes (e.g., R = COOR', CN, Ph) could be extended to nonactivated alkenes. Moreover, a significant number of new radical scavengers have been found, which now allow the introduction of a broad variety of atoms or functional groups Y.<sup>2,3</sup>

Along with these expansions of the scope, the attractiveness of the Meerwein arylation was further increased through the development of transition-metal-free,<sup>4</sup> catalytic,<sup>5</sup> as well as photocatalytic versions.<sup>6,7</sup> Beautiful examples are the TEMPO– Na-mediated carboxygenation<sup>8</sup> developed by Studer<sup>9</sup> (Scheme 1, B) and the photocatalyzed carboamination reported by König<sup>10</sup> (Scheme 1, C). In this context, it is, however, surprising that the base-induced generation of aryl radicals from diazonium salts,<sup>11,12</sup> which is well established for Gomberg–Bachmann reactions,<sup>13–15</sup> has so far only rarely been exploited in alkene functionalizations.<sup>16,17</sup> Especially from an economical point of view, such base-mediated variants could be fully competitive with catalytic and photocatalytic versions.

Since carboaminations of alkenes is a highly topical field of research, with many metal-catalyzed versions having been reported recently,<sup>18</sup> we chose a Meerwein-type carboamination reaction as a starting point (Scheme 1, D). In this reaction, aryldiazonium salts play a double role and act as aryl radical sources as well as nitrogen-centered radical scavengers.<sup>19,20</sup> A major drawback of these versatile transformations so far, however, was that a large excess of either iron(II) sulfate or titanium(III) chloride<sup>21</sup> was required to obtain the desired azo compounds in good yields.<sup>22</sup>

In a series of optimization experiments aiming at a baseinduced radical carboamination reaction, solutions of 4chlorophenyldiazonium tetrafluoroborate (1a) were added to mixtures of 3-methyl-3-buten-1-ol (2a) and a base in aqueous acetonitrile under various conditions (Scheme 2; see the Supporting Information for details).<sup>23</sup> The best yield for the desired azo compound 3aa was obtained with sodium carbonate at a temperature of 50 °C. Pyrazolidinone 4aa, which was the most important byproduct of the reaction, was formed in largest quantity when the weaker base potassium acetate was used. By





Received: October 30, 2015



employing the conditions optimized for **3aa**, a scaling-up experiment was also successful in which **3aa** was obtained from **1a** and **2a** on a 10 mmol scale in 72% yield (1.04 g). In parallel, we evaluated the base-induced carboamination starting from the corresponding 4-chlorophenyldiazonium chloride (see the Supporting Information). The best yield for **3aa** (63%), along with 7% of **4aa**, was obtained with 4.5 equiv of Na<sub>2</sub>CO<sub>3</sub> in a 1:1 mixture of acetonitrile and water at 50 °C and at an identical addition time of 5 min. On this basis, the tetrafluoroborate-based version was chosen for an extensive evaluation of scope and limitations (Tables 1 and 2), and fewer reactions were carried out under those conditions optimized for the diazonium chlorides (see the Supporting Information).

Table 1. Evaluation of Substrate Scope and Limitations (I)				
R <sup>1</sup> N2 <sup>⊖</sup> + N2 <sup>⊖</sup> + 1a-t (slow add. over 5 min)	Na₂CO₃           (1.5 equiv)           50 °C           (12 equiv)	R <sup>1</sup> N <sup>N</sup> Jaa-3ta		
entry o	liazonium salt 1: R <sup>1</sup> =	yield <sup><math>a</math></sup> of $3b$ (%)		
1	1a: 4-Cl	<b>3aa</b> (83) <sup>c</sup>		
2	1b: 3-Cl	<b>3ba</b> (60) <sup><i>c</i></sup>		
3	1c: 2-Cl	<b>3ca</b> (73)		
4	1d: 3,4-Cl <sub>2</sub>	<b>3da</b> (78) <sup>c</sup>		
5	1e: 3,5-Cl <sub>2</sub>	<b>3ea</b> (48) <sup>c</sup>		
6	1f: 4-F	<b>3fa</b> (51) <sup>c</sup>		
7	1g: 2-F	3ga (62)		
8	1h: 4-Br	<b>3ha</b> (63)		
9	1i: 3-Br	<b>3ia</b> (63)		
10	1j: 2-Br	3ja (59)		
11	1k: 4-I	<b>3ka</b> (43)		
12	1l: 4-OMe	3la (30) <sup>d</sup>		
13	1m: 3,4-(OMe) <sub>2</sub>	<b>3ma</b> (51)		
14	<b>1n</b> : 4-CF <sub>3</sub>	<b>3na</b> (51) <sup><i>d</i></sup>		
15	10: 4-CN	<b>30a</b> (48) <sup>d</sup>		
16	1p: 3-CN	3pa (47) <sup>d</sup>		
17	1q: 4-CO <sub>2</sub> Me	3qa (50)		
18	1r: 4-COMe	<b>3ra</b> (40)		
19	<b>1s</b> : 4-NO <sub>2</sub>	<b>3sa</b> (41) <sup>c</sup>		
20	1t: 4-H	<b>3ta</b> (34)		

<sup>*a*</sup>Diazonium tetrafluoroborate 1 (1.0 mmol) in CH<sub>3</sub>CN/H<sub>2</sub>O (5:1, 4 mL) added over 5 min to a mixture of alkene 2 (12 mmol) and Na<sub>2</sub>CO<sub>3</sub> (1.5 mmol) in CH<sub>3</sub>CN/H<sub>2</sub>O (5:1, 5 mL) at 50 °C. <sup>*b*</sup>Yields after purification by column chromatography. <sup>*c*</sup>Byproduct pyrazolidinone 4 observed in low yield <8%. <sup>*d*</sup>Byproduct pyrazolidinone: 41a (23%), 4na (15%), 4oa (23%), 4pa (16%).

The evaluation of the substrate scope was started with different substitution patterns on the aryldiazonium salt (Table 1). On average, good yields were obtained for halogenated compounds 1a-k (entries 1-11). Donor- and acceptor-substituted diazonium salts 11-s were tolerated as well (entries 12-19). Within these groups, the largest amounts of the pyrazolidinones 4 were observed, with a maximum yield of 23% for 40a (entry 15). The comparably low yield of 34% obtained for the unsubstituted diazonium salt 1t (entry 20) was not completely surprising as a lack of substituents on the aromatic core had shown negative effects in earlier studies.<sup>15a</sup> Variation of the alkene in the second part of the study (Table 2) at first demonstrated that the



	$\begin{bmatrix} R_{1}^{1} & BF_{4}^{\Theta} \\ N_{2}^{\Theta} \end{bmatrix} + \begin{bmatrix} 1a,c,d,k & 2b \\ slow add. over 5 min \end{bmatrix}$ (12 e	$R^{2} \xrightarrow{(1.5 \text{ equiv})}_{(CH_{3}CN-H_{2}O)} R^{1}$	$N = N$ $R^{1}$ $N = N$ $R^{2}$ $R^{2$
entry	diazonium salt 1: $R^1 =$	alkene <b>2</b> : $R^2 =$	yield <sup><math>a</math></sup> of $3b$ (%)
1	1a: 4-Cl	<b>2b</b> : CH <sub>2</sub> OH	<b>3ab</b> (72)
2	1c: 2-Cl	<b>2b</b> : CH <sub>2</sub> OH	3cb (54)
3	1d: 3,4-Cl <sub>2</sub>	<b>2b</b> : CH <sub>2</sub> OH	3db (70)
4	1k: 4-I	<b>2b</b> : CH <sub>2</sub> OH	<b>3kb</b> (61)
5	1a: 4-Cl	<b>2c</b> : (CH <sub>2</sub> ) <sub>3</sub> C(Me) <sub>2</sub> OH	<b>3ac</b> (66)
6	1a: 4-Cl	<b>2d</b> : (CH <sub>2</sub> ) <sub>2</sub> OAc	<b>3ad</b> (46)
7	1a: 4-Cl	<b>2e</b> : (CH <sub>2</sub> )OAc	<b>3ae</b> (43)
8	1a: 4-Cl	<b>2f</b> : $(CH_2)_2COMe$	<b>3af</b> (71)
9	1a: 4-Cl	2g: CH <sub>2</sub> OC <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	<b>3ag</b> (56)
10	1a: 4-Cl	<b>2h</b> : (CH <sub>2</sub> )OTHP <sup>c</sup>	<b>3ah</b> (61)
11	1a: 4-Cl	<b>2i</b> : (CH <sub>2</sub> ) <sub>2</sub> CO <sub>2</sub> Me	3ai (65)
12	1a: 4-Cl	<b>2</b> j: (CH <sub>2</sub> ) <sub>2</sub> OCOPh	<b>3aj</b> (44)
13	1a: 4-Cl	<b>2k</b> : (CH <sub>2</sub> ) <sub>2</sub> N <sub>3</sub>	<b>3ak</b> (56)
Diazo	onium totrafluoroborat	a <b>1</b> (10  mmol) in CH	CN/H O (5.1 4

<sup>*a*</sup>Diazonium tetrafluoroborate **1** (1.0 mmol) in CH<sub>3</sub>CN/H<sub>2</sub>O (5:1, 4 mL) added over 5 min to a mixture of alkene **2** (12 mmol) and Na<sub>2</sub>CO<sub>3</sub> (1.5 mmol) in CH<sub>3</sub>CN/H<sub>2</sub>O (5:1, 5 mL) at 50 °C. <sup>*b*</sup>Yields after purification by column chromatography. <sup>*c*</sup>THP = tetrahydropyran-2-yl.

undesired pyrazolidinone formation is a particular feature of alcohol 2a (c.f. entries 1–5), which can be explained by the mechanism presented below (Scheme 3). Second, this series of experiments showed that functional groups other than alcohols, such as esters (entries 6, 7, 11, and 12), ketones (entry 8), phenyl ethers (entry 9), cyclic acetals (entry 10), and azides (entry 13), do not interfere with the chosen reaction conditions.

Additional attempts with 3-butenol demonstrated that the optimized reaction conditions are not equally well suited for monosubstituted alkenes, as the tetrafluoroborate salt **1a** and its corresponding chloride salt gave the related azo compound only in yields of 30% and 25%, respectively (see the Supporting Information). As a lower stability of an azo compound derived from a monosubstituted alkene under the reaction conditions and an isomerization to the corresponding hydrazone could be ruled out,<sup>24</sup> we assume at this point that the slower trapping of less nucleophilic secondary carbon-centered radicals by the diazonium ions is responsible for this trend.<sup>25</sup>

Under conditions identical to those applied in Tables 1 and 2, methylenecyclohex-2-yl acetate and 4-methylpent-3-en-2-one provided the azo compounds 3al (2:1 mixture of stereoisomers) and 3am, respectively (Figure 1). A first example for a basemediated Meerwein-type carbohydroxylation was found in the



Figure 1. Additional products.

в

reaction of 2-nitrophenyldiazonium tetrafluoroborate with alkene **2a**. This reaction gave the tertiary alcohol **5**, probably through a fast intramolecular alkyl radical attack onto the nitro group and subsequent oxidative ring-opening,<sup>26</sup> and no yield of the corresponding azo compound.<sup>27</sup> After unsuccessful attempts with sodium dithionite,<sup>28</sup> the reduction of **3aa** to amine **6** and 4-chloroaniline was achieved through hydrogenation over Raney nickel<sup>29</sup> in 90% yield. Regarding carboamination, this reduction thus allows recovery of the second aniline equivalent, which may then be reused for the reaction via diazotization.

Taking a closer look at the mechanism of the base-induced carboamination, the aryl radical generation is likely to proceed via the intermediates known from Gomberg–Bachmann reactions (Scheme 3).<sup>13b</sup> The diazonium ion 1 is stepwise converted to

## Scheme 3. Plausible Reaction Mechanisms



diazohydroxide 7, to diazotate 8, and then to diazoanhydride 9 before homolytic cleavage into an aryl radical 10, nitrogen, and diazenyloxyl radical 11 occurs.<sup>13,30</sup> Aryl radical addition to 2a leads to alkyl radical 12, which may be trapped by a diazonium ion to give at first radical cation 13 and then azo compound 14 via a final reductive step (path A).<sup>20a</sup> As the reaction can also be conducted under strongly basic conditions (see Table S1, Supporting Information), under which no free diazonium ions should be available for the trapping of 12, it appears plausible that 9 itself may act as radical scavenger to give a well-stabilized nitrogen-centered radical 15.<sup>31</sup> Fragmentation of 15 could then occur via either path B or C. The formation of pyrazolidinone 4 from 13 can be rationalized by an intramolecular electron transfer out of a suitably positioned and activated C–H bond to give the stabilized radical 16.<sup>31–33</sup> The intramolecular nature of this step is supported by the fact that the CH<sub>2</sub>-shortened alkene **2b** (c.f. Table 2) did not undergo a similar reaction. Later, **4** is formed from 16 via oxidation and cyclization.<sup>34</sup> An alternative mechanism leading to 4 via a Cannizarro-like hydride transfer can currently however not be ruled out.<sup>35</sup> Investigation of the byproducts suggested that the electrons required for the steps 11  $\rightarrow$  8 and 13  $\rightarrow$  14 come from an oxidation of the alkene (e.g., 2a) in allylic position.

To maximize the yield of pyrazolidinone 4,<sup>36</sup> we combined the most suitable base potassium acetate (Scheme 2) with the best diazonium salt 10 (Table 1, entry 15). This experiment indeed gave the pyrazolidinone 40a in 47% yield along with only 18% of 30a (see the Supporting Information). When the same reactants 10 and 2a were instead submitted to an iron(II)-mediated reaction,<sup>21b,c</sup> only the carboamination product 30a was obtained in 47% yield, thereby showing the critical importance of radical cation 13 for the formation of pyrazolidinones.

In summary, a new transition-metal-free Meerwein-type carboamination reaction has been developed, which relies on the generation of aryl radicals under basic conditions that were so far typical for Gomberg–Bachmann reactions. The product yields compare well and do sometimes even exceed those, which could so far only be reached by employing iron(II) sulfate as reductant in large excess.<sup>21d</sup> The variation of reaction conditions further gave hints that not only diazonium ions but also diazoanhydrides may act as nitrogen-centered radical scavengers. Most importantly, this study shows that base-mediated variants represent, besides recent catalytic and photocatalytic improvements of the Meerwein arylation, a third option to further increase the attractiveness of this powerful radical transformation.

### ASSOCIATED CONTENT

## Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b03143.

General experimental methods, characterization data, and NMR spectra (PDF)

## AUTHOR INFORMATION

## **Corresponding Author**

\*E-mail: markus.heinrich@fau.de. Tel: +49-9131-85-24115. Fax: +49-9131-85-22585.

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We thank Clara Schäfer (Pharmaceutical Chemistry, FAU Erlangen-Nürnberg) for experimental assistance.

#### REFERENCES

(1) Meerwein, H.; Büchner, E.; van Emster, K. J. Prakt. Chem. 1939, 152, 237.

(2) For review articles, see: (a) Heinrich, M. R. Chem. - Eur. J. 2009, 15, 820. (b) Vaillard, S. E.; Schulte, B.; Studer, A. In Modern Arylation Methods; Ackermann, L., Ed.; Wiley–VCH, 2009. (c) Pratsch, G.; Heinrich, M. R. In Radicals in Synthesis III. In Topics in Current Chemistry; Gansäuer, A., Heinrich, M. R., Eds.; Springer, 2012; Vol. 320. (d) Fehler, S. K.; Heinrich, M. R. Synlett 2015, 26, 580.

(3) For a recent extension to carbofluorination, see: (a) Kindt, S.; Heinrich, M. R. Chem. - Eur. J. **2014**, 20, 15344. (b) Guo, R.; Yang, H.; Tang, P. Chem. Commun. **2015**, *51*, 8829.

(4) (a) Grishchuk, B. D.; Baranovskii, V. S.; Simchak, R. V.; Tulaidan, G. N.; Gorbovoi, P. M. Russ. J. Gen. Chem. 2006, 76, 936. (b) Gorbovoi, P. M.; Tulaidan, G. N.; Grishchuk, B. D. Russ. J. Gen. Chem. 2008, 78, 133.

(5) (a) Brunner, H.; Blüchel, C.; Doyle, M. P. J. Organomet. Chem. 1997, 541, 89. (b) Mastrorilli, P.; Nobile, C. F.; Taccardi, N. Tetrahedron Lett. 2006, 47, 4759. (c) Obushak, N. D.; Lesyuk, A. I.; Gorak, Y. I.; Matiichuk, V. S. Russ. J. Org. Chem. 2009, 45, 1375.

(6) For reviews on photocatalyzed reactions, see: (a) Hari, D. P.; König, B. Angew. Chem., Int. Ed. **2013**, 52, 4734. (b) Mo, F.; Dong, G.; Zhang, Y.; Wang, J. Org. Biomol. Chem. **2013**, 11, 1582.

(7) For recent examples of photocatalyzed Meerwein arylations, see:
(a) Hari, D. P.; Hering, T.; König, B. Org. Lett. 2012, 14, 5334.
(b) Schroll, P.; Hari, D.; König, B. ChemistryOpen 2012, 1, 130. (c) Hari, D. P.; Schroll, P.; König, B. J. Am. Chem. Soc. 2012, 134, 2958. (d) Yao, C.-J.; Sun, Q.; Rastogi, N.; König, B. ACS Catal. 2015, 5, 2935. (e) Chen, Y.-H.; Lee, M.; Lin, Y.-Z.; Leow, D. Chem. - Asian J. 2015, 10, 1618.

(8) For Meerwein-type carbooxygenation reactions, see: (a) Heinrich, M. R.; Wetzel, A.; Kirschstein, M. Org. Lett. **2007**, *9*, 3833. (b) Dickschat,

A.; Studer, A. Org. Lett. **2010**, *12*, 3972. (c) Taniguchi, T.; Zaimoku, H.; Ishibashi, H. Chem. - Eur. J. **2011**, *17*, 4307. (d) Kindt, S.; Jasch, H.; Heinrich, M. R. Chem. - Eur. J. **2014**, *20*, 6251. (e) See also refs 7e and 9.

(9) Hartmann, M.; Li, Y.; Studer, A. J. Am. Chem. Soc. 2012, 134, 16516.

(10) Prasad Hari, D.; Hering, T.; König, B. *Angew. Chem., Int. Ed.* **2014**, 53, 725.

(11) Galli, C. Chem. Rev. 1988, 88, 765.

(12) (a) Canning, P. S. J.; McCrudden, K.; Maskill, H.; Sexton, B. J. J. Chem. Soc., Perkin Trans. 2 1999, 2, 2735. (b) Pazo-Llorente, R.; Bravo-Diaz, C.; Gonzalez-Romero, E. Eur. J. Org. Chem. 2004, 3221.

(13) For original work and mechanism of the Gomberg–Bachmann reaction, see: (a) Gomberg, M.; Bachmann, W. E. J. Am. Chem. Soc. **1924**, 46, 2339. (b) Rüchardt, C.; Merz, E. Tetrahedron Lett. **1964**, 5, 2431. (c) Bolton, R.; Williams, G. H. Chem. Soc. Rev. **1986**, 15, 261.

(14) For reviews of radical aryl-aryl coupling, see: (a) Dermer, O. C.; Edmison, M. T. *Chem. Rev.* **1957**, *57*, 77. (b) Studer, A.; Bossart, M. In *Radicals in Organic Synthesis*, 1st ed.; Renaud, P., Sibi, M., Eds.; Wiley-VCH, 2001; Vol. 2, p 62. (c) Bowman, W. R.; Storey, J. M. D. *Chem. Soc. Rev.* **2007**, *36*, 1803. (d) Studer, A.; Curran, D. *Angew. Chem., Int. Ed.* **2011**, *50*, 5018. (e) See also ref 2b-d.

(15) Recent developments in base-mediated radical aryl-aryl coupling: (a) Pratsch, G.; Wallaschkowski, T.; Heinrich, M. R. Chem. -Eur. J. 2012, 18, 11555. (b) Wu, Y.; Wong, S. M.; Mao, F.; Chan, T. L.; Kwong, F. Y. Org. Lett. 2012, 14, 5306. (c) De, S.; Ghosh, S.; Bhunia, S.; Sheikh, J. A.; Bisai, A. Org. Lett. 2012, 14, 4466. (d) Chen, W. C.; Hsu, Y. C.; Shih, W. C.; Lee, C. Y.; Chuang, W. H.; Tsai, Y. F.; Chen, P. P.; Ong, T. G. Chem. Commun. 2012, 48, 6702. (e) Cheng, Y.; Gu, X.; Li, P. Org. Lett. 2013, 15, 2664. (f) Dewanji, A.; Murarka, S.; Curran, D.; Studer, A. Org. Lett. 2013, 15, 6102. (g) Zhao, H.; Shen, J.; Guo, J.; Ye, R.; Zeng, H. Chem. Commun. 2013, 49, 2323. (h) Budén, M. E.; Guastavino, J. F.; Rossi, R. A. Org. Lett. 2013, 15, 1174. (i) Hofmann, J.; Jasch, H.; Heinrich, M. R. J. Org. Chem. 2014, 79, 2314. (j) Ghosh, D.; Lee, J.-Y.; Liu, C. Y.; Chiang, Y.-H.; Lee, H. M. Adv. Synth. Catal. 2014, 356, 406. (16) (a) Ek, F.; Axelsson, O.; Wistrand, L.-G.; Frejd, T. J. Org. Chem. 2002, 67, 6376. (b) Molinaro, C.; Mowat, J.; Gosselin, F.; O'Shea, P. D.; Marcoux, J. F.; Angelaud, R.; Davies, I. W. J. Org. Chem. 2007, 72, 1856. (17) For a related addition to isonitriles, see: Xia, Z.; Zhu, Q. Org. Lett. 2013, 15, 4110.

(18) (a) Zeng, W.; Chemler, S. R. J. Am. Chem. Soc. 2007, 129, 12948.
(b) Zavesky, B. P.; Babij, N. R.; Fritz, J. A.; Wolfe, J. P. Org. Lett. 2013, 15, 5420. (c) Choi, G. J.; Knowles, R. R. J. Am. Chem. Soc. 2015, 137, 9226. (d) Faulkner, A.; Scott, J. S.; Bower, J. F. J. Am. Chem. Soc. 2015, 137, 7224.

(19) For early examples, see: (a) Al Adel, I.; Salami, B. A.; Levisalles, J.; Rudler, H. *Bull. Soc. Chim. Fr.* **1976**, 934. (b) Citterio, A.; Minisci, F.; Albinati, A.; Bruckner, S. *Tetrahedron Lett.* **1980**, *21*, 2909. (c) Citterio, A.; Minisci, F.; Vismara, E. J. Org. Chem. **1982**, *47*, 81.

(20) For reviews on nitrogen-centered radical scavengers, see:
(a) Höfling, S. B.; Heinrich, M. R. Synthesis 2011, 173. (b) Ollivier,
C.; Renaud, P. In Radicals in Organic Synthesis, 1st ed.; Renaud, P., Sibi,
M., Eds.; Wiley–VCH, 2001; Vol. 2, p 93.

(21) (a) Heinrich, M. R.; Blank, O.; Wölfel, S. Org. Lett. 2006, 8, 3323.

(b) Heinrich, M. R.; Blank, O.; Wetzel, A. J. Org. Chem. 2007, 72, 476.
(c) Blank, O.; Wetzel, A.; Ullrich, D.; Heinrich, M. R. Eur. J. Org. Chem. 2008, 3179. (d) Prechter, A.; Gröger, H.; Heinrich, M. R. Org. Biomol.

Chem. 2012, 10, 3384. (22) For related radical carbonitrosation reactions, see: (a) de Salas, C.; Blank, O.; Heinrich, M. R. Chem. - Eur. J. 2011, 17, 9306. (b) de Salas, C.; Heinrich, M. R. Green Chem. 2014, 16, 2982.

(23) Yields determined by <sup>1</sup>H NMR using dimethyl terephthalate as internal standard. For inert behavior of acetonitrile, see: Kosynkin, D.; Bockman, T. M.; Kochi, J. K. *J. Am. Chem. Soc.* **1997**, *119*, 4846.

(24) For the isomerization of azo compounds to hydrazones, see: Landge, S. M.; Tkatchouk, E.; Benítez, D.; Lanfranchi, D. A.; Elhabiri, M.; Goddard, W. A.; Aprahamian, I. J. Am. Chem. Soc. 2011, 133, 9812.
(25) De Vleeschouwer, F.; Van Speybroeck, V.; Waroquier, M.; Geerlings, P.; De Proft, F. Org. Lett. 2007, 9, 2721. (26) For intramolecular radical attack on nitro groups, see: Heinrich, M. R.; Blank, O.; Ullrich, D.; Kirschstein, M. J. Org. Chem. 2007, 72, 9609.

(27) An iron(II)-mediated reaction of **1u** and **2a** gave azo compound **5** (7%) and also no yield of **3ua** (see ref 21c and the Supporting Information).

(28) Phatangare, K. R.; Borse, B. N.; Padalkar, V. S.; Patil, V. S.; Gupta, V. D.; Umape, P. G.; Sekar, N. J. Chem. Sci. **2013**, *125*, 141.

(29) Papanastassiou, Z. B.; McMillan, A.; Czebotar, V. J.; Bardos, T. J. J. Am. Chem. Soc. **1959**, 81, 6056.

(30) For characterization and stability of aryl diazotates, see: (a) Elofson, R. M.; Cyr, N.; Laidler, J. K. *Tetrahedron Lett.* **1990**, *31*, 7205. (b) Kauffmann, T.; Friestad, H. O.; Henkeler, H. Justus Liebigs Ann. Chem. **1960**, *634*, 64.

(31) Substituent effects on radical stabilization: Hioe, J.; Zipse, H. In *Encyclopedia of Radicals and Chemistry, Biology and Materials*, 1st ed.; Chatgilialoglu, C., Studer, A., Eds.; Wiley–VCH, 2012; Vol. 1, p 449.

(32) For related C-H activation reactions, see: (a) Robertson, J.; Pillai, J.; Lush, R. K. Chem. Soc. Rev. 2001, 30, 94. (b) Chiba, S.; Chen, H. Org. Biomol. Chem. 2014, 12, 4051. (c) Amaoka, Y.; Nagatomo, M.; Watanabe, M.; Tao, K.; Kamijo, S.; Inoue, M. Chem. Sci. 2014, 5, 4339. (d) Guptill, D. M.; Davies, H. M. L. J. Am. Chem. Soc. 2014, 136, 17718. (e) Soldi, C.; Lamb, K. N.; Squitieri, R. A.; González-López, M.; Di Maso, M. J.; Shaw, J. T. J. Am. Chem. Soc. 2014, 136, 15142. (f) Shi, H.; Babinski, D. J.; Ritter, T. J. Am. Chem. Soc. 2015, 137, 3775.

(33) For related Hofmann–Löffler–Freytag reactions, see: Paz, N. R.; Rodríguez-Sosa, D.; Valdés, H.; Marticorena, R.; Melián, D.; Copano, M. B.; González, C. C.; Herrera, A. J. *Org. Lett.* **2015**, *17*, 2370.

(34) For radical oxidations of alcohols and ethers via H-abstraction, see: Feray, L.; Kuznetsov, N.; Renaud, P. In *Radicals in Organic Synthesis*, 1st ed.; Renaud, P., Sibi, M., Eds.; Wiley–VCH, 2001; Vol. 2, p 246.

(35) Swain, C. G.; Powell, A. L.; Sheppard, W. A.; Morgan, C. R. J. Am. Chem. Soc. 1979, 101, 3576.

(36) Application of pyrazolidinones: White, J. D.; Toske, S. G. Bioorg. Med. Chem. Lett. **1993**, 3, 2383.