



Cite this: DOI: 10.1039/c5cc08181e

Received 2nd October 2015,
Accepted 16th November 2015

DOI: 10.1039/c5cc08181e

www.rsc.org/chemcomm

Even the normal is abnormal: N-heterocyclic carbene C² binding to a phosphalkene without breaking the P=C π -bond†

 Paresh Kumar Majhi,^{ab} Keith C. F. Chow,^b Tom H. H. Hsieh,^b Eric G. Bowes,^b
 Gregor Schnakenburg,^a Pierre Kennepohl,^{*b} Rainer Streubel^{*a} and Derek P. Gates^{*b}

The reaction of MesP=CPh₂ with the least sterically demanding N-heterocyclic carbene (NHC = IMe) results in formation of the 'abnormal' (C⁴-substituted) 4-phosphino-NHC (1). In contrast, reaction with Me₂Ime gives the unprecedented 'normal' C² adduct, Me₂Ime → P(Mes)=CPh₂ (2). Particularly striking is the asymmetric and weak bonding of the NHC to the P=C moiety in 2. DFT calculations indicate that the P=C natural bond order in 2 (1.54) still reflects significant π -character to the bond (cf. MesP=CPh₂: NBO = 1.98). Further computational analysis suggests that π -delocalization into the remote C-phenyl substituents is key to stabilizing the NHC adduct.

In just a few decades, N-heterocyclic carbenes (NHCs), the most important of which are the imidazol-2-ylidenes,¹ have risen from relative obscurity to becoming competitive with phosphines as important ligands in organometallic chemistry and catalysis.² The carbene carbon at the C² position in such an NHC (Fig. 1) is by far the most reactive site (so-called "classical" or "normal" binding).³

A growing number of NHCs have been observed to react through the C⁴ position (so-called "mesoionic" or "abnormal" binding).^{4–8} To obtain abnormal N-heterocyclic carbenes (aNHCs), the C²-position is usually blocked and the C⁴-position bound to a d- or f-block element. For example, the isolation of the first free aNHC by Bertrand and co-workers required phenyl moieties at both the C² and C⁵ positions.⁵ In 2009, we reported the "abnormal" reaction of the unblocked 1,3-dimesityl-imidazol-2-ylidene (IMes) with the phosphalkene MesP=CPh₂ to afford the first example of an NHC with a phosphine substituent at C⁴.^{9,10} The direct functionalization of unprotected NHCs with p-block elements at the C⁴ position has grown to include: halogenation,^{11,12} deuteration,¹³ and silylation,¹⁴ addition of Si=N¹⁵ and B=N¹⁶ compounds,




Fig. 1 Structures and numbering scheme for normal (NHC) and abnormal imidazol-2-ylidenes (aNHC).

s-block metallation,^{17,18} and borate formation.¹⁹ An elegant method to induce aNHC reactivity involves temporarily blocking C² with an electrophile to permit base-assisted functionalization selectively at C⁴, thereby affording NHCs with C⁴ substituents such as: Br, Cl, C(O)R, OTf, PPh₂, SiMe₃.^{20–22}

It was proposed that the sterically demanding N-Mes substituents of IMes might impede the formation of the "normal" (C²) product in the reaction with phosphalkenes thus leading to the "abnormal" (C⁴) 4-phosphino-NHC.²³ To evaluate this hypothesis, we sought to probe the reactivity of less sterically-demanding NHCs with the phosphalkene MesP=CPh₂.


Significantly, we have now isolated the "abnormal" product 1 and the "normal" adduct 2, both of which have been crystallographically characterized (1 as its BH₃ complex, 1·BH₃). Adduct 2 is particularly remarkable in that the metrical data along with DFT calculations reveal that the NHC binds to the phosphalkene π^* with a low degree of disruption of the P=C π -bond.

1,3-Dimethyl-imidazol-2-ylidene (IMe) was generated *in situ* in THF²⁴ and treated with MesP=CPh₂ (1 equiv.) at –78 °C followed by warming to room temperature (Scheme 1). Analysis of an aliquot removed from the reaction mixture by ³¹P{¹H} NMR spectroscopy revealed that the signal for MesP=CPh₂ ($\delta^{31}\text{P}$ = 234 ppm) was not present and a new signal at –37.2 ppm had appeared, assigned to 1. Importantly, the chemical shift of this new signal is comparable to the abnormal product derived from IMes and MesP=CPh₂ (δ = –37.3 ppm).⁹ The isolated product (yield 78%) was analyzed by ¹³C{¹H} NMR spectroscopy in THF-*d*⁸ to reveal a broad resonance at δ = 222.5 ppm, assigned to the carbene carbon (C²) atom of 1; the ³J_{P,C} coupling constant could not be resolved.

^a Institut für Anorganische Chemie, Rheinische Friedrich-Wilhelms-Universität Bonn, Gerhard-Domagk-Straße 1, D-53121 Bonn, Germany. E-mail: r.streubel@uni-bonn.de

^b Department of Chemistry, University of British Columbia, 2036 Main Mall, Vancouver, British Columbia, V6T 1Z1, Canada. E-mail: dgates@chem.ubc.ca

† Electronic supplementary information (ESI) available: Experimental procedures, characterization data, X-ray data, DFT procedures and results, selected spectra. CCDC 1428162 (1·BH₃) and 1428161 (2). For crystallographic data in CIF or other electronic format see DOI: 10.1039/c5cc08181e

Scheme 1 Synthesis of compounds **1** and **2**.

In order to obtain X-ray crystallographic confirmation of **1**, several complexes **1**·ML_n [ML_n = BH₃, Rh(cod)Cl, Cr(CO)₅, Mo(CO)₅, W(CO)₅] were prepared. The C² bound complexes were characterized by ³¹P, ¹H, ¹³C{¹H} NMR spectroscopy²⁵ and mass spectrometry (see ESI†). In the case of **1**·BH₃, single crystals suitable for X-ray crystallography were obtained from a saturated THF solution at −20 °C. The molecular structure (Fig. 2) confirms the formulation as the 4-phosphinyl-substituted NHC borane complex **1**·BH₃. The metrical parameters in **1**·BH₃ do not differ significantly from those of IMe·BH₃²⁶ and, hence, shall not be discussed further.

Despite monitoring the abnormal reaction of IMe with MesP=CPh₂ by ³¹P NMR spectroscopy at variable temperatures (−80 to 25 °C), no P-containing intermediates have been observed. In contrast, Braunschweig and co-workers have detected a “normal” C²-adduct as a short-lived intermediate in the abnormal reaction of IPr with the iminoborane, *t*BuB=N*t*Bu.¹⁶ In an effort to see whether the formation of the normal C² adduct was possible with phosphoralkenes, the C⁴/C⁵-blocked 1,3,4,5-tetramethyl-imidazol-2-ylidene

(Me₂IMe)²⁷ was treated with MesP=CPh₂ in Et₂O. ³¹P{¹H} NMR spectroscopic analysis of the reaction mixture revealed that the signal for MesP=CPh₂ (δ = 234) had been replaced by a broad signal at δ = 206.1 (w_{1/2} = 124 Hz) along with about 7% of unidentified products. Interestingly, the chemical shift is highly dependent on the solvent in which the reaction is conducted (THF: δ = 185.1, w_{1/2} = 608 Hz; toluene: δ = 213.4, w_{1/2} = 122 Hz). Given the similarity in chemical shift to that of MesP=CPh₂, we suspected that the product, regardless of solvent, retains significant P=C bond character. In addition, the phosphoralkene is released from **2** upon the addition of boranes [BR₃ = BPh₃, B(C₆F₅)₃] to yield a mixture of MesP=CPh₂ and Me₂IMe·BR₃.

X-ray crystallography permitted the product described above to be identified as the unprecedented NHC-phosphaalkene adduct **2** (Fig. 3). This “normal” NHC adduct **2** features a short P–C bond [P(1)–C(17) = 1.7420(19) Å] that is closer to that of a double bond (*e.g.* for MesP=CPh₂ [1.692(3) Å]²⁸) than that of a typical P–C single bond (1.85 Å).²⁹ In addition, the carbon atom C(17) retains a planar geometry Σ∠C(17) = 359.7°, which is consistent with retention of its sp² hybridization. The distortion of the P=C bond upon binding of NHC is small with a 36.4° twisting of the C(1)–P(1)–C(10) plane with respect to the C(18)–C(17)–C(24) plane; for comparison, the same angle in MesP=CPh₂ is 0.9°³⁰ or 7.5°.³¹ Particularly striking is the almost perpendicular position of the NHC donor with respect to the best plane of the P=C bond (102.6° is max. angle between NHC–P bond and P=C best plane). In addition, the C(10)–P(1) single bond length [P(1)–C(10) = 1.8512(18) Å] is long. These observations stand in stark contrast to the Me₂IPr adduct of Mes*N=P–Cl³² which displays significant torsion of the C_{Mes}–N=P–Cl unit and in the Me₂IPr adduct of *t*BuN=B*t*Bu¹⁶ where C_{*t*Bu}–N=B–C_{*t*Bu} and C² are coplanar. The molecular structures of related NHC-complexes to E=E bonds also exhibit significant distortions at the E=E bond.³³ The above




Fig. 2 Molecular structure of complex **1**·BH₃. Disordered solvent molecule (THF) and hydrogen atoms (except B–H and C(6)–H) have been omitted for clarity (50% probability level). Selected bond distances (Å) and angles (°): C(2)–B 1.609(4), C(2)–N(1) 1.343(3), C(2)–N(2) 1.357(3), C(1)–C(3) 1.352(3), C(1)–P 1.817(3), C(6)–P 1.881(2), C(1)–N(1) 1.407(3), C(3)–N(2) 1.374(3); N(1)–C(2)–N(2) 104.7(2), N(1)–C(2)–B 128.9(2), N(2)–C(2)–B 126.4(3), N(1)–C(1)–P 121.21(18).




Fig. 3 Molecular structure of complex **2**. Hydrogen atoms have been omitted for clarity (50% probability level); side view of the reduced structure (top left). Selected bond distances (Å) and angles (°): P(1)–C(10) 1.8512(18), P(1)–C(17) 1.7420(19), P(1)–C(1) 1.8416(17), N(1)–C(10) 1.354(2), N(2)–C(10) 1.345(2), C(17)–C(18) 1.479(2); N(1)–C(10)–N(2) 106.14(15), N(1)–C(10)–P(1) 121.94(14), N(2)–C(10)–P(1) 128.95(13), C(10)–P(1)–C(1) 100.98(7), C(10)–P(1)–C(17) 112.73(9), C(18)–C(17)–C(24) 118.57(16).

metrical data for **2** combined with the ^{31}P chemical shift and the ready displacement of the NHC with electrophiles are all consistent with the interpretation of the NHC as a weak σ -donor in **2**.

The fascinating metrical parameters in **2** led us to probe the nature of bonding in this compound and in $\text{MesP}=\text{CPh}_2$ by computational methods (DFT-B3LYP/TZVP). The optimized gas phase geometry was in good agreement with the solid state molecular structure (see ESI†) and natural bond orbital (NBO) analysis suggests that both P(1) and C(17) retain significant π -bond character. On first glance, the data is consistent with adduct formation between the C^2 lone pair of Me_2IME and the $\text{P}=\text{C}$ π^* orbital of $\text{MesP}=\text{CPh}_2$. Surprisingly, this adduct formation only leads to a small increase in the π^* population in **2** (by 0.08 e^- compared to $\text{MesP}=\text{CPh}_2$) and displays a concomitant decrease in the NBO $\text{P}=\text{C}$ bond order (from 1.98 to 1.54). The $\text{P}=\text{C}$ π bond polarizes substantially upon binding of Me_2IME (Fig. 4); the phosphorus natural hybrid orbital (NHO) shows increased 3d character in the π_{PC} bond leading to pd hybridization, and this pd hybrid functions as the acceptor orbital for the $\text{Me}_2\text{IME} \rightarrow \text{P}$ σ bond. Nevertheless, the actual contributions from the putative pd hybrid NHOs in both the $\pi_{\text{P(1)-C(17)}}$ and the $\sigma_{\text{P(1)-C(10)}}$ orbitals are quite small (11% and 23%, respectively).

The calculated atomic charges obtained by natural population analysis (NPA) are shown in Table 1 and the changes (Δq) reveal the overall effect of adduct formation. Predictably, there is an electron redistribution towards the carbon of the phosphalkene moiety that is accompanied by a concomitant decrease in charge at phosphorus. The significant redistribution of charge density from the NHC to the remote phenyl substituents was unanticipated, particularly when noting the asymmetry in negative charge accumulation between the two substituents (see Δq for Ph_{cis} vs. Ph_{trans} in Table 1). However, these observations are in agreement with the different angles between the $\text{P}=\text{C}$ bond plane and each phenyl substituent in **2** (*cis*, 17.5° ; *trans*, 51.9°); the metrical data contrasts the analogous angles within the free $\text{MesP}=\text{CPh}_2$ ³⁴ ($\text{Ph}_{\text{cis}} > \text{Ph}_{\text{trans}}$). This finding is consistent with asymmetric allylic π -delocalization across the CPh_2 moiety. In fact, a comparison to the hypothetical Me_2IME adduct of $\text{MesP}=\text{CMe}_2$ shows that the phenyl substituents are critical to stabilizing **2**. Namely, the dimethyl derivative exhibits a much longer $\text{P}-\text{C}_{\text{NHC}}$ bond (2.00 \AA vs. 1.86 \AA) and this bond has a significantly lower bond dissociation energy ($\sim 45\text{ kJ mol}^{-1}$ lower than that for **2**).




Fig. 4 NBO representation of the NBO-derived π_{PC} orbital in $\text{MesP}=\text{CPh}_2$ (left) and **2** (right). Orbital surfaces are plotted at $0.05\text{ e}^- \text{ \AA}^{-3}$ using WebMO.

Table 1 Calculated NPA atomic and fragment charges for $\text{MesP}=\text{CPh}_2$, Me_2IME , and the resultant adduct **2**. Δq is the change in charge between **2** and the sum of the charges in the reactants

	NPA atomic/fragment charges					
	Mes	Me_2IME	P	CPh_2	Ph_{cis}	Ph_{trans}
$\text{MesP}=\text{CPh}_2$	−0.30	—	0.66	−0.38	0.02	0.01
IME_4	—	0.00	—	—	—	—
2	−0.32	0.35	0.90	−0.65	−0.16	−0.13
Δq	−0.01	0.35	0.24	−0.26	−0.18	−0.14

Analysis by natural resonance theory (NRT) suggests that the major contributors to the canonical forms of **2** ($\text{R} = \text{Ph}$) are the simple adduct (**2A**) and the asymmetric allylic form (**2B**) (Fig. 5). It is interesting that the structure **2B** showing π -delocalization to Ph_{cis} is considerably more important than that involving Ph_{trans} (**2D**). Importantly, this result is consistent with the crystal structure data described above. Moreover, this analysis confirms that significant $\text{P}=\text{C}$ character remains even though charge is highly delocalized. Overall, the results are consistent with an interpretation of the bonding within **2** where the $\pi_{\text{P}=\text{C}}$ bond is perturbed by the NHC donor but is not broken entirely resulting in the unique geometry of the isolated adduct.

Herein, we have shown that steric bulk at the *N*-substituents of NHCs is not required for abnormal products to be formed in their reactions with phosphalkenes. By contrast, we find that blocking the $\text{C}^{4,5}$ positions of the NHC leads to the formation of an unprecedented weakly bound C^2 -adduct that retains significant $\text{P}=\text{C}$ double bond character. The formation of such adducts is predicated on the ability for charge delocalization in the substituents of the phosphalkene and suggests that modulation of the stability of the NHC adduct is strongly dependent on the nature of the phosphalkene.

DG and PK are grateful to the Natural Sciences and Engineering Research Council (NSERC) of Canada for financial support of this work. RS thanks the Deutsche Forschungsgemeinschaft (DFG) SFB 813 “Chemistry at Spin Centers” and the COST action (CM1302). RS and PKM are grateful to the DAAD PPP Program for Canada. Computation work was enabled by support by WestGrid (www.westgrid.ca) and Compute Canada Calcul Canada (www.computeCanada.ca). We are grateful to Mr. S. Serin




Fig. 5 Generalized canonical forms illustrating the bonding within $\text{MesP}=\text{CR}_2$ (**A**, $\text{R} = \text{Ph}$ or Me) and a $\text{NHC} \rightarrow \text{MesP}=\text{CR}_2$ (**2A–D**, $\text{R} = \text{Ph}$ or Me). The percent contributions are obtained from an NRT analysis of the NBO computational results (see ESI† for details).

for assistance in the preparation of Fig. 1 and 2 and determination of some metrical parameters.

Notes and references

- 1 A. J. Arduengo III, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1991, **113**, 361.
- 2 M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485.
- 3 D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39.
- 4 S. Grundemann, A. Kovacevic, M. Albrecht, J. W. Faller Robert and H. Crabtree, *Chem. Commun.*, 2001, 2274.
- 5 E. Aldeco-Perez, A. J. Rosenthal, B. Donnadieu, P. Parameswaran, G. Frenking and G. Bertrand, *Science*, 2009, **326**, 556.
- 6 O. Schuster, L. Yang, H. G. Raubenheimer and M. Albrecht, *Chem. Rev.*, 2009, **109**, 3445.
- 7 P. L. Arnold and S. Pearson, *Coord. Chem. Rev.*, 2007, **251**, 596.
- 8 J. B. Waters and J. M. Goicoechea, *Coord. Chem. Rev.*, 2015, **293**, 80.
- 9 J. I. Bates, P. Kennepohl and D. P. Gates, *Angew. Chem., Int. Ed.*, 2009, **48**, 9844.
- 10 NHCs bearing an anionic PR-substituent at C⁴ have subsequently been reported: P. K. Majhi, G. Schnakenburg, Z. Kelemen, L. Nyulaszi, D. P. Gates and R. Streubel, *Angew. Chem., Int. Ed.*, 2013, **52**, 10080.
- 11 A. J. Arduengo III, F. Davidson, H. V. R. Dias, J. R. Goerlich, D. Khasnis, W. J. Marshall and T. K. Prakasha, *J. Am. Chem. Soc.*, 1997, **119**, 12742.
- 12 M. L. Cole, C. Jones and P. C. Junk, *New J. Chem.*, 2002, **26**, 1296.
- 13 M. K. Denk and J. M. Rodezno, *J. Organomet. Chem.*, 2001, **617–618**, 737.
- 14 H. Cui, Y. Shao, X. Li, L. Kong and C. Cui, *Organometallics*, 2009, **28**, 5191.
- 15 R. S. Ghadwal, H. W. Roesky, M. Granitzka and D. Stalke, *J. Am. Chem. Soc.*, 2010, **132**, 10018.
- 16 H. Braunschweig, W. C. Ewing, K. Geetharani and M. Schafer, *Angew. Chem., Int. Ed.*, 2015, **54**, 1662.
- 17 Y. Wang, Y. Xie, M. Y. Abraham, P. Wei, H. F. Schaefer, P. v. R. Schleyer and G. H. Robinson, *J. Am. Chem. Soc.*, 2010, **132**, 14370.
- 18 D. R. Armstrong, S. E. Baillie, V. L. Blair, N. G. Chabloz, J. Diez, J. Garcia-Alvarez, A. R. Kennedy, S. D. Robertson and E. Hevia, *Chem. Sci.*, 2013, **4**, 4259.
- 19 S. Kronig, E. Theuergarten, C. G. Daniliuc, P. G. Jones and M. Tamm, *Angew. Chem., Int. Ed.*, 2012, **51**, 3240.
- 20 D. Mendoza-Espinosa, B. Donnadieu and G. Bertrand, *J. Am. Chem. Soc.*, 2010, **132**, 7264.
- 21 J. Ruiz and A. F. Mesa, *Chem. – Eur. J.*, 2012, **18**, 4485.
- 22 K. Schwedtmann, M. H. Holthausen, K.-O. Feldmann and J. J. Weigand, *Angew. Chem., Int. Ed.*, 2013, **52**, 14204.
- 23 J. I. Bates and D. P. Gates, *Organometallics*, 2012, **31**, 4529.
- 24 A. Wacker, H. Pritzkow and W. Siebert, *Eur. J. Inorg. Chem.*, 1998, 843.
- 25 Consistent with that observed for the Rh(i) and Ir(i) complexes of 4-phosphinyl NHC derived from IMes (see ref. 23), two signals were observed in the ³¹P NMR spectra of complexes 1-ML_n [ML_n = BH₃, Rh(cod)Cl, Cr(CO)₅, Mo(CO)₅, W(CO)₅]. Although the origin of these signals is not known, we tentatively attribute these signals to isomers.
- 26 P. Bissinger, H. Braunschweig, T. Kupfer and K. Radacki, *Organometallics*, 2010, **29**, 3987.
- 27 N. Kuhn and T. Kratz, *Synthesis*, 1993, 561.
- 28 T. A. van der Knaap, T. C. Klebach, F. Visser, F. Bickelhaupt, P. Ros, E. J. Baerends, C. H. Stam and M. Konijn, *Tetrahedron*, 1984, **40**, 765.
- 29 F. H. Allen, O. Kennard, D. G. Watson, L. Brammer, A. G. Orpen and R. Taylor, *J. Chem. Soc., Perkin Trans. 2*, 1987, S1.
- 30 T. A. Vanderknaap, T. C. Klebach, F. Visser, F. Bickelhaupt, P. Ros, E. J. Baerends, C. H. Stam and M. Konijn, *Tetrahedron*, 1984, **40**, 765.
- 31 O. Mundt, G. Becker, W. Uhl, W. Massa and M. Birkhahn, *Z. Anorg. Allg. Chem.*, 1986, **541**, 319.
- 32 N. Burford, T. S. Cameron, D. J. LeBlanc, A. D. Phillips, T. E. Concolino, K. C. Lam and A. L. Rheingold, *J. Am. Chem. Soc.*, 2000, **122**, 5413.
- 33 For recent examples, see: (a) D. Geiss, M. I. Arz, M. Strassmann, G. Schnakenburg and A. C. Filippou, *Angew. Chem., Int. Ed.*, 2015, **54**, 2739; (b) S. Roy, P. Stollberg, R. Herbst-Irmer, D. Stalke, D. M. Andrada, G. Frenking and H. W. Roesky, *J. Am. Chem. Soc.*, 2015, **137**, 150; (c) F. Dahcheh, D. W. Stephan and G. Bertrand, *Chem. – Eur. J.*, 2015, **21**, 199; (d) F. Dahcheh, D. Martin, D. W. Stephan and G. Bertrand, *Angew. Chem., Int. Ed.*, 2014, **53**, 13159; (e) A. Jana, V. Huch and D. Scheschkewitz, *Angew. Chem., Int. Ed.*, 2013, **52**, 12179.
- 34 M. Yam, J. H. Chong, C.-W. Tsang, B. O. Patrick, A. E. Lam and D. P. Gates, *Inorg. Chem.*, 2006, **45**, 5225.