

Wagner–Meerwein-Type Rearrangements of Germapolysilanes - A Stable Ion Study

Lena Albers,[†] Mohammad Aghazadeh Meshgi,[§] Judith Baumgartner,^{*,‡} Christoph Marschner,^{*,§} and Thomas Müller^{*,†}


[†]Institut für Chemie, Carl von Ossietzky Universität Oldenburg, Carl von Ossietzky-Strasse 9-11, D-26129 Oldenburg, Federal Republic of Germany

[‡]Institut für Chemie, Karl Franzens Universität Graz, Stremayrgasse 9, 8010 Graz, Austria

[§]Institut für Anorganische Chemie, Technische Universität Graz, Stremayrgasse 9, 8010 Graz, Austria

Supporting Information

ABSTRACT: The rearrangement of tris(trimethylsilyl)silyl-trimethylgermane **1** to give tetrakis(trimethylsilyl)germane **2** was investigated as a typical example for Lewis acid catalyzed Wagner–Meerwein-type rearrangements of polysilanes and polygermasilanes. Direct ²⁹Si NMR spectroscopic evidence is provided for several cationic intermediates during the reaction. The identity of these species was verified by independent synthesis and NMR characterization, and their transformation was followed by NMR spectroscopy.



INTRODUCTION


Despite the most interesting properties and the widespread potential use of oligo- and polysilanes, synthetic access to this class of compounds is mainly limited to the Wurtz-type coupling reaction, in which the Si–Si chain is built up in a reductive process using silicon halides and alkali metals.¹ Up to date alternative methods for the synthesis of oligosilanes such as dehydrocoupling of hydrosilanes² or electrochemically mediated Si–Si bond forming processes³ do not reach the prominence of the Wurtz-type coupling. While this process works fairly well for the preparation of dialkylated linear polysilane polymers and small cyclosilanes, it does not provide satisfactory access to compounds of even slightly enhanced structural complexity. In particular, the generation of well-defined polycyclic molecules in acceptable yields has proven to be very challenging by this method.¹ Regarding the potential for such materials being useful for technical applications, there is intrinsic interest in novel approaches that open the possibility to develop new synthetic tools for the preparation of open chain, cyclic, and polyhedral compounds with silicon containing backbones. Pioneering work by the group of Kumada and Ishikawa,⁴ further developed by the groups of West,⁵ Märkl,^{6a} and Pannell,^{6b} revealed that the Lewis acid catalyzed Wagner–Meerwein-type rearrangement of oligosilanes is a powerful method to create structures of higher complexity. The high yield synthesis of an all-silicon analogue of adamantane from a structural isomer, reported previously by two of us, is certainly one of the more spectacular examples for the potential of this synthetic methodology.⁷

The description of this process as being a silicon variation of the Wagner–Meerwein rearrangement suggests the presence of cationic intermediates.^{4,5} However, despite the fact that the chemistry of silyl cations has experienced immense popularity in recent years,⁸ experimental evidence for polysilanyl-substituted silyl cations and their possible involvement in rearrangement or fragmentation reactions is indeed rather scarce. Lambert and co-workers reported on the formation of tris(trimethylsilyl)silylium, (Me₃Si)₃Si⁺, most probably in the form of its arene complexes; the obtained ²⁹Si NMR spectra suggest, however, substantial decomposition of the material.⁹ Structural aspects and the question of aromaticity and homoaromaticity had been the focus in several publications of the Sekiguchi group concerning some cyclic polysilasil cations.^{10,11} Sekiguchi and co-workers also demonstrated the occurrence of a degenerated intramolecular 1,3-methyl shift occurring in a dialkyloligosilanyl silyl cation.¹² Interestingly, in none of these reports were parallels drawn between the subject of study and the likely involvement of similar species in the sila-Wagner–Meerwein rearrangement. During our own experimental studies of the Wagner–Meerwein rearrangement in oligosilanes,¹³ we found that for germysilanes a particular preference exists for the formation of products in which the germanium atoms occupy central positions in the oligosilane frameworks.¹⁴ A deeper insight was provided by a computational study of the fundamental reaction of germysilane **1** to

Received: May 21, 2015


give germane **2** (reaction 1, Scheme 1). The results of density functional theory (DFT) calculations suggested an exothermic

Scheme 1. Rearrangement of Germylsilane 1 to Silylgermane 2 Using Lewis Acids



multistep rearrangement via several isomeric silyl- and germyl cations **3**–**8** (Scheme 2). According to this investigation, the isomerization occurs via consecutive 1,2-silyl and 1,2-methyl shifts with an overall barrier of less than 50 kJ mol^{−1}.¹⁴

Scheme 2. Suggested Isomerization Cascade of Germylum Ion 3 to Silylum Ion 8¹⁴




An experimental investigation on the course of the principal reaction 1 would greatly benefit from the simplicity of the reactants, proposed intermediates, and products and from the straightforward interpretation of their NMR spectra. The close chemical relationship between silicon and germanium suggests that conclusions drawn from a mechanistic study of reaction 1 will be of significance also in the case of polysilanes. Here, we report on a ²⁹Si NMR investigation of the exemplary reaction 1, which provides direct evidence for cationic intermediates in the sila-Wagner–Meerwein rearrangement and for the clean transformation of the incipient germyl cation **3** into the product-forming silyl cation **8** at low temperatures along a cascade of isomeric cations **4**–**7** (Scheme 2).

RESULTS AND DISCUSSION

As a starting point for our experimental investigation, we tested additional cationic Lewis acids as stoichiometric reagents or catalysts for the rearrangement of model compound **1**. The idea behind using a cationic Lewis acid is to introduce weakly coordinating anions such as the tetrakis(pentafluorophenyl)-borate, [B(C₆F₅)₄][−],¹⁵ which will be beneficial for the stabilization of the postulated cationic intermediates. Indeed, we found that already 4 mol % of trityl borate [Ph₃C][B(C₆F₅)₄], **9**, induces the complete transformation of germylsilane **1** into the symmetric germane **2** in dichloromethane at room temperature (Scheme 1).¹⁶ For the generation of stoichiometric amounts of the postulated cationic intermediates in the rearrangement reaction shown in Scheme 1, we switched to the even more electrophilic trialkylsilyl arenium borates,



[R₃Si(arene)] [B(C₆F₅)₄] (R = Me, Et, ⁱPr; arene = C₆H₅CH₃, C₆H₆, C₆H₅Cl), **10**,⁸ and used arenes as solvents. In the past, it was shown⁸ that the combination of arene solvents and weakly coordinating anions is suitable for the detection of highly reactive silylium ions. In order to allow the NMR spectroscopic detection of the postulated cationic intermediates, we tested at first the stoichiometric reaction between germylsilane **1** and triethylsilyl benzenium borate, [Et₃Si(C₆D₆)] [B(C₆F₅)₄], **10a**. The reaction in benzene at 6 °C was completed after 2 h and resulted in a biphasic reaction mixture that is typical for solutions of tetrakis(pentafluorophenyl)borates in aromatic hydrocarbons.⁸ ²⁹Si NMR spectra obtained from the ionic phase showed many signals, which indicated the formation of a complicated reaction mixture. The ²⁹Si NMR analysis of the second, nonpolar layer showed the presence of at least nine different neutral compounds. These nine compounds were identified by GC/MS spectrometry and NMR spectroscopy as the expected germane **2**, three different methylethylsilanes, and a series of silylated germanes **11** and **12** (Scheme 3 and

Scheme 3. Stoichiometric Reaction between Triethylsilyl Benzenium Borate 10a and Silylgermane 1 in Benzene at T = 6 °C



Supporting Information for details). Noteworthy, germanes **11** and **12**, which all have cation **8** as a common intermediate, are substituted with a different number of ethyl groups. When the same reaction was performed with tri-*iso*-propylsilyl toluenium borate, [Pr₃Si(C₇D₈)] [B(C₆F₅)₄], **10b**, only two side products were detected, namely, (Me₃Si)₃GeSiMe₂ⁱPr and Pr₃SiMe. Finally, when trimethylsilyl toluenium borate [Me₃Si(C₇D₈)] [B(C₆F₅)₄], **10c**, was applied as reaction partner, only the expected rearrangement product **2** and tetramethylsilane were identified in the nonpolar phase (see Supporting Information). Although with all three silylarenium borates **10** the clean production of cationic species at room temperature was not achieved, the results of these stoichiometric reactions indicate that the initial step of the rearrangement reaction is cation formation by cleavage of either a Ge–C- or Si–C bond. Judged from the calculated bond strengths, the cleavage of a Ge–C bond is preferred over that of a Si–C bond (D_e(Ge–C) = 305 kJ mol^{−1}; D_e(Si–C) = 347 kJ mol^{−1}).¹⁴ This clearly supports our initial assumption that the first cationic intermediate formed should be germyl cation **3** which undergoes a rearrangement reaction to give silyl cation **8**. Furthermore, the results suggest that the terminating step consists in back transfer of an alkyl group to the rearranged cation which yields besides germane **2** the alkylated products **13** (Scheme 4). The multiple substitution⁵ in the case of the ethyl substituted silyl arenium ion demonstrates the ease of alkyl group exchange under the applied conditions, which is only limited by the steric requirements of the exchanging groups.

In contrast to the reaction at room temperature, the reaction of silylgermane **1** with tri-*iso*-propylsilyl toluenium borate, **10b**, in toluene-*d*₈ at −20 °C yielded cleanly a single ionic compound as indicated by only two resonances that were detected in the ²⁹Si NMR spectra of the ionic phase. The NMR

Scheme 4. Reaction of Silylgermane **1** with Different Trialkylsilyl Arenium Borates **10** at r.t. (Borate Anion Not Shown)Scheme 5. Formation of Silyl Tolenium Ion **8**(C₇D₈) from Different Precursor CompoundsTable 1. Experimental and Calculated ²⁹Si NMR Chemical Shifts of Solvent Complexes of Gernylium Ions **3** and **7** and Silylium Ion **8**^a

compound	$\delta^{29}Si$ (Si ^q)	$\delta^{29}Si$ (SiMe ₃)	$\delta^{29}Si$ (Si ⁺ Me ₂)	$\delta^{29}Si$ (SiMe ₂)
[3 (C ₇ D ₈)]	-87.9 (-93)	-7.9 (-7)		
[7 (C ₇ D ₈)]		(8; -13)		(-13)
[8 (C ₇ D ₈)]		-2.4 (0)	98.1 (108)	
[8 (ClC ₆ D ₅)] ^b		-1.5 (0)	154.3 (157)	

^aCalculated values for the nondeuterated compounds at GIAO/M06-L/6-311G(2d,p)//M06-2X/6-311+G(d,p) are in parentheses. ^bCalculated for the chloronium ion structure [**8**-Cl-C₆H₅]. Chloronium ions are the dominating species when silyl cations are generated in chlorinated arenes; see ref 22.

data ($\delta^{29}Si = -87.9$ (Si^q) and $\delta^{29}Si = -7.9$ (SiMe₃), integral ratio in the ²⁹Si{¹H} inverse gated experiment, 1:3) is consistent with the structure of germyl tolenium ion **3**(C₇D₈). This assignment was supported by the fact that the same cationic species was formed by the conventional Bartlett Condon Schneider hydride transfer¹⁷ reaction of hydridogermene **14** with trityl borate **9** (Scheme 5). In addition, the results of quantum mechanical calculations^{18,19} of ²⁹Si NMR chemical shifts for an optimized molecular structure of **3**(C₇H₈) predict values that are very close to the experiment (see Table 1).

Interestingly, already after 2 h at -20 °C two new ²⁹Si NMR resonances were detected in both cases ($\delta^{29}Si = -2.4$ and +98.1, integral ratio in the ²⁹Si{¹H} inverse gated experiment, 3:1; see Figure 1 and Supporting Information).²⁰ Over time, the intensity of the new signals increased significantly while the original ²⁹Si NMR signals of germyl cation **3**(C₇D₈) vanished (see Figure 1), thereby confirming the clean conversion of silylgermyl tolenium ion **3**(C₇D₈) into a new cationic species in toluene at -20 °C. The new high field signal appears in the typical region for trimethylsilyl groups attached to germanium,²¹ but the measured chemical shift is not specific. More informative is the low field resonance at $\delta^{29}Si = 98.1$, which is

characteristic for silyl arenium ions.⁸ The ²⁹Si NMR chemical shifts of these cationic species are reported to be very sensitive to changes of the solvent due to the replacement of the arene molecule attached to the silicon atom. Indeed, the ²⁹Si NMR chemical shift of the low field resonance varies significantly with the arene solvent from $\delta^{29}Si = 98.1$ in toluene-*d*₈ to $\delta^{29}Si = 154.3$ in chlorobenzene-*d*₅ (see Table 1).²² On the basis of these results, the two new ²⁹Si NMR signals were tentatively assigned to silyl tolenium ion **8**(C₇D₈). This assignment was supported by its independent synthesis by hydride transfer reaction from the appropriate silane **15** (see Scheme 5 and Figure 2) and by quantum mechanical calculations of the ²⁹Si NMR chemical shifts of optimized molecular structures of arene complexes of **8** (see Table 1). According to the NMR results, solutions of the silyl tolenium borate [**8**(C₇D₈)] [B-(C₆F₅)₄] in toluene at *T* = -20 °C are stable for at least 3 days. When the temperature of the NMR probe was gradually raised in steps of 10 °C, already at *T* = -10 °C new signals in the ²⁹Si NMR spectrum appeared, and exposure of the NMR sample to temperatures of about +20 °C resulted in complete decomposition of cation **8**(C₇D₈) (see Supporting Information, Figure S21).²³ The thermal instability of arenium borate [**8**(C₇D₈)] [B(C₆F₅)₄] in solution prevents its isolation in




Figure 1. 99 MHz $^{29}\text{Si}\{^1\text{H}\}$ NMR spectra (toluene- d_8 , 253 K) of the rearrangement of germyl toluenium ion $3(\text{C}_7\text{D}_8)$ (●) to silyl toluenium $8(\text{C}_7\text{D}_8)$ (*) ($\downarrow$ $^i\text{Pr}_3\text{SiMe}$). (a) Spectrum 2 h after the addition of $[\text{Pr}_3\text{Si}(\text{C}_7\text{D}_8)][\text{B}(\text{C}_6\text{F}_5)_4]$; (b) after 9 h; (c) after 25 h; (d) after 120 d at 210 K.




Figure 2. 99 MHz $^{29}\text{Si}\{^1\text{H}\}$ NMR spectrum (toluene- d_8 , 253 K) of silyl toluenium ion $8(\text{C}_7\text{D}_8)$ at 253 K in toluene synthesized by hydride transfer from silane 15.

substance and additional characterization. These NMR experiments demonstrated the clean conversion of silylgermyl toluenium ion $3(\text{C}_7\text{D}_8)$ to germylsilyl toluenium $8(\text{C}_7\text{D}_8)$ in toluene at -20°C . According to our previous computational study, this sila-variant of the Wagner–Meerwein rearrangement is expected to proceed via several cationic intermediates (see Scheme 2).¹⁴ Additional DFT computations at the B3LYP/6-311+G(d,p) level of theory^{18,19} indicated that complexation of the formed silyl and germyl cations with the arene solvent has no significant influence on the calculated reaction coordinate. The calculated relative energies of the toluene complexes of cations 3–8 reveal only slight modifications of the previously

predicted stability sequence for the free silylium and germylium ions (see Figure S22, Supporting Information). In particular, the silyltoluenium ion $8(\text{C}_7\text{H}_8)$ is predicted to be the most stable along the complete reaction sequence, and the overall reaction $3(\text{C}_7\text{H}_8) \rightarrow 8(\text{C}_7\text{H}_8)$ is found to be exothermic by 44 kJ mol^{-1} . Experimentally, however, no indications for the cationic intermediates 4–7 or for the related arenium ions were detected. In order to test our mechanistic proposal, we decided to generate one specific cation along the reaction cascade, shown in Scheme 2, and to monitor its rearrangement. Therefore, we subjected silylgermane 16 to the hydride transfer reaction in toluene in an attempt to generate the corresponding germyltoluenium $7(\text{C}_7\text{D}_8)$ (Scheme 5). This last step of the reaction sequence, $7(\text{C}_7\text{H}_8) \rightarrow 8(\text{C}_7\text{H}_8)$, is predicted by the calculations to be exothermic by 15 kJ mol^{-1} (Figure S22, Supporting Information). $^{29}\text{Si}\{^1\text{H}\}$ NMR spectra recorded after 15 min at -20°C showed no signals expected for cation $7(\text{C}_7\text{H}_8)$ (compare Figure S19c (Supporting Information) and calculated values in Table 1), but the characteristic resonances for silyltoluenium $8(\text{C}_7\text{D}_8)$ already dominated. In agreement with the computational results, this experiment shows that germyltoluenium ion $7(\text{C}_7\text{D}_8)$ undergoes a fast 1,2-shift of the trimethylsilyl group to form the more stable cation $8(\text{C}_7\text{D}_8)$.

Obviously, the barrier for this rearrangement is rather low and in a similar range as the 18 kJ mol^{-1} that is predicted for the transformation of the free cations $7 \rightarrow 8$ (Figure S22, Supporting Information),¹⁴ and therefore, germyltoluenium ion $7(\text{C}_7\text{D}_8)$ cannot be detected even at temperatures as low as -20°C .

CONCLUSIONS

We were able to synthesize polysilanyl-substituted silyl and germyl cations, such as silyl cation 8 and germyl cation 3 in the form of their arene complexes at low temperatures using the weakly coordinating $[\text{B}(\text{C}_6\text{F}_5)_4]^-$ anion. The cation synthesis was done using either the stoichiometric reaction of germa-oligosilanes with a cationic Lewis acid such as tri-*iso*-propylsilyltoluenium, or, in a more regio-controlled way, by the standard hydride transfer reaction between hydrido-silanes and -germanes with the trityl cation. The cationic species were identified by low-temperature NMR spectroscopy supported by the results of quantum mechanical NMR chemical shift calculations. Applying these techniques, we were able to verify several silyl- and germyl cations and their solvent complexes as intermediates in the sila-Wagner–Meerwein rearrangement of silagermane 1 (eq 1, Scheme 1) and to provide strong evidence for their clean interconversion at low temperatures. We are convinced that this reaction is archetypical for many synthetic useful Lewis acid catalyzed skeletal rearrangements in polysilanes and polygermasilanes. Additional stable ion studies for sila-Wagner–Meerwein rearrangement of oligosilanes of higher complexity are currently under investigation in our laboratory to identify general rules and reaction patterns.

EXPERIMENTAL SECTION

All manipulations of air- and moisture-sensitive compounds were carried out under argon or nitrogen atmosphere using Schlenk techniques or a standard glovebox (Braun Unilab). Glassware was dried in an oven at 120°C and evacuated prior to use. The solvents tetrahydrofuran (THF), dimethoxyethane (DME), *n*-pentane, benzene, and toluene were dried over sodium and distilled under nitrogen atmosphere. Chlorobenzene was dried over CaCl_2 and stored over molecular sieves. Deuterated benzene and toluene were stored over

molecular sieves after drying over sodium. Dichlorodimethylgermane, chlorodimethylsilane, triethylsilane, tri-*iso*-propylsilane and trimethylchlorosilane were obtained from commercial suppliers, and the silanes were dried over molecular sieves. Sodium methanolate was prepared by addition of sodium to an excess of *abs.* methanol. After all sodium was consumed, the solvent was removed *in vacuo*. Triphenylmethyl tetrakis(pentafluorophenyl) borate **9** ($[\text{Ph}_3\text{C}][\text{B}(\text{C}_6\text{F}_5)_4]$) was prepared according to a modified literature procedure.²⁴ Tetrakis(trimethylsilyl)silane,²⁵ tetrakis(trimethylsilyl)germane²⁶ **2**, tris(trimethylsilyl)silypotassium,²⁷ tris(trimethylsilyl)germylpotassium,²⁸ tris(trimethylsilyl)silyltrimethylgermane²⁹ **1**, chloropentamethyldisilane,³⁰ and trimethylsilane³¹ were synthesized according to reported procedures. GC-MS spectra were performed on a Thermo Focus DSQ. NMR spectra were recorded on Bruker Avance 500, Avance III 500 and Varian Inova 300 spectrometers. ^1H NMR spectra were calibrated against the residual proton signal of the solvent as internal reference (benzene- $d_6/\delta^1\text{H}(\text{C}_6\text{D}_6\text{H}) = 7.20$; toluene- $d_8/\delta^1\text{H}(\text{CD}_2\text{H}) = 2.08$; chloroform- $d_1/\delta^1\text{H}(\text{CHCl}_3) = 7.24$; chlorobenzene- $d_5/\delta^1\text{H}(\text{C}_6\text{D}_4\text{HCl}) = 7.14$) and ^{13}C NMR spectra by using the central line of the solvent signal (benzene- $d_6/\delta^{13}\text{C}(\text{C}_6\text{D}_6) = 128.0$, toluene- $d_8/\delta^{13}\text{C}(\text{C}_6\text{D}_5\text{CD}_3) = 20.4$, chloroform- $d_1/\delta^{13}\text{C}(\text{CDCl}_3) = 77.0$, chlorobenzene- $d_5/\delta^{13}\text{C}(\text{C}_6\text{D}_5\text{Cl}) = 134.2$). $^{29}\text{Si}\{^1\text{H}\}$ NMR spectra were calibrated against an external standard ($^{29}\text{Si}(\text{Me}_2\text{SiHCl}) = 11.1$ versus tetramethylsilane (TMS)). The $^{29}\text{Si}\{^1\text{H}\}$ NMR inverse gated spectra were recorded with a relaxation delay $D1 = 10$ s. On the basis of our experiences, at -20°C this delay is long enough to allow a reliable integration of the peaks. The $^{29}\text{Si}\{^1\text{H}\}$ INEPT spectra were recorded with delays $D3 = 8.4$ ms and $D4 = 31.3$ ms. IR spectra were recorded on a Bruker Tensor 27 instrument. Analysis values for carbon show often too low values, which we attribute to the formation and incomplete combustion of silicon carbide, although vanadium pentoxide as combustion aid was used.

Dimethoxydimethylgermane (17). This compound was prepared according to slightly modified literature procedures.³² NaOMe (2.84 g) (3.5 eq., 52.57 mmol) was suspended in 40 mL of pentane, and 1.74 mL (15.00 mmol) of dichlorodimethylgermane was slowly added with a syringe. The mixture was stirred overnight at room temperature. The excess of NaOMe and formed NaCl were separated from the solution by using a centrifuge (20 min, 2000 rpm), and then the product-containing pentane solution was decanted using a Teflon tube. The salts were washed with 10 mL of pentane and again centrifuged and decanted. The pentane solutions were combined, and the product was separated from the solvent by fractionated distillation (bp.: 118°C at normal pressure (1.45 g, 58%). ^1H NMR (500.13 MHz, 297.9 K, C_6D_6 , δ): 0.30 (s, 6H, $(\text{CH}_3)_2\text{Ge}$), 3.54 (s, 6H, $\text{Ge}(\text{OCH}_3)_2$). $^{13}\text{C}\{^1\text{H}\}$ NMR (125.77 MHz, 298.1 K, C_6D_6 , δ): -2.9 ($(\text{CH}_3)_2\text{Ge}$), 51.6 ($\text{Ge}(\text{OCH}_3)_2$). Mass required for $\text{C}_4\text{H}_{12}\text{GeO}_2$: 166.0. Mass found GC/MS: 164.9 (0.5) $[\text{M}^+-\text{H}]$, 150.8 (100) $[\text{M}^+-\text{Me}]$, 135.9 (72) $[\text{M}^+-\text{OMe}]$, 120.9 (88) $[\text{M}^+-\text{OMe-Me}]$, 104.9 (84) $[\text{M}^+-\text{OMe-2Me}]$.

Tris(trimethylsilyl)silyldimethylmethoxygermane (18). A solution of 4.50 mmol tris(trimethylsilyl)silypotassium²⁷ in 40 mL of pentane and a solution of 0.75 g (4.50 mmol) of dimethoxydimethylgermane **17** in 10 mL of pentane were cooled to 0°C . The silyl potassium compound was added dropwise to the germane solution. The ice bath was allowed to warm to room temperature overnight. The reaction mixture was then hydrolyzed with 1 M hydrochloric acid. The organic layer was separated and dried over sodium sulfate. The solvent was removed under reduced pressure, and the product was purified by Kugelrohr distillation (0.62 g, 36%). Because of the use of hydrochloric acid, about 14% of the corresponding germyl chloride was formed as a byproduct, which was detected in the GC chromatograms and NMR spectra. ^1H NMR (499.87 MHz, 305.0 K, C_6D_6 , δ): 0.34 (s, 27H, $(\text{CH}_3)_3\text{Si}$), 0.60 (s, 6H, $(\text{CH}_3)_2\text{Ge}$), 3.54 (s, 3H, CH_3OGe). $^{13}\text{C}\{^1\text{H}\}$ NMR (125.69 MHz, 305.0 K, C_6D_6 , δ): 2.6 ($(\text{CH}_3)_3\text{Si}$), 4.1 ($(\text{CH}_3)_2\text{Ge}$), 52.6 (CH_3OGe). $^{29}\text{Si}\{^1\text{H}\}$ INEPT NMR (99.31 MHz, 305.0 K, C_6D_6 , δ): -124.6 ($(\text{CH}_3)_3\text{Si}$), -9.9 ($(\text{CH}_3)_3\text{Si}$). Mass required for $\text{C}_{12}\text{H}_{36}\text{GeOSi}_4$: 382.1. Mass found GC/MS: 367.2 (1) $[\text{M}^+-\text{Me}]$, 351.2 (0.5) $[\text{M}^+-\text{OMe}]$, 278.0 (8) $[\text{M}^+-\text{SiMe}_3-\text{OMe}]$, 205.1 (13) $[\text{M}^+-$

$2\text{SiMe}_3-\text{OMe}]$, 73.0 (100) $[\text{Me}_3\text{Si}^+]$. No satisfactory combustion analysis was available due to contamination with the chloride side-product.

Tris(trimethylsilyl)silyldimethylgermane (14). A solution of 0.62 g (1.62 mmol) tris(trimethylsilyl)silyldimethylmethoxygermane **18** in 30 mL of THF and a suspension of 0.062 g (1.62 mmol) of LiAlH_4 in 50 mL of THF were cooled to 0°C with an ice bath. The solution of silagermane **18** was added to the LiAlH_4 suspension and the reaction mixture was stirred for 20 min at 0°C before it was allowed to warm to room temperature and stirred for another 20 min. The mixture was slowly added to ice cold 2 M sulfuric acid. The phases were separated, and the aqueous phase was extracted two times with 50 mL of diethyl ether. The combined organic phases were dried over sodium sulfate and filtered, and the solvent was removed under reduced pressure. The product was crystallized from ethanol as a waxy, colorless solid (0.38 g, 1.09 mmol, 67%). ^1H NMR (499.87 MHz, 305.0 K, C_6D_6 , δ): 0.30 (s, 27H, $(\text{CH}_3)_3\text{Si}$), 0.50 (d, $^3J_{\text{H-H}} = 4.2$ Hz, 6H, $(\text{CH}_3)_2\text{Ge}$), 4.04 (sept, $^3J_{\text{H-H}} = 4.2$ Hz, 1H, GeH). $^{13}\text{C}\{^1\text{H}\}$ NMR (125.69 MHz, 305.0 K, C_6D_6 , δ): -2.3 ($(\text{CH}_3)_2\text{Ge}$), 2.5 ($(\text{CH}_3)_3\text{Si}$). $^{29}\text{Si}\{^1\text{H}\}$ NMR (99.31 MHz, 305.0 K, C_6D_6 , δ): -128.3 ($(\text{CH}_3)_3\text{Si}$), -9.4 ($(\text{CH}_3)_3\text{Si}$). Mass required for $\text{C}_{11}\text{H}_{34}\text{GeSi}_4$: 352.1. Mass found GC/MS: 351.1 (0.1) $[\text{M}^+-\text{H}]$, 337.1 (0.6) $[\text{M}^+-\text{Me-H}]$, 278.0 (30) $[\text{M}^+-\text{SiMe}_3-\text{H}]$, 189.1 (13) $[\text{M}^+-2\text{SiMe}_3-\text{Me-H}]$, 174.0 (4) $[\text{M}^+-\text{SiMe}_3-\text{GeMe}_2\text{H}]$, 73.1 (100) $[\text{Me}_3\text{Si}^+]$. IR (ATR, neat): $\nu_{\text{Ge-H}}$ 1982 cm^{-1} . Anal. found/calcd. for $\text{C}_{11}\text{H}_{34}\text{GeSi}_4$: C 37.63/37.60, H 10.67/9.75.

Tris(trimethylsilyl)germyldimethylsilane (15).³⁴ Solutions of 2.99 mmol tris(trimethylsilyl)germylpotassium²⁸ in 30 mL of DME and of 0.6 mL (excess, 5.52 mmol) of chlorodimethylsilane in 30 mL of DME were cooled to 0°C with an ice bath. The germypotassium compound was slowly added to the chlorosilane solution during 1 h. The ice bath was allowed to warm to room temperature overnight. The reaction mixture was then hydrolyzed with 1 M sulfuric acid. The organic layer was separated, and the aqueous phase was extracted with 10 mL of diethyl ether. The combined organic phases were dried over sodium sulfate, and the filtrate was concentrated to 5 mL under reduced pressure. The product was crystallized by adding 2 mL of acetonitrile as a colorless, waxy solid (0.847 g, 80.6%). ^1H NMR (499.87 MHz, 305.0 K, CDCl_3 , δ): 0.22 (s, 27H, $(\text{CH}_3)_3\text{Si}$), 0.27 (d, $^3J_{\text{H-H}} = 4.2$ Hz, 6H, $(\text{CH}_3)_2\text{Si}$), 4.12 (sept, $^3J_{\text{H-H}} = 4.2$ Hz, 1H, SiH). $^{13}\text{C}\{^1\text{H}\}$ NMR (125.71 MHz, 305.0 K, CDCl_3 , δ): -1.4 ($(\text{CH}_3)_2\text{Si}$), 3.1 ($(\text{CH}_3)_3\text{Si}$). ^{29}Si INEPT NMR (99.31 MHz, 305.0 K, CDCl_3 , δ): -29.8 (dsept, $^1J_{\text{Si-H}} = 180.5$ Hz, $^2J_{\text{Si-H}} = 7.0$ Hz, SiH), -4.7 ($(\text{CH}_3)_3\text{Si}$). Mass required for $\text{C}_{11}\text{H}_{34}\text{GeSi}_4$: 352.1. Mass found GC/MS: m/z (%) = 351.1 (0.3) $[\text{M}^+-\text{H}]$, 337.1 (2.5) $[\text{M}^+-\text{Me-H}]$, 278.1 (64) $[\text{M}^+-\text{SiMe}_3-\text{H}]$, 189.9 (22) $[\text{M}^+-2\text{SiMe}_3-\text{Me-H}]$, 174.0 (2) $[\text{M}^+-\text{SiMe}_3-\text{GeMe}_2\text{H}]$, 73.0 (100) $[\text{Me}_3\text{Si}^+]$. IR (ATR, neat): $\nu_{\text{Si-H}}$ 2085 cm^{-1} . Anal. found/calcd. for $\text{C}_{11}\text{H}_{34}\text{GeSi}_4$: C 36.31/37.60, H 9.98/9.75.

Tris(trimethylsilyl)pentamethyldisilanylgermane (19). A solution of 1.37 mmol tris(trimethylsilyl)germylpotassium·18-crown-6²⁸ in 3 mL of benzene was added dropwise to a solution of 0.25 g (1.51 mmol) of chloropentamethyldisilane³⁰ in 3 mL of benzene. After 5 h, the solution mixture was quenched with 1 M sulfuric acid, and the phases were separated. The aqueous phase was extracted with pentane, and the combined organic phases were dried over sodium sulfate and filtered, and the solvent was removed under reduced pressure. The product was obtained as colorless crystals by crystallization from methanol/diethyl ether 1:2 (0.42 g, 73%). ^1H NMR (299.94 MHz, 298.0 K, C_6D_6 , δ): 0.22 (s, 9H, $\text{Si}(\text{CH}_3)_3$), 0.36 (s, 27H, $(\text{CH}_3)_3\text{Si}$), 0.40 (s, 6H, $\text{Si}(\text{CH}_3)_2$), 0.40 (s, 6H, $\text{Si}(\text{CH}_3)_2$). $^{13}\text{C}\{^1\text{H}\}$ NMR (75.43 MHz, 298.0 K, C_6D_6 , δ): -0.8 ($\text{Si}(\text{CH}_3)_2$), -0.4 ($\text{Si}(\text{CH}_3)_3$), 4.0 ($(\text{CH}_3)_3\text{Si}$). $^{29}\text{Si}\{^1\text{H}\}$ INEPT NMR (59.59 MHz, 295.0 K, C_6D_6 , δ): -34.0 ($\text{Si}(\text{CH}_3)_2$), -15.5 ($\text{Si}(\text{CH}_3)_3$), -5.2 ($(\text{CH}_3)_3\text{Si}$). Mass required for $\text{C}_{14}\text{H}_{42}\text{GeSi}_5$: 424.1. Mass found GC/MS: m/z (%) = 424 (1) $[\text{M}^+]$, 408 (1) $[\text{M}^+-\text{Me-H}]$, 351 (3) $[\text{M}^+-\text{SiMe}_3]$, 278 (10) $[\text{M}^+-2\text{SiMe}_3]$, 259 (1) $[\text{GeSi}_3\text{C}_7\text{H}_{17}^+]$, 243 (1) $[\text{GeSi}_3\text{C}_6\text{H}_{13}^+]$, 219 (3) $[\text{GeSi}_2\text{C}_6\text{H}_{17}^+]$, 203 (11) $[\text{M}^+-3\text{SiMe}_3-2\text{H}]$, 187(8) $[\text{M}^+-3\text{SiMe}_3-\text{Me-4H}]$;

147 (7) [GeSiMe₃⁺]; 131 (35) [SiMe₃SiMe₂⁺]; 73 (100) [SiMe₃⁺]. Anal. found/calcd. for C₁₄H₄₂GeSi₃ C 39.33/39.70, H 9.50/9.99.

Bis(trimethylsilyl)pentamethyldisilanylgermane (16). A mixture of 0.21 g (0.49 mmol) of germapolysilane **19**, 0.062 g (0.51 mmol) of KO^tBu, and 0.134 g (0.51 mmol) of 18-crown-6 ether was dissolved in 2 mL of benzene. After the complete formation of germlypotassium compound **20** was confirmed by NMR spectroscopy, the solution was added to a stirred mixture of 10 mL of degassed diethyl ether and 20 mL of degassed 2 M sulfuric acid cooled with an ice bath. The phases were separated, the aqueous phase was extracted with degassed diethyl ether, and the combined organic phases dried over sodium sulfate. The solvents were removed under reduced pressure, and the product was obtained as a colorless oil (0.15 g, 91%). The compound is sensitive to oxygen and should be stored under argon at −20 °C.

Germlypotassium Compound 18-Crown-6 (20). ¹H NMR (299.94 MHz, 298.0 K, C₆D₆, δ ppm): 0.39 (s, 9H, Si(CH₃)₂Si(CH₃)₃), 0.59 (s, 18H, ((CH₃)₃Si)₂GeK), 0.64 (s, 6H, Si(CH₃)₂Si(CH₃)₃), 3.25 (s, 24H, CH₂O). ¹³C{¹H} NMR (75.43 MHz, 295.0 K, C₆D₆, δ ppm): 0.0 (Si(CH₃)₂Si(CH₃)₃), 3.5 (Si(CH₃)₂Si(CH₃)₃), 8.5 (((CH₃)₃Si)₂GeK), 70.0 (CH₂O). ²⁹Si{¹H} INEPT NMR (59.59 MHz, 295.0 K, C₆D₆, δ ppm): −33.0 (Si(CH₃)₂Si(CH₃)₃), −16.6 (Si(CH₃)₂Si(CH₃)₃), −3.4 (((CH₃)₃Si)₂GeK). Mass required for C₁₃H₃₈GeSi₄ after ethyl bromide derivatization: 380.1. Mass found after ethyl bromide derivatization GC/MS: *m/z* (%) = 380 (1) [M⁺]; 365 (1) [M⁺-Me]; 350 (3) [M⁺-Et-H]; 307 (1) [M⁺-SiMe₃]; 292 (1) [M⁺-SiMe₃-Me]; 277 (3) [M⁺-SiMe₃-2Me]; 262 (1) [M⁺-SiMe₃-3Me]; 234 (3) [M⁺-2SiMe₃]; 219 (8) [M⁺-2SiMe₃-Me]; 203 (17) [M⁺-2SiMe₃-2Me-H]; 187 (6) [M⁺-2SiMe₃-3Me-2H]; 159 (4) [M⁺-3SiMe₃-H]; 145 (10) [M⁺-3SiMe₃-Me]; 131 (33) [GeSiC₂H₅⁺]; 115 (9) [GeSiCH⁺]; 73 (100) [SiMe₃⁺].

Bis(trimethylsilyl)pentamethyldisilanylgermane (16). ¹H NMR (499.87 MHz, 305.1 K, C₆D₆, δ ppm): 0.22 (s, 9H, Si(CH₃)₂Si(CH₃)₃), 0.35 (s, 18H, ((CH₃)₃Si)₂Ge), 0.40 (s, 6H, Si(CH₃)₂Si(CH₃)₃), 2.25 (s, 1H, GeH). ¹³C{¹H} NMR (125.71 MHz, 305.0 K, CDCl₃, δ ppm): −1.7 (Si(CH₃)₂Si(CH₃)₃), −1.6 (Si(CH₃)₃Si(CH₃)₂), 3.1 (((CH₃)₃Si)₂Ge). ²⁹Si{¹H} NMR (99.31 MHz, 305.0 K, C₆D₆, δ ppm): −34.3 (Si(CH₃)₂Si(CH₃)₃), −16.2 (Si(CH₃)₂Si(CH₃)₃), −5.7 (((CH₃)₃Si)₂Ge). Mass required for C₁₁H₃₄GeSi₄: 352.1. Mass found GC/MS: 335 (2) [M⁺-Me-2H]; 278 (27) [M⁺-SiMe₃-H]; 263 (2) [M⁺-SiMe₃-Me-H]; 203 (13) [M⁺-2SiMe₃-2H]; 189 (10) [M⁺-2SiMe₃-Me-2H]; 173 (2) [M⁺-2SiMe₃-2Me-2H]; 131 (24) [SiMe₃SiMe₂⁺]; 115 (14) [Si₂C₄H₁₁⁺]; 73 (100) [SiMe₃⁺]. IR (ATR, neat) ν_{Ge-H} 1951 cm^{−1}. Anal. found/calcd. for C₁₁H₃₄GeSi₄ C 38.85/37.60, H 9.41/9.75.

General Preparation of Trialkylsilyl Arenium Borates (10a–d).³³ Triphenylmethyl tetrakis(pentafluorophenyl)borate was dissolved in 3 mL of the indicated solvent, and the silane was added. The formation of two phases could be observed, and the biphasic reaction mixture was vigorously stirred for 30 min. The upper, nonpolar phase was removed, and the lower, polar phase was washed with 2 mL of the used solvent, and again the nonpolar phase was removed. The polar phase was dried under reduced pressure for 30 min and then dissolved in the respective deuterated solvent.

Triethylsilyl Benzenium Borate (10a). Triphenylmethyl tetrakis(pentafluorophenyl)borate (0.50 g (0.54 mmol)) was dissolved in 3 mL of benzene, and 0.14 mL (1.6 eq., 0.87 mmol) of triethylsilane was added.

Tri-*iso*-propylsilyl Toluenum Borate (10b). Triphenylmethyl tetrakis(pentafluorophenyl)borate (0.46 g (0.50 mmol)) was dissolved in 3 mL of toluene, and 0.11 mL (1.1 eq., 0.55 mmol) of tri-*iso*-propylsilane was added.

Trimethylsilyl Toluenum Borate (10c). Triphenylmethyl tetrakis(pentafluorophenyl)borate (0.46 g (0.50 mmol)) was dissolved in 3 mL of toluene, and 0.06 mL (1.1 eq., 0.55 mmol) of trimethylsilane was added.

General Procedure for the Rearrangement of Tris(trimethylsilyl)silyltrimethylgermane (1) with Trialkylsilyl Arenium Borates (10a–c). A solution of 0.18 g (0.50 mmol) of silagermane **1** in 1 mL of the named deuterated solvent was added to a

precooled solution of the named trialkylsilyl arenium borate **10a–c**. The reaction mixture was stirred for 2 h at the specified temperature and then allowed to warm to room temperature. The polar phase and the nonpolar phase were each transferred to separate NMR tubes to be analyzed independently. In the following reactions, the NMR spectra of the polar phase showed too many signals to be analyzable, but the compounds in the nonpolar phase were identified by NMR and GC/MS spectroscopy (see Figures S8–11 in the [Supporting Information](#) for details).

Rearrangement of Tris(trimethylsilyl)silyldimethylgermyl Toluenum Borate (3(C₇D₈)[B(C₆F₅)₄]) to Tris(trimethylsilyl)germyldimethylsilyl Toluenum Borate (8(C₇D₈)[B(C₆F₅)₄]) Starting from Silagermane (1). To a solution of 0.18 g (0.50 mmol) of tris(trimethylsilyl)silyltrimethylgermane **1** in 1 mL of toluene-*d*₈ cooled to −20 °C, 1 equiv of tri-*iso*-propylsilyl toluenum borate **10b** was slowly added via a Teflon tube. The mixture was stirred for 1 h at −20 °C. The brown polar phase and the light yellow nonpolar phase were each transferred to separate NMR tubes at −20 °C and stored at −60 °C overnight until the NMR spectra were recorded the next morning. The polar phase contained borates [3(C₇D₈)] [B(C₆F₅)₄] and [8(C₇D₈)] [B(C₆F₅)₄]. The nonpolar phase contained methyl-tri-*iso*-propylsilane and the rearrangement product **2**. **Polar phase** ¹H NMR (499.87 MHz, 253.0 K, C₇D₈, δ ppm): −0.15 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 0.07 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 0.12 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), 0.28 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺). ¹³C{¹H} NMR (125.71 MHz, 253.0 K, C₇D₈, δ ppm): 1.6 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), 2.5 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 5.5 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 12.1 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺). ²⁹Si{¹H} NMR (99.31 MHz, 253.0 K, C₇D₈, δ ppm): −87.9 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), −7.9 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), −2.4 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 98.1 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺) (see Figures S12a–c in the [Supporting Information](#)). **Nonpolar phase** ¹H NMR (499.87 MHz, 305.0 K, C₇D₈, δ ppm): −0.17 (s, ((CH₃)₂CH)₃SiCH₃), 0.26 (s, ((CH₃)₃Si)₄Ge), 0.86 (sept, ³J_{H,H} = 7.3 Hz, ((CH₃)₂CH)₃SiCH₃), 0.97 (d, ³J_{H,H} = 7.3 Hz, ((CH₃)₂CH)₃SiCH₃). ¹³C{¹H} NMR (127.71 MHz, 305.0 K, C₇D₈, δ ppm): −10.1 (((CH₃)₂CH)₃SiCH₃), 3.5 (((CH₃)₃Si)₄Ge), 11.8 (((CH₃)₂CH)₃SiCH₃), 18.8 (((CH₃)₂CH)₃SiCH₃). ²⁹Si{¹H} NMR (99.31 MHz, 305.0 K, C₇D₈, δ ppm): −5.1 (((CH₃)₃Si)₄Ge), 9.0 (((CH₃)₂CH)₃SiCH₃) (see Figures S13a–d in the [Supporting Information](#)).

Rearrangement of Tris(trimethylsilyl)silyldimethylgermyl toluenum Borate (3(C₇D₈)[B(C₆F₅)₄]) to Tris(trimethylsilyl)germyldimethylsilyl Toluenum Borate (8(C₇D₈)[B(C₆F₅)₄]) Starting from Hydrogen Substituted Silagermane (14). tris(trimethylsilyl)silyldimethylgermane **14** (0.14 g (0.40 mmol)) and 0.37 g (0.40 mmol) of triphenylmethyl tetrakis(pentafluorophenyl)borate were each dissolved in 1 mL of toluene-*d*₈ and cooled to −20 °C. Silagermane **14** was slowly added to the borate salt via a Teflon tube, and the mixture was stirred at −20 °C for 1.5 h. The brown polar phase and the light yellow nonpolar phase were each transferred to NMR tubes at −20 °C. The NMR spectra were recorded at −20 °C. The polar phase contained borates [3(C₇D₈)] [B(C₆F₅)₄] and [8(C₇D₈)] [B(C₆F₅)₄]. The nonpolar phase contained triphenylmethane and the rearrangement product **2**. **Polar phase** ¹H NMR (499.87 MHz, 253.0 K, C₇D₈, δ ppm): −0.15 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 0.07 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), 0.12 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 0.28 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺). ¹³C{¹H} NMR (125.71 MHz, 253.0 K, C₇D₈, δ ppm): 1.6 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), 2.4 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 5.5 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 12.1 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺). ²⁹Si{¹H} NMR (99.31 MHz, 253.0 K, C₇D₈, δ ppm): −87.9 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), −7.9 (((CH₃)₃Si)₃SiGe(CH₃)₂⁺), −2.4 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺), 98.1 (((CH₃)₃Si)₃GeSi(CH₃)₂⁺) (see Figures S14a–c and S15a–b in the [Supporting Information](#)). **Nonpolar phase** ¹H NMR (499.87 MHz, 305.0 K, C₇D₈, δ ppm): 0.30 (s, ((CH₃)₃Si)₄Ge), 5.38 (s, Ph₃CH), 6.98–7.09 (m, Ph₃CH). ¹³C{¹H} NMR (127.71 MHz, 305.0 K, C₇D₈, δ ppm): 3.5 (((CH₃)₃Si)₄Ge), 57.1 (Ph₃CH), 125.4 (Ph₃CH), 128.3 (Ph₃CH), 129.2 (Ph₃CH), 144.3 (Ph₃CH). ²⁹Si{¹H} NMR (99.31

MHz, 305.0 K, C_7D_8 , δ ppm): -5.1 ($((CH_3)_3Si)_4Ge$) (see Figures S16a–c in the Supporting Information).

Tris(trimethylsilyl)germyldimethylsilyl Toluonium Borate ($8(C_7H_9)[B(C_6F_5)_4]$) from Silane 15. Tris(trimethylsilyl)germyldimethylsilane **15** (0.18 g (0.50 mmol)) and 0.46 g (0.50 mmol) of triphenylmethyl tetrakis(pentafluorophenyl)borate were both dissolved in 1 mL of toluene- d_8 and cooled to -20 °C. Germysilane **15** was slowly added to the borate salt via a Teflon tube, and the mixture was stirred at -20 °C for 1.5 h. The brown polar phase and the light yellow nonpolar phase were each transferred to separate NMR tubes at -20 °C and stored at -60 °C overnight until the NMR spectra were recorded the next morning. The polar phase contained borate $[8(C_7D_8)][B(C_6F_5)_4]$. The nonpolar phase contained triphenylmethane and the rearrangement product **2**. **Polar phase** 1H NMR (499.87 MHz, 253.0 K, C_7D_8 , δ ppm): -0.15 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$), 0.12 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$). $^{13}C\{^1H\}$ NMR (125.71 MHz, 253.0 K, C_7D_8 , δ ppm): 2.5 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$), 5.5 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$). $^{29}Si\{^1H\}$ NMR (99.31 MHz, 253.0 K, C_7D_8 , δ ppm): -2.4 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$), 98.1 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$) (see Figures S17a–c in the Supporting Information). **Nonpolar phase** 1H NMR (499.87 MHz, 305.0 K, C_7D_8 , δ ppm): 0.29 (s, $((CH_3)_3Si)_4Ge$), 5.38 (s, Ph_3CH), 6.98 – 7.10 (m, Ph_3CH). $^{13}C\{^1H\}$ NMR (125.69 MHz, 305.0 K, C_7D_8 , δ ppm): 3.5 ($((CH_3)_3Si)_4Ge$), 57.2 (Ph_3CH), 125.4 (Ph_3CH), 128.3 (Ph_3CH), 129.2 (Ph_3CH), 144.3 (Ph_3CH). $^{29}Si\{^1H\}$ NMR (99.31 MHz, 305.0 K, C_7D_8 , δ ppm): -5.1 ($((CH_3)_3Si)_4Ge$) (see Figure S18a–c in the Supporting Information).

Tris(trimethylsilyl)germyldimethylsilyl-phenylchloronium Borate ($8(C_6D_5Cl)[B(C_6F_5)_4]$) from Silane 15. Tris(trimethylsilyl)germyldimethylsilane **15** (0.09 g (0.25 mmol)) and 0.23 g (0.25 mmol) of triphenylmethyl tetrakis(pentafluorophenyl)borate were both dissolved in 0.5 mL of chlorobenzene- d_5 and cooled to -20 °C. Germysilane **15** was slowly added to the borate salt via a Teflon tube, and the mixture was stirred at -20 °C for 1.5 h. The brown solution was transferred into a NMR tube at -20 °C and stored at -60 °C overnight until the NMR spectra were recorded the next morning. The mixture contained borate $[8(C_6D_5Cl)][B(C_6F_5)_4]$, the rearrangement product **2**, and triphenylmethane. $^{29}Si\{^1H\}$ NMR (99.31 MHz, 253.0 K, C_6D_5Cl , δ ppm): -5.1 ($((CH_3)_3Si)_4Ge$), -1.4 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$), 154.3 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$).

Tris(trimethylsilyl)germyldimethylsilyl Toluonium Borate ($8(C_7D_8)[B(C_6F_5)_4]$) from Germene 16. Bis(trimethylsilyl)pentamethyldisilanylgermane **16** (0.11 g (0.32 mmol)) and 0.29 g (0.32 mmol) of triphenylmethyl tetrakis(pentafluorophenyl)borate were both dissolved in 1 mL of toluene- d_8 and cooled to -20 °C. Silylgermane **16** was slowly added to the borate salt via a Teflon tube, and the mixture was stirred at -20 °C for 5 min. The brown polar phase and the light yellow nonpolar phase were each transferred to separate NMR tubes at -20 °C and stored at -60 °C for 5 h until the NMR spectra were recorded. At -60 °C, the polar phase solidifies, and no further reaction is expected. The NMR spectra of the polar phase recorded at -20 °C contained nearly exclusively borate $[8(C_7D_8)][B(C_6F_5)_4]$. The nonpolar phase contained triphenylmethane and the rearrangement product **2**. **Polar phase** 1H NMR (499.87 MHz, 253.0 K, C_7D_8 , δ ppm): -0.16 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$), 0.12 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$). $^{13}C\{^1H\}$ NMR (125.71 MHz, 253.0 K, C_7D_8 , δ ppm): 2.5 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$), 5.5 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$). $^{29}Si\{^1H\}$ NMR (99.31 MHz, 253.0 K, C_7D_8 , δ ppm): -2.4 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$), 98.2 ($((CH_3)_3Si)_3GeSi(CH_3)_2^+$) (see Figures S19a–c in the Supporting Information). **Nonpolar phase** 1H NMR (499.87 MHz, 305.0 K, C_7D_8 , δ ppm): 0.30 (s, $((CH_3)_3Si)_4Ge$), 5.38 (s, Ph_3CH), 6.98 – 7.10 (m, Ph_3CH). $^{13}C\{^1H\}$ NMR (125.69 MHz, 305.0 K, C_7D_8 , δ ppm): 3.5 ($((CH_3)_3Si)_4Ge$), 57.1 (Ph_3CH), 125.4 (Ph_3CH), 128.2 (Ph_3CH), 129.1 (Ph_3CH), 144.3 (Ph_3CH). $^{29}Si\{^1H\}$ NMR (99.31 MHz, 305.0 K, C_7D_8 , δ ppm): -5.1 ($((CH_3)_3Si)_4Ge$) (see Figure S20a–c in the Supporting Information).

■ ASSOCIATED CONTENT

Supporting Information

Experimental and theoretical characterization, including the preparation of all compounds of interests and computational details. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.organomet.5b00431.

■ AUTHOR INFORMATION

Corresponding Authors

*(J.B.) E-mail: baumgartner@tugraz.at.

*(C.M.) E-mail: christoph.marschner@tugraz.at.

*(T.M.) E-mail: thomas.mueller@uni-oldenburg.de.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the ERA-chemistry program (DFG-Mu1440/8-1, FWF-I00669) and by Carl von Ossietzky University Oldenburg. The simulations were performed at the HPC Cluster HERO (High End Computing Resource Oldenburg), located at the University of Oldenburg (Germany) and funded by the DFG through its Major Research Instrumentation program (INST 184/108-1 FUGG) and the Ministry of Science and Culture (MWK) of the Lower Saxony State.

■ REFERENCES

- (1) Recent review: Marschner, C. *Struct. Bonding* **2014**, *155*, 163–288.
- (2) Recent review: Corey, J. Y. *Adv. Organomet. Chem.* **2004**, *51*, 1–52.
- (3) Ishifune, M.; Kashimura, S.; Kogai, Y.; Fukuhara, Y.; Kato, T.; Bu, H.-B.; Yamashita, N.; Murai, Y.; Murase, H.; Nishida, R. *J. Organomet. Chem.* **2000**, *611*, 26–31.
- (4) (a) Ishikawa, M.; Kumada, M. *J. Chem. Soc., Chem. Commun.* **1969**, 567–568. (b) Ishikawa, M.; Kumada, M. *J. Chem. Soc., Chem. Commun.* **1970**, 157. (c) Ishikawa, M.; Iyoda, J.; Ikeda, H.; Kotake, K.; Hashimoto, T.; Kumada, M. *J. Am. Chem. Soc.* **1981**, *103*, 4845–4850. (d) Ishikawa, M.; Watanabe, M.; Iyoda, J.; Ikeda, H.; Kumada, M. *Organometallics* **1982**, *1*, 317–322.
- (5) Blinks, A.; West, R. *Organometallics* **1986**, *5*, 128–133.
- (6) (a) Märkl, G.; Wagner, R. *Tetrahedron Lett.* **1986**, *27*, 4015–4018. (b) Sharma, S.; Caballero, N.; Li, H.; Pannell, K. H. *Organometallics* **1999**, *18*, 2855–2860.
- (7) Fischer, J.; Baumgartner, J.; Marschner, C. *Science* **2005**, *310*, 825.
- (8) Recent reviews: (a) Müller, T. *Struct. Bonding* **2014**, *155*, 107–162. (b) Müller, T. In *Science of Synthesis: Knowledge Updates 2013/3*; Oestreich, M., Ed.; Thieme: Stuttgart, Germany, 2013; pp 1–42. (c) Klare, H. F. T.; Oestreich, M. *Dalton Trans.* **2010**, *39*, 9176–9184. (d) Müller, T. *Adv. Organomet. Chem.* **2005**, *155*–215.
- (9) (a) Lambert, J. B.; Zhang, S. *J. Chem. Soc., Chem. Commun.* **1993**, 383–384. (b) Lambert, J. B.; Zhang, S. J.; Ciro, S. M. *Organometallics* **1994**, *13*, 2430–2443. (c) Ottosson, C.-H.; Cremer, D. *Organometallics* **1996**, *15*, 5495–5501. (d) Ottosson, C.-H.; Szabó, K. J.; Cremer, D. *Organometallics* **1997**, *16*, 2377–2385.
- (10) (a) Sekiguchi, A.; Matsuno, T.; Ichinohe, M. *J. Am. Chem. Soc.* **2000**, *122*, 11250–11251. (b) Inoue, S.; Ichinohe, M.; Yamaguchi, T.; Sekiguchi, A. *Organometallics* **2008**, *27*, 6056–6058.
- (11) (a) Ichinohe, M.; Igarashi, M.; Sanuki, K.; Sekiguchi, A. *J. Am. Chem. Soc.* **2005**, *127*, 9978–9979. (b) Igarashi, M.; Ichinohe, M.; Sekiguchi, A. *J. Am. Chem. Soc.* **2007**, *129*, 12660–12661.
- (12) Nakamoto, M.; Fukawa, T.; Sekiguchi, A. *Chem. Lett.* **2004**, *33*, 38–39.

- (13) Wagner, H.; Wallner, A.; Fischer, J.; Flock, M.; Baumgartner, J.; Marschner, C. *Organometallics* **2007**, *26*, 6704–6717.
- (14) Wagner, H.; Baumgartner, J.; Müller, T.; Marschner, C. *J. Am. Chem. Soc.* **2009**, *131*, 5022–5023.
- (15) Reviews: (a) Reed, C. A. *Acc. Chem. Res.* **2010**, *43*, 121–128. (b) Krossing, I.; Raabe, I. *Angew. Chem., Int. Ed.* **2004**, *43*, 2066–2090.
- (16) No silyl- or germlychlorides have been detected as by-products. Obviously, the rearrangement of the intermediate cations and alkylation of the final cation is faster than chloride abstraction from the solvent.
- (17) (a) Bartlett, P. D.; Condon, F. E.; Schneider, A. *J. Am. Chem. Soc.* **1944**, *66*, 1531–1539. (b) Corey, J. Y. *J. Am. Chem. Soc.* **1975**, *97*, 3237–3238.
- (18) All computations were performed using the Gaussian 09 package: Frisch, M. J. et al. *Gaussian 09*, revision B.01; Gaussian, Inc.: Wallingford, CT, 2009.
- (19) See [Supporting Information](#) for further details.
- (20) At temperatures lower than $-20\text{ }^{\circ}\text{C}$, the ionic phase became solid, which excludes further NMR investigations.
- (21) Krempner, C. *Polymers* **2002**, *4*, 408–447.
- (22) In chlorobenzene solution, usually the corresponding silylated chloronium ions predominate over the silylated arenium ion, see: (a) Schäfer, A.; Saak, W.; Haase, D.; Müller, T. *Angew. Chem., Int. Ed.* **2012**, *51*, 2881–2984. (b) Hoffmann, S. P.; Kato, T.; Tham, F. S.; Reed, C. A. *Chem. Commun.* **2006**, 767–769.
- (23) In addition, NMR detection of toluenium ion $8(\text{C}_7\text{D}_8)$ at temperatures around $0\text{ }^{\circ}\text{C}$ is severely hampered by significant line broadening of the ^{29}Si NMR resonances, most probably due to fast exchange of the toluene molecule in cation $8(\text{C}_7\text{D}_8)$ with the solvent (for example, the line width at half height, $w_{1/2}$ of the ^{29}Si NMR signal at $\delta^{29}\text{Si} = 98.1$ is $w_{1/2} = 17\text{ Hz}$ at $T = -20\text{ }^{\circ}\text{C}$ and $w_{1/2} = 40\text{ Hz}$ at $T = +10\text{ }^{\circ}\text{C}$).
- (24) (a) Massey, A. G.; Park, A. J. *J. Organomet. Chem.* **1964**, *2*, 245–250. (b) Chien, J. C. W.; Tsai, W. M.; Rausch, M. D. *J. Am. Chem. Soc.* **1991**, *113*, 8570–8571.
- (25) Gilman, H.; Smith, C. L. *J. Organomet. Chem.* **1967**, *8*, 245–253.
- (26) Brook, A. G.; Abdesaken, F.; Söllradl, H. *J. Organomet. Chem.* **1986**, *299*, 9–13.
- (27) Marschner, C. *Eur. J. Inorg. Chem.* **1998**, 1998, 221–226.
- (28) Fischer, J.; Baumgartner, J.; Marschner, C. *Organometallics* **2005**, *24*, 1263–1268.
- (29) Mallela, S. P.; Ghuman, M. A.; Geanangel, R. A. *Inorg. Chim. Acta* **1992**, *202*, 211–217.
- (30) Gluyas, J. B. G.; Burschka, C.; Kraft, P.; Tacke, R. *Organometallics* **2010**, *29*, 5897–5903.
- (31) Lehmann, M.; Schulz, A.; Villinger, A. *Angew. Chem., Int. Ed.* **2009**, *48*, 7444–7447.
- (32) (a) West, R.; Hunt, H. R.; Whipple, R. O. *J. Am. Chem. Soc.* **1954**, *76*, 310–310. (b) Moedritzer, K.; Van Wazer, J. R. *Inorg. Chem.* **1965**, *4*, 1753–1760.
- (33) Schäfer, A.; Schäfer, A.; Müller, T. *Dalton Trans.* **2010**, 39, 9296–9303.
- (34) Sharma, S.; Pannell, K. H. *Organometallics* **2000**, *19*, 1225.