#### Total Synthesis

### Total Synthesis of Ramonanins A-D\*\*

Ross S. Harvey, Emily G. Mackay, Lukas Roger, Michael N. Paddon-Row,\* Michael S. Sherburn,\* and Andrew L. Lawrence\*

Abstract: The first total synthesis of the ramonanin family of lignan natural products is described. The short synthesis involves a 2,5-diaryl-3,4-dimethylene tetrahydrofuran intermediate, which participates in an unexpectedly facile Diels-Alder dimerization, generating all four natural products. Insights into the reactivity and stereoselectivity of the key dimerization are provided through computational studies employing B3LYP/6-31G(d) and M06-2X/6-31G(d) model chemistries.

Guaiacum officinale and Guaiacum sanctum are the sources of the highly prized lignum vitae wood, known for its extraordinary strength and durability. The resin of lignum vitae, known as gum guaiac, or guaiac resin, is also valuable. It is reported to possess various medicinal properties,<sup>[1]</sup> and has also found application in colorimetric tests for oxidative conditions (e.g., Nobles' test,<sup>[2]</sup> Guaiac occult blood test,<sup>[3]</sup> and Schönbein–Pagenstecher test<sup>[4]</sup>).  $\alpha$ -Guaiaconic acid (1) is the constituent of gum guaiac that is readily oxidized, forming the vividly colored guaiacum blue (2; Scheme 1 a).<sup>[5]</sup> Sadly, the highly sought-after attributes of lignum vitae wood and gum guaiac have led to overexploitation of Guaiacum officinale, which is now listed as an endangered species on the Red List of the International Union for the Conservation of Nature.<sup>[6]</sup>

The ramonanins A-D (3-6) are a group of spirocyclic phenylpropanoid tetramers isolated from the heartwood of Guaiacum officinale by Schroeder and co-workers in 2011 (Scheme 1 b).<sup>[7]</sup> The ramonanins (3-6) formulate as dimers of  $\alpha$ -guaiaconic acid (1) and are reported to exhibit cytotoxic

[\*] R. S. Harvey, E. G. Mackay, L. Roger, Prof. M. S. Sherburn

	Research School of Chemistry, Australian National University Canberra, ACT 2601 (Australia) E-mail: michael.sherburn@anu.edu.au
	Prof. M. N. Paddon-Row School of Chemistry, The University of New South Wales Sydney, NSW 2052 (Australia) E-mail: m.paddonrow@unsw.edu.au
	Dr. A. L. Lawrence School of Chemistry, The University of Edinburgh Joseph Black Building, West Mains Road, Edinburgh, EH9 3JJ (UK) E-mail: a.lawrence@ed.ac.uk
[**]	This work was supported by the Australian Research Council and a Marie Curie Career Integration Grant. M.N.PR. acknowledges that this research was undertaken with the assistance of resources provided at the NCI National Facility through the National Computational Merit Allocation Scheme supported by the Austral- ian Government. The authors thank Prof. Schroeder (Cornell University) for kindly supplying samples of the ramonanin natural products and Tony Herlt (Australian National University) for assistance with HPLC separations.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201409818.

a) Structure and reactivity of α-guaiaconic acid (1)



Guaiac occult blood test

Used to detect the presence of blood in stool samples. Heme catalyzes the oxidation of 1

Nobles' test

Used to detect the presence of extracellular oxidase enzymes

Schönbein-Pagenstecher test

Used to detect the presence of HCN, which reacts with  $CuSO_4$  to generate  $O_3$ .

b) Schroeder's proposed biosynthesis of the ramonanins A-D (3-6)



**Scheme 1.** a) The structure and reactivity of  $\alpha$ -guaiaconic acid (1). b) Schroeder's proposed biosynthesis of the ramonanins A-D (3-6).<sup>[7]</sup>

activity against human breast cancer cell lines.<sup>[7]</sup> The endangered status of Guaiacum officinale, together with the promising biological activity of its natural products, are classic motivations for total synthesis efforts.<sup>[8]</sup> In truth, however, our attention was drawn to these molecules because of their intriguing structures, which are fascinating from both a topological and biosynthetic perspective. The spirocyclic core of the ramonanins is unprecedented among natural products and Schroeder and co-workers proposed a biosynthetic pathway that involved dimerization of a lignan precursor, 7, through a "Diels-Alder-like mechanism" (Scheme 1b).<sup>[7]</sup>

Schroeder's proposed biosynthetic intermediate 7 presumably also serves as the biosynthetic precursor to  $\alpha$ guaiaconic acid (1). There are no reports of 1,2-dimethylenecyclopentane-type structures undergoing Diels-Alder dimerizations under ambient conditions.<sup>[9]</sup> It is, therefore, reason-

Angew. Chem. Int. Ed. 2014, 53, 1-5

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

# Wiley Online Library

These are not the final page numbers!

Angewandte Communications



Scheme 2. Synthesis of lignan 7.

able to assume that lignan **7** would not undergo a spontaneous Diels–Alder dimerization.<sup>[10]</sup> Our previous studies on the kingianin natural products utilized a radical-cation-catalyzed formal Diels–Alder (RCDA) reaction to dimerize a thermally unreactive bicyclo[4.2.0]octadiene structure.<sup>[11]</sup> We considered that the ramonanins A–D (**3–6**) might also be the result of a biosynthetic RCDA dimerization.<sup>[12]</sup> To investigate the nature of this fascinating biosynthetic dimerization we chose to pursue a total synthesis of the ramonanin natural products (**3–6**).<sup>[13]</sup>

The synthesis began with the high yielding conversion of vanillin into its benzene sulfonate ester, a process that could be easily conducted on large scale (Scheme 2). Two molecules of the resulting aldehyde 8 were united through an acetylene unit in a one-pot sequence to afford multigram quantities of diacetate 9 as a 1:1 mixture of diastereomers. Following a strategy pioneered by Mori and co-workers,<sup>[14]</sup> alkyne 9 was subjected to an envne metathesis with ethylene using the Hoveyda–Grubbs II catalyst, to afford diene 10 on multigram scale. Deprotection of the secondary alcohols was accomplished using KOH in MeOH to afford diol 11. Camphor sulfonic acid was then used to promote 3,4-dimethylene tetrahydrofuran ring formation, which generated protected lignan 12 as a 2:1 mixture of diastereomers, both of which would be required for the synthesis of ramonanins A-D(3-6). This short synthetic sequence allowed access to multigram quantities of key intermediate 12, which could be stored at -15 °C for weeks without deterioration. When required, the benzene sulfonate protecting groups could be cleanly removed with NaOH in MeOH/THF to afford lignan 7 (Scheme 2).

To our surprise and delight, dimerization of lignan **7** was observed to occur spontaneously at room temperature, albeit slowly.<sup>[15]</sup> Under optimized conditions, dissolution of the monomer *cis*-**7**<sup>[16]</sup> in a minimal amount of DMF and warming at 50 °C provided, after chromatographic purification, a 54:9:37 mixture of ramonanins A (**3**), B (**4**), and C (**5**) in a combined yield of 45%, in addition to 24% recovered starting material *cis*-**7** (Scheme 3).<sup>[17]</sup> Preparative HPLC



Scheme 3. Dimerization of lignan cis-7.

allowed the isolation of analytically pure samples of ramonanins A–C (**3–5**) and the spectroscopic data for each synthetic compound matched those reported for the corresponding natural product, thus securing the total synthesis of ramonanins A–C (**3–5**). Optical rotation values for the natural products were reported in the isolation paper [**3**:  $[\alpha]_D^{22} - 2.2$ (*c* 0.6, MeOH); **4**:  $[\alpha]_D^{22} - 4.0$  (*c* 0.2, MeOH); **5**:  $[\alpha]_D^{22} - 5.2$  (*c* 0.2, MeOH); **6**:  $[\alpha]_D^{22} - 5.1$  (*c* 0.3, MeOH)].<sup>[7]</sup> Nevertheless, since ramonanins A–C are the Diels–Alder dimers of the achiral diene *cis-***7**, we strongly suspected that these natural products were racemates. To test this hypothesis, we performed chiral HPLC analysis on samples of ramonanins A and C, kindly provided by Professor Schroeder,<sup>[18]</sup> which confirmed their racemic nature.

Our attention now turned to the final member of the family: ramonanin D (6; Scheme 1). A unique stereochemical feature of ramonanin D (6) is the *trans* relationship between the two aryl substituents on the dihydrofuran ring. As would

www.angewandte.org

2

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!



**Scheme 4.** The eight stereoisomeric TSs for the dimerization of **13** to afford the ramonanin simulacra (ram A, ram B, ram C, and  $ram_{noe}$ ), with explanation of the TS notation.

be expected, upon warming a mixture of *cis*-7 and *trans*-7, complex mixtures of products were obtained. Nevertheless, an analytically pure sample of ramonanin D (6) was isolated, albeit in low (<1%) yield, after multiple HPLC purification steps.

Given the surprisingly facile nature of the Diels-Alder dimerization of *cis*-7, we chose to investigate the reactivity and stereochemical control of this process computationally, using the B3LYP/6-31G(d) and M06-2X/6-31G(d) model chemistries. The 2,5-diphenyl analogue of cis-7, namely 13 (Scheme 4), was used as a simulacrum.<sup>[19]</sup> Eight distinct stereochemical modes of the Diels-Alder dimerization, and hence eight transition structures (TSs), are possible, and these are depicted in Scheme 4. The TSs are described using a notation, in which the Alder-Stein mode may be either endo (N) or exo (X) and the pair of phenyl substituents on the diene and on the dienophile may be either anti (a) or syn (s) with respect to the bond-forming zone. The eight TSs comprise four pairs, both members of each pair giving one of the ramonanin simulacra adducts (ram X = simulacra of ramonanin X;  $ram_{noe} = not$  observed experimentally).

The computed product distribution for the Diels–Alder dimerization of **13** is presented in Table 1, together with the experimentally determined product distribution from dimerization of *cis*-**7** (Scheme 3). The agreement between the predicted and experimental adduct distribution is good and strongly suggests that not only is the model chemistry used in this study reliable, but also that the OH and OMe substituents in *cis*-**7** do not significantly affect the distribution of the ramonanin stereoisomers. As expected, the *Naa* and *Xaa* TSs are the most favorable, presumably because both pairs of phenyl groups (i.e., from diene and dienophile) occupy the *anti* space with respect to the bond-forming zone, with the former lying 3 kJ mol<sup>-1</sup> below the latter, in accordance with

**Table 1:** B3LYP/6-31G(d) and experimental product distributions [%] for the Diels–Alder dimerization of **13** and *cis*-**7**, respectively.

Adduct	Mode	Predicted	Total predicted	Experimental
ram A	Naa Xsa	58.6 0.1	59	54
ram B	Nas Xss	3.7 0	4	9
ram C	Nsa Xaa	17.4 17.8	35	37
ram <sub>noe</sub>	Nss Xas	0.1 2.3	2	-

Alder's *endo* rule. TSs with all phenyl groups positioned in the *syn* space, are 16 (*Nss*) and 24 kJ mol<sup>-1</sup> (*Xss*) above the *Naa* TS, reflecting steric congestion of the phenyl groups, as may be seen by comparing the geometries of *Naa* and *Nss* TSs in Figure 1. All TSs were found to be highly asynchronous, with the shorter of the newly formed bonds between 1.72 and 1.97 Å and the longer one between 3.04 and 3.69 Å, the most asynchronous being the *Nss* TS ( $\Delta r = 1.97$  Å, Figure 1).



*Figure 1.* Transition Structures *Naa* and *Nss* with bond lengths of the newly formed bonds.

It seems surprising that neat samples of *cis*-**7** readily dimerize, as does cyclopentadiene (CPD), given that the terminal methylene carbon atoms in **13** are calculated to be 3.139 Å apart, compared to only 2.363 Å in CPD.<sup>[20]</sup> The M06-2X computed activation enthalpies and free energies for the dimerization of CPD, 1,2-dimethylenecyclopentane, 3,4-dimethylenetetrahydrofuran, and **13** are listed in Table 2.

As may be seen from Table 2,  $\Delta H^{\pm}$  for Diels–Alder dimerization is approximately the same for all four reactants and, in particular, neither the presence of phenyl groups nor that of the ring oxygen has any significant effect on  $\Delta H^{\pm}$ (compare entries 3 with 4 and 2 with 3). However,  $\Delta G^{\pm}$  for the dimerization of **13** is about 10 kJ mol<sup>-1</sup> higher than in the other three systems, for which similar values are obtained. This free energy difference translates into a 50-fold decrease in the Diels–Alder dimerization rate for **13**, compared to the other dienes listed in Table 2. The origin of this effect is probably due to the phenyl groups in **13** experiencing a steeper potential for their torsional motion about the bonds connecting them to the tetrahydrofuran ring in the TS, compared to

www.angewandte.org

#### Angewandte Communications

**Table 2:**  $\Delta H^+$  and  $\Delta G^+$  values (kJ mol<sup>-1</sup>, 25 °C) for the Diels–Alder dimerization of various dienes.<sup>[a]</sup>

Entry	Reactant	$\Delta H^{+}$	$\Delta {\sf G}^{st}$
1		60.7	121.7
2		64.2	125.0
3		59.8	122.2
4	Ph O Ph 13	58.0	132.2

[a] M06-2X/6-31G(d)-[PCM = THf]//M06-2X/6-31G(d) activation energies from the lowest energy (*endo*) path.

the reactant. In conclusion, these calculations support the notion of a concerted cycloaddition event, both in the synthesis and biogenesis of the ramonanin natural products.

In summary, a seven-step total synthesis of the ramonanin natural products has been achieved. An investigation of the chemistry of Schroeder's proposed biosynthetic pathway has pinpointed the most likely origin of these fascinating structures. The natural products have been shown to be racemates. Our experimental results demonstrate that lignan **7** is predisposed toward thermal Diels–Alder dimerization, and our computational findings predict this behavior more widely in the 1,2-dimethylene cyclopentane series. It remains unclear whether this predisposed reactivity is being harnessed in nature constructively, or is generating products that are artifacts of isolation.

Received: October 6, 2014 Published online:

**Keywords:** biomimetic synthesis · Diels–Alder reactions · enyne metathesis · lignans · natural products

- a) R. S. Munger, J. Hist. Med. Allied Sci. 1949, IV, 196-229;
  b) J. F. Morton, Atlas of Medicinal Plants of Middle America: Bahamas to Yucatan, Charles C. Thomas, Springfield, 1981;
   c) J. L. Hartwell, Lloydia 1971, 34, 386-425;
   d) L. P. Somogyi, Food Additives, Wiley, New York, 2000.
- [2] M. K. Nobles, Can. J. Bot. 1958, 36, 91-99.
- [3] D. C. Rockey, N. Engl. J. Med. 1999, 341, 38-46.
- [4] W. Autenrieth, Laboratory manual for the detection of poisons and powerful drugs, 4th ed. (Ed.: W. H. Warren), Blakiston, Philadelphia, 1915, pp. 21-22.
- [5] The structure of guaiacum blue (2) in Scheme 1a is that shown in the following publication, but it should be noted that this compound has never been fully characterized, see: J. F. Kratochvil, R. H. Burris, M. K. Seikel, J. M. Harkin, *Phytochemistry* 1971, 10, 2529–2531.
- [6] "Guaiacum officinale L." Taxonomic Serial No.: 897104 Retrieved [07-18-2014], from the ITIS (Integrated Taxonomic

Information System) on-line database, http://www.itis.gov. Listed as endangered on the IUCN (International Union for the Conservation of Nature) Red List. Retrieved [07–18–2014], from http://www.iucnredlist.org.

- [7] K. J. Chavez, X. Feng, J. A. Flanders, E. Rodriguez, F. C. Schroeder, J. Nat. Prod. 2011, 74, 1293–1297.
- [8] For a discussion of the role of total synthesis in addressing the problem of natural product supply, see J. D. Keasling, A. Mendoza, P. S. Baran, *Nature* 2012, 492, 188–189.
- [9] This lack of literature precedent perhaps explains why Schroeder and co-workers described the key step in their proposed biosynthesis as having a "Diels-Alder-like mechanism" (Scheme 1). The most closely related example of this type of Diels-Alder reaction comes from Bloomquist and co-workers who, in 1956, synthesized 1,2-dimethylenecyclopentane and reported that attempts to distil the product at 97°C resulted in dimerization, see: A. T. Blomquist, J. Wolinsky, Y. C. Meinwald, D. T. Longone, J. Am. Chem. Soc. 1956, 78, 6057-6063.
- [10] For an excellent review on biosynthetic Diels-Alder reactions, see: E. M. Stocking, R. M. Williams, *Angew. Chem. Int. Ed.* 2003, 42, 3078-3115; *Angew. Chem.* 2003, 115, 3186-3223.
- [11] S. L. Drew, A. L. Lawrence, M. S. Sherburn, Angew. Chem. Int. Ed. 2013, 52, 4221–4224; Angew. Chem. 2013, 125, 4315–4318.
- [12] Relatively few examples of total syntheses containing proposed biomimetic radical cation Diels-Alder reactions have been reported, see Ref. [11] and: a) H. Cong, C. F. Becker, S. J. Elliot, M. W. Grinstaff, J. A. Porco, J. Am. Chem. Soc. 2010, 132, 7514–7518; b) S. Lin, M. A. Ischay, C. G. Fry, T. P. Yoon, J. Am. Chem. Soc. 2011, 133, 19350–19353; c) H. N. Lim, K. A. Parker, Org. Lett. 2013, 15, 398–401; d) C. Qi, H. Cong, K. J. Cahill, P. Müller, R. P. Johnson, J. A. Porco, Angew. Chem. Int. Ed. 2013, 52, 8345–8348; Angew. Chem. 2013, 125, 8503–8506; e) H. N. Lim, K. A. Parker, J. Org. Chem. 2014, 79, 919–926.
- [13] For our previous biomimetic studies on dimeric natural products, see Ref. [11] and: a) P. D. Brown, A. C. Willis, M. S. Sherburn, A. L. Lawrence, *Angew. Chem. Int. Ed.* **2013**, *52*, 13273–13275; *Angew. Chem.* **2013**, *125*, 13515–13517; b) P. D. Brown, A. C. Willis, M. S. Sherburn, A. L. Lawrence, *Org. Lett.* **2012**, *14*, 4537–4539.
- Selected examples: a) A. Kinoshita, N. Sakakibara, M. Mori, J. Am. Chem. Soc. 1997, 119, 12388-12389; b) K. Tonogaki, M. Mori, Tetrahedron Lett. 2002, 43, 2235-2238; c) M. Mori, K. Tonogaki, N. Nishiguchi, J. Org. Chem. 2002, 67, 224-226.
- [15] The half-lives of both diastereomers of lignan 7 were determined to be approximately 2 days, for mixtures of diastereomers dissolved in a minimum of DMF (14.6 M) at 25 °C; see the Supporting Information for details.
- [16] The *cis* and *trans* diastereomers of compound 12 could be separated by preparative HPLC and then used to prepare pure samples of *cis-7* and *trans-7*, see the Supporting Information for details.
- [17] The reaction was halted before completion to avoid the formation of decomposition products. Attempts to bring about a RCDA reaction using aminium radical catalysis (D. A. Bellville, D. D. Wirth, N. L. Bauld, *J. Am. Chem. Soc.* **1981**, *103*, 718– 720), or photoredox catalysis (Ref. [12b]), were met with failure.
- [18] An authentic sample of ramonanin B was not available.
- [19] Full details of these calculations are given in the Supporting Information.
- [20] The fast Diels-Alder dimerization of cyclopentadiene has been ascribed to its short C1-C4 distance, see: R. Sustmann, M. Böhm, J. Sauer, *Chem. Ber.* **1979**, *112*, 883-889.

www.angewandte.org

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

## Communications



thesis of the ramonanin family of natural

products has been achieved in short

order. These natural phenylpropanoid tetramers were assembled in seven steps

from the starting materials vanillin, ethylene, and the ethynyl Grignard reagent. Computational studies shed light on a surprisingly facile Diels–Alder dimerization.