

Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: B. Breit, Y. Zhou, A. -. Nikbakht and F. Bauer, *Chem. Sci.*, 2019, DOI: 10.1039/C9SC00980A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Rhodium catalyzed cycloisomerization and tandem Diels-Alder reaction for facile access to diverse bicyclic and tricyclic heterocycles

Yirong Zhou,^{a,b} Ali Nikbakht,^a Felix Bauer^a and Bernhard Breit^{*a}Received 00th January 20xx,
Accepted 00th January 20xx



DOI: 10.1039/x0xx00000x

A regioselective distal cycloisomerization of 1,6-allenenes was successfully developed to afford six-membered ring exocyclic 1,3-dienes employing a rhodium/diphosphine catalyst system. Deuterium labelling experiments and DFT calculations were performed to provide insights into the reaction mechanism of this unprecedented transformation. In addition, one-pot tandem Diels-Alder reactions with various dienophiles could readily construct diverse bicyclic and tricyclic nitrogen heterocycles, which are ubiquitous core scaffolds for a variety of natural products and bioactives. High efficiency as well as exclusive chemo- and regioselectivities for a broad substrate scope were achieved under mild conditions using a low catalyst loading of 0.5 mol%.

Transition metal catalyzed cycloisomerization reactions have proved to be synthetically useful and elegant methods to construct structurally diverse all carbon and heterocyclic frameworks in high efficiency as well as excellent atom and step economy.¹ In particular, cycloisomerization reactions of linear di-unsaturated systems with a suitable linker chain, such as 1,6-diynes,² 1,6-enynes,³ allenynes,⁴ bisallenenes,⁵ and allenedienes⁶ have been intensely investigated. Furthermore, readily available 1,6-allenenes have attracted extensive synthetic interest and their cycloisomerization has been successfully realized to access diverse heterocycles by employing various transition metal catalysts, including nickel,⁷ palladium,⁸ ruthenium,⁹ rhodium¹⁰ and gold,¹¹ etc. To the best of our knowledge, all the previously reported examples involved a proximal allene π -bond activation and produced five-membered ring dienes as major product (Scheme 1a). In view of diversity-oriented synthesis,¹² it is therefore in high demand to develop a new versatile catalytic system to furnish alternative cycloisomerization products which enable for subsequent multiple functionalizations.

Recently, our group has made significant progress on the development of rhodium catalyzed atom-efficient addition reactions of various pronucleophiles to unactivated allenes and alkynes to provide enantioenriched branched allylic products.¹³ To further expand the synthetic potential, we envisioned that 1,6-allenene would be an interesting substrates for

cycloisomerization to construct useful cyclic 1,3-dienes by using our rhodium/diphosphine catalytic system. Herein, we present an unprecedented rhodium catalyzed regioselective distal cycloisomerization of 1,6-allenenes to provide six-membered ring exocyclic 1,3-dienes exclusively (Scheme 1b). Based on deuterium labelling experiments and DFT calculations a plausible reaction mechanism could be elucidated. Moreover, a one-pot tandem Diels-Alder reaction with various dienophiles furnished diverse bicyclic and tricyclic nitrogen heterocycles in high atom- and step-economy.¹⁴ Such fused bicyclic and tricyclic nitrogen containing heterocycles constitute privileged core skeletons for a variety of natural products and drugs as well as fluorescent probes (Fig. 1), which highlights the manifold potential applications of this new methodology in medicinal and material chemistry.¹⁵

a) Previous work: proximal π -bond activationb) This work: distal π -bond activation

Scheme 1 Transition metal catalyzed cycloisomerizations of allenenes.

^a Institut für Organische Chemie und Biochemie
Albert-Ludwigs-Universität, Alberstr. 21, 79104 Freiburg (Germany)
E-mail: bernhard.breit@chemie.uni-freiburg.de

^b Key Laboratory of Functional Small Organic Molecules, Ministry of Education,
College of Chemistry and Chemical Engineering
Jiangxi Normal University, Nanchang, 330022, China

†Electronic Supplementary Information (ESI) available: CCDC 1579171 (2a) and CCDC 1834751 (4c). Experimental procedures and detailed characterization data of all new compounds. See DOI: 10.1039/x0xx00000x







Fig. 1 Selected examples for drugs and natural product containing bicyclic or tricyclic nitrogen heterocycles as core structures.

To test our hypothesis, unactivated terminal 1,6-allenene **1a** was chosen as a privileged model substrate for initial reactivity assays. Surprisingly, in contrast to literature reported results, no proximal coupled five-membered ring 1,3-diene product (as shown in Scheme 1a) was detected. Conversely, a completely new kind of six-membered ring exocyclic 1,3-diene was obtained as the sole product with exclusive regioselectivity and in good yield. The molecular structure of product **2a** was unambiguously confirmed by X-ray crystallographic analysis. This unexpected preliminary result induced us to systematically optimize the reaction conditions (Table 1). Firstly, several solvents were examined for the new cycloisomerization reaction (Table 1, entries 1-3) with DCE (1,2-dichloroethane) proving to be superior to THF (tetrahydrofuran) and toluene. Further investigations on the reaction temperature revealed that 60 °C was most suitable for this new transformation. The yield decreased with either higher or lower temperatures (Table 1, entry 5). Finally, the best result (80% yield) was obtained employing a lower substrate concentration of 0.1 M (Table 1, entry 6).

Table 1 Reaction conditions optimization for the cycloisomerization to access exocyclic 1,3-dienes^a




Entry	Solvent	Temp./°C	Yield ^b [%]
1	DCE	80	65
2	THF	80	58
3	Toluene	80	55
4	DCE	95	51
5	DCE	60	75
6 ^c	DCE	60	80

^aThe reactions were carried out on a 0.2 mmol scale of **1a** in the presence of 2.5 mol% of [Rh(COD)Cl]₂ and 5.0 mol% of DPEphos in DCE (1.0 mL) at Temp./°C for 20 h. ^bIsolated yield. ^c2.0 mL DCE was used (0.1 M).

With the optimized reaction conditions in hand, the substrate scope was explored. The results are shown in Table 2. First, substrates having different protecting groups at the nitrogen linker atom were investigated (**2a** to **2g**). Thus, in addition to sulfonyl groups, easily removable Boc and Cbz carbamates were well tolerated. Second, a variety of allenenes with an all-carbon linker were examined (**2h** to **2n**). In these cases, a slight increase of the reaction temperature to 80 °C was necessary in order to complete the cycloisomerization process. Different ester functions, such as methyl, ethyl and benzyl were all compatible in this reaction (**2h** to **2j**). Moreover, a group of masked hydroxyl functions were also suitable for the transformation (**2l** to **2n**). It is noteworthy that an interesting diene product **2k** with a spiro ketal structure was obtained in good yield.


Table 2 Substrate scope for the cycloisomerization to access exocyclic 1,3-dienes^a



^aThe reactions were carried out on a 0.2 mmol scale of **1** in the presence of 2.5 mol% of [Rh(COD)Cl]₂ and 5.0 mol% of DPEphos in DCE (2.0 mL) for 20 h. For products **2b** to **2g**, the reactions were performed at 60 °C. For products **2h** to **2n**, the reactions were performed at 80 °C. All the yields were isolated yields.


To gain deeper insights into the reaction mechanism of this new cycloisomerization reaction, a series of deuterium labelled substrates was prepared and subjected to standard reaction conditions (Scheme 2). The results with substrates **1a-1** and **1a-2** indicated that the deuterium atoms at terminal positions of alkene and allene completely remained in their original position. Hence, these carbon-hydrogen bonds should not change during the process (equation 1 and 2). However, the deuterium atom at the internal position of the alkene in **1a-3** completely shifted into the 5 position of the piperidine ring (**2a-3**), which indicated that an isomerization step might be involved to form the exocyclic carbon-carbon double bonds (equation 3).





Scheme 2 Deuterium labelling experiments.

To shed further light on the mechanism of this new cycloisomerization reaction, DFT computations were performed.¹⁶ Based on the labeling experiments and the DFT calculations the mechanism depicted in Scheme 3 is proposed. Starting from a monomeric [ClRh(DPEphos)] complex ([Rh]) the substrate **1a** could coordinate to form the chelate intermediate **I1**. An oxidative coupling via **TS1** furnishes the σ -allyl intermediate **I2**. This is followed by a β -hydride elimination (**TS2**) to give the σ -allyl hydride complex **I3**. Isomerization to the π -allyl intermediate **I4** (via **TS3**) and reductive elimination (via **TS4**) delivers product **2a** and regenerates the rhodium catalyst.




Scheme 3 Proposed mechanism for the rhodium-catalyzed cycloisomerization of 1,6-allenenes to six-membered exocyclic 1,3-dienes.

Fig. 2 displays the energy profile of the reaction. The highest energetic barriers are the oxidative coupling to form the five-membered metallacycle (**TS1**) with a ΔG of 14.6 kcal/mol (M06/def2SVP), and the reductive elimination step (**TS4**) with a ΔG of 14.5 kcal/mol (M06/def2SVP). The calculated reaction mechanism is in accord with the results of the deuterium labelling experiments. Furthermore, calculations of the

complete catalytic cycle for the traditional 5-membered ring cycloisomerization were performed. However, the energy barrier of the rate determining step was found to be significantly higher and is therefore unfavored.¹⁶

Considering that exocyclic 1,3-dienes **2** are potentially good reaction partners in Diels-Alder reactions, we anticipated that in the presence of suitable dienophiles, a one-pot tandem cycloisomerization/Diels-Alder reaction could be developed. This could become an efficient synthetic method to prepare diverse bicyclic and tricyclic nitrogen heterocycles.¹⁶ Indeed, as summarized in Table 3, we found that a wide range of 1,6-allenenes reacted smoothly in the presence of the rhodium catalyst and *N*-phenyl maleimide to furnish the desired tricyclic heterocycles **4** in good yields along with exclusive regio-, and diastereoselectivities. The constitution and relative configuration of **4c** were determined by X-ray crystallographic analysis, while the others were assigned by analogy. For the protected amide allenenes (**4a-4g**), as low as 0.5 mol% of rhodium catalyst was sufficient to achieve full conversion. Comparable yields showed that the protecting groups on the nitrogen atom exhibited negligible influences on the reaction. For carbon-linked allenene substrates (**4h-4o**) a slightly increased catalyst loading of 1 mol% was needed to allow for smooth and complete transformation. Various functional

Table 3 Substrate scope for the cycloisomerization and tandem Diels-Alder reaction^a

^aThe reactions were carried out on a 0.2 mmol scale of **1** with 1.0 equivalent of **3a** at 80 °C for 20 h. For products **4b** to **4g**, the reactions were performed in THF (1.0 mL) using 0.5 mol% of [Rh(COD)Cl]₂ and 1.0 mol% of DPEphos. For products **4h** to **4o**, the reactions were performed in DCE (1.0 mL) using 1.0 mol% of [Rh(COD)Cl]₂ and 2.0 mol% of DPEphos. All the yields were isolated yields.






Fig. 2 Computed catalytic cycle of the Rh catalyzed cycloisomerization of 1,6-allenes to six-membered exocyclic 1,3-dienes. All energies (PCM-M06/def2SVP//BP86/def2SVP) given with respect to energies of the $[CIRh(DPEphos)]$ complex and the substrate.

groups, such as ester, ketone, ketal and ethers were all well tolerated.


Next, the scope of dienophiles for the rhodium catalyzed domino cycloisomerization/Diels-Alder reaction was evaluated. As illustrated in Table 4, a wide range of symmetrical dienophiles proved suitable, providing the desired tricyclic and bicyclic heterocycles in good to high yields.

Table 4 Substrate scope for various dienophiles^a

^aThe reactions were carried out on a 0.2 mmol scale of **1a** with 1.0 equivalent of **5** in the presence of 0.5 mol% of $[Rh(COD)Cl]_2$ and 1.0 mol% of DPEphos in THF (1.0 mL) at 80 °C for 20 h. For products **6s** to **6u**, the reactions were performed in DCE (1.0 mL) using 2.0 mol% of $[Rh(COD)Cl]_2$ and 4.0 mol% of DPEphos and the dienophiles were added after the first cycloisomerization step completed. All the yields were isolated yields.

Thus, a series of *N*-aryl maleimides with either electron poor or electron rich aryl substituents behaved well and provided the tandem products in good to high yields (**6a-6n**). A variety of aryl halides including F, Cl, Br and even I were well tolerated enabling a subsequent derivatization through diverse cross coupling methods (**6e-6h**). Other well behaving dienophiles where benzoquinone (**6o**), the diphenyl ketone derived from fumaric acid (**6p**), dimethyl fumarate (**6q**), *trans*-dicyano ethylene (**6r**), acetylene dicarboxylate (**6s**), tetra-cyano ethylene (**6f**) and azo dicarboxylate (**6u**).

The obtained fused tricyclic heterocyclic products contain an internal tetra-substituted alkene function, which permits a variety of functionalization reactions. Towards this goal the tricyclic products **4a** and **4h** were selected for preliminary studies (Scheme 4).



Scheme 4 Diverse Synthetic transformations of **4a** and **4h**.

First, the palladium catalyzed diastereoselective hydrogenation delivered the saturated azatricyclic **7** as a single diastereomer in high yield. By treatment of suitable bromination reagent (PyH^+Br_3) in DCM, the dibrominated product **8** was obtained as a mixture of diastereomers. Epoxidation with *m*-CPBA furnished the epoxide **9** as a single diastereomer. A dihydroxylation yielded the vicinal diols **10a** and **10h** as single diastereomers from **4a** and **4h**, respectively.



These diols could be oxidatively cleaved to give the ten-membered diketones **11a** and **11h**. Alternatively, tricyclic product **4h** could be directly transformed into **11h** via ozonolysis. It is noteworthy, that such medium-sized fused bicyclic structures are difficult to access by other methods, highlighting the significance of this new method for potential applications in natural product synthesis and medicinal chemistry.

Conclusions

In conclusion, a novel regioselective cycloisomerization of 1,6-allenenes was successfully developed to generate six-membered ring exocyclic 1,3-dienes by using a rhodium/diphosphine catalyst system. Based on labelling experiments corroborated by DFT computations a plausible reaction mechanism could be suggested. Moreover, one-pot tandem Diels-Alder reactions with various dienophiles led to the efficient and rapid construction of diverse bicyclic and tricyclic nitrogen heterocycles. The new method displays high efficiency, broad substrate scope, complete atom and step economy, low catalyst loading of 0.5 mol%, excellent chemo-, regio-, and diastereoselectivity. Further expeditions on the application of this new method are currently underway in our laboratory.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the DFG. Y. Z. thanks the Sino-German (CSC-DAAD) Postdoc Scholarship and the National Natural Science Foundation of China (no. 21602089) for support. We thank Dr. D. Kratzert for the X-ray diffraction analysis; Dr. M. Keller for NMR analysis; Solvias, Umicore, BASF and Wacker for generous gifts of chemicals.

Notes and references

- Selected recent reviews: (a) Y. Lee, K. Kumar and H. Waldmann, *Angew. Chem. Int. Ed.*, 2018, **57**, 5212–5226; (b) D. Zhang, J. Liu, A. Córdova and W. Liao, *ACS Catal.*, 2017, **7**, 7051–7063; (c) C. I. Stathakis, P. L. Gkizis and A. L. Zografos, *Nat. Prod. Rep.*, 2016, **33**, 1093–1117; (d) D. Zhang, Z. Zhang and M. Shi, *Chem. Commun.*, 2012, **48**, 10271–10279; (e) A. Marinetti, H. Jullien and A. Voituriez, *Chem. Soc. Rev.*, 2012, **41**, 4884–4908; (f) I. D. G. Watson and F. Dean Toste, *Chem. Sci.*, 2012, **3**, 2899–2919; (g) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria, and A. Simonneau, *Chem. Rev.*, 2011, **111**, 1954–1993; (h) I. Nakamura and Y. Yamamoto, *Chem. Rev.*, 2004, **104**, 2127–2198.
- Selected recent reviews and examples: (a) S. Saito and Y. Yamamoto, *Chem. Rev.*, 2000, **100**, 2901–2915; (b) P. R. Chopade and J. Louie, *Adv. Synth. Catal.*, 2006, **348**, 2307–2327; (c) Y. Shibata and K. Tanaka, *Synthesis*, 2012, **44**, 323–350; (d) H. Jang and M. J. Krische, *J. Am. Chem. Soc.*, 2004, **126**, 7875–7880; (e) K. Tanaka, Y. Otake and M. Hirano, *Org. Lett.*, 2007, **9**, 3953–3956.
- Selected recent reviews: (a) Y. Hu, M. Bai, Y. Yang and Q. Zhou, *Org. Chem. Front.*, 2017, **4**, 2256–2275; (b) R. Dorel and A. M. Echavarren, *J. Org. Chem.*, 2015, **80**, 7321–7332; (c) S. I. Lee and N. Chatani, *Chem. Commun.*, 2009, 371–384; (d) V. Michelet, P. Y. Toullec and J. Gen  t, *Angew. Chem. Int. Ed.*, 2008, **47**, 4268–4315; (e) L. Zhang, J. Sun and S. A. Kozmin, *Adv. Synth. Catal.*, 2006, **348**, 2271–2296; (f) C. Bruneau, *Angew. Chem. Int. Ed.*, 2005, **44**, 2328–2334; (g) C. Aubert, O. Buisine and M. Malacria, *Chem. Rev.*, 2002, **102**, 813–834.
- (a) Y. Oonishi, Y. Kitano and Y. Sato, *Angew. Chem. Int. Ed.*, 2012, **51**, 7305–7308; (b) S. I. Lee, S. H. Sim, S. M. Kim, K. Kim and Y. K. Chung, *J. Org. Chem.*, 2006, **71**, 7120–7123; (c) K. M. Brummond, H. Chen, P. Sill and L. You, *J. Am. Chem. Soc.*, 2002, **124**, 15186–15187.
- (a) S. Kitagaki, M. Kajita, S. Narita and C. Mukai, *Org. Lett.*, 2012, **14**, 1366–1369; (b) S. M. Kim, J. H. Park, Y. K. Kang and Y. K. Chung, *Angew. Chem. Int. Ed.*, 2009, **48**, 4532–4535; (c) P. Lu and S. Ma, *Org. Lett.*, 2007, **9**, 2095–2097; (d) X. Jiang, X. Cheng and S. Ma, *Angew. Chem. Int. Ed.*, 2006, **45**, 8009–8013.
- (a) I. Alonso, H. Faustino, F. L  pez and J. L. Mascare  as, *Angew. Chem. Int. Ed.*, 2011, **50**, 11496–11500; (b) M. Gul  as, A. Collado, B. Trillo, F. L  pez, E. O  ate, M. A. Esteruelas and J. L. Mascare  as, *J. Am. Chem. Soc.*, 2011, **133**, 7660–7663; (c) P. Maule  n, R. M. Zeldin, A. Z. Gonz  lez and F. D. Toste, *J. Am. Chem. Soc.*, 2009, **131**, 6348–6349; (d) P. A. Wender, T. E. Jenkins and S. Suzuki, *J. Am. Chem. Soc.*, 1995, **117**, 1843–1844.
- (a) N. N. Noucti and E. J. Alexanian, *Angew. Chem. Int. Ed.*, 2015, **54**, 5447–5450; (b) B. M. Trost and J. M. Tour, *J. Am. Chem. Soc.*, 1988, **110**, 5231–5233.
- (a) A. K.   . Persson and J.-E. B  ckvall, *Angew. Chem., Int. Ed.*, 2010, **49**, 4624–4627; (b) B. Trillo, M. Gul  as, F. L  pez, L. Castedo and J. L. Mascare  as, *Adv. Synth. Catal.*, 2006, **348**, 2381–2384.
- (a) T. Nada, Y. Yoneshige, Y. Ii, T. Matsumoto, H. Fujioka, S. Shuto and M. Arisawa, *ACS Catal.*, 2016, **6**, 3168–3171; (b) S. Kang, B. Ko and D. Lee, *Tetrahedron Lett.*, 2002, **43**, 6693–6696.
- (a) K. Sugikubo, F. Omachi, Y. Miyanaga, F. Inagaki, C. Matsumoto and C. Mukai, *Angew. Chem. Int. Ed.*, 2013, **52**, 11369–11372; (b) K. M. Brummond, H. Chen, B. Mitasev and A. D. Casarez, *Org. Lett.*, 2004, **6**, 2161–2163; (c) T. Makino and K. Itoh, *J. Org. Chem.*, 2004, **69**, 395–405; (d) T. Makino and K. Itoh, *Tetrahedron Lett.*, 2003, **44**, 6335–6338; (e) P. A. Wender, F. Glorius, C. O. Husfeld, E. Langkopf and J. A. Love, *J. Am. Chem. Soc.*, 1999, **121**, 5348–5349.
- (a) P. Aillard, P. Retailleau, A. Voituriez and A. Marinetti, *Chem. Eur. J.*, 2015, **21**, 11989–11993; (b) H. Teller, M. Corbet, L. Mantilli, G. Gopakumar, R. Goddard, W. Thiel and A. F  rstner, *J. Am. Chem. Soc.*, 2012, **134**, 15331–15342; (c) A. Z. Gonz  lez, D. Benitez, E. Tkatchouk, W. A. Goddard and F. D. Toste, *J. Am. Chem. Soc.*, 2011, **133**, 5500–5507; (d) H. Teller, S. Fl  gge, R. Goddard and A. F  rstner, *Angew. Chem. Int. Ed.*, 2010, **49**, 1949–1953; (e) M. R. Luzung, P. Maule  n and F. D. Toste, *J. Am. Chem. Soc.*, 2007, **129**, 12402–12403; (f) M. A. Tarselli, A. R. Chianese, S. J. Lee and M. R. Gagn  , *Angew. Chem. Int. Ed.*, 2007, **46**, 6670–6673.
- Selected recent reviews: (a) S. L. Schreiber, *Science*, 2000, **287**, 1964–1969; (b) C. J. O' Connor, H. S. G. Beckmann and D. R. Spring, *Chem. Soc. Rev.*, 2012, **41**, 4444–4456; (c) J. Kim, H. Kim and S. B. Park, *J. Am. Chem. Soc.*, 2014, **136**, 14629–14638; (d) M. Garcia-Castro, S. Zimmermann, M. G. Sankar and K. Kumar, *Angew. Chem. Int. Ed.*, 2016, **55**, 7586–7605.
- (a) P. Koschker and B. Breit, *Acc. Chem. Res.*, 2016, **49**, 1524–1536; (b) A. M. Haydl, B. Breit, T. Liang and M. J. Krische, *Angew. Chem. Int. Ed.*, 2017, **56**, 11312–11325; (c) Y. Zhou and B. Breit, *Chem. Euro. J.*, 2017, **23**, 18156–18160; (d) Z. Liu and B. Breit, *Org. Lett.*, 2018, **20**, 300–303; (e) D. Berthold and B.



ARTICLE


Journal Name

View Article Online

DOI: 10.1039/C9SC00980A

- Breit, *Org. Lett.*, 2018, **20**, 598–601; (f) C. Grugel and B. Breit, *Org. Lett.*, 2018, **20**, 1066–1069; (g) P. Steib and B. Breit, *Angew. Chem. Int. Ed.*, 2018, **57**, 6572–6576.
- 14 Selected recent reviews: (a) J. Funel and S. Abele, *Angew. Chem. Int. Ed.*, 2013, **52**, 3822–3863; (b) S. Kotha, M. Meshram and A. Tiwari, *Chem. Soc. Rev.*, 2009, **38**, 2065–2092; (c) E. J. Corey, *Angew. Chem. Int. Ed.*, 2002, **41**, 1650–1667. During the preparation of this manuscript a related work using cycloisomerization of 1,6-enynes and tandem Diels-Alder reaction strategy has appeared: H. Zheng, Y. Wang, C. Xu, Q. Xiong, L. Lin and X. Feng, *Angew. Chem. Int. Ed.*, DOI: 10.1002/anie.201900421.
- 15 (a) P. A. Wender, M. S. Jeffreys and A. G. Raub, *J. Am. Chem. Soc.*, 2015, **137**, 9088–9093; (b) W. E. Childers, Jr. and R. B. Baudy, *J. Med. Chem.*, 2007, **50**, 2557–2562; (c) R. D. Clark, N. C. Ray, P. Blaney, P. H. Crackett, C. Hurley, K. Williams, H. J. Dyke, D. E. Clark, P. M. Lockey, R. Devos, M. Wong, A. White and J. K. Belanoff, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 5704–5708; (d) F. Chen and J. Huang, *Chem. Rev.*, 2005, **105**, 4671–4706.
- 16 For detailed information, see supporting information.





A new rhodium catalyzed regioselective cycloisomerization of 1,6-allenenes was successfully developed to prepare six-membered ring exocyclic 1,3-dienes. Moreover, one-pot tandem Diels-Alder reaction with various dienophiles proved to be a high efficient construction of diverse bicyclic and tricyclic nitrogen heterocycles. This methodology was distinguished by complete atom and step economy, low catalyst loading of 0.5 mol%, excellent chemo-, regio-, and diastereoselectivity.

