

Supplementary data for this paper are available from the IUCr electronic archives (Reference: LN1067). Services for accessing these data are described at the back of the journal.

References

- Bentele, H. J., Sünkel, K. & Beck, W. (1997). *Chem. Ber.* **130**, 1475–1477.
- Chiaroni, A., Riche, C., El Firdoussi, L., Benharref, A. & Karim, A. (1993). *Acta Cryst.* **C49**, 365–368.
- Cotton, F. A. (1990). *Chemical Applications of Group Theory*, 3rd ed., pp. 247–249. New York: Wiley Interscience.
- El Firdoussi, L., Allaoud, S., Karim, A., Barrero, A. F., Quirós, M., Castanet, Y. & Morreux, A. (1997). *Acta Cryst.* **C53**, 710–712.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Oike, H., Kai, Y., Miki, K., Tanaka, N. & Kasai, N. (1987). *Bull. Chem. Soc. Jpn.* **60**, 1993–1999.
- Sardone, N., Seresini, P. & Zecchi, G. (1996). *Acta Cryst.* **C52**, 613–615.
- Sheldrick, G. M. (1994). *SHELXTL/PC*. Version 5. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Solladie-Cavallo, A. (1989). *Advances in Metalorganic Chemistry*, Vol. 1, pp. 99–133. London: JAI Press.
- Stoe & Cie (1996a). *STADIA. Diffractometer Control Program*. Version 1.06. Stoe & Cie, Darmstadt, Germany.
- Stoe & Cie (1996b). *X-RED. Data Reduction Program*. Version 1.08. Stoe & Cie, Darmstadt, Germany.

Acta Cryst. (1999). **C55**, 363–365

(Z)- and (E)-1-methylthio-1-triphenylstannyl-2-phenylethene

CLEMENS BRUHN,^a DIRK STEINBORN,^a TOMÁŠ LÉBL,^b AND JAROSLAV HOLEČEK^b

^a*Institut für Anorganische Chemie, Martin-Luther-Universität Halle-Wittenberg, Kurt-Mothes-Straße 2, D-06120 Halle (Saale), Germany, and* ^b*Department of General and Inorganic Chemistry, University of Pardubice, Cs. legii 565, CZ-53210 Pardubice, Czech Republic. E-mail: bruhn@chemie.uni-halle.de*

(Received 17 September 1998; accepted 19 November 1998)


Abstract

The structures of the title compounds, (Z)-Ph₃Sn—C(SMe)=CHPh, (I), and (E)-Ph₃Sn—C(SMe)=CHPh, (II) {both [Sn(C₆H₅)₃(C₉H₉S)]}, (1-methylthio-2-phenylethenyl)triphenyltin}, were determined by single-crystal X-ray diffraction at room temperature. Both structures consist of discrete molecules in which the Sn atom is tetrahedrally coordinated by three phenyl groups and one 1-methylthio-2-phenylethenyl group. The planar ethenyl skeleton forms angles of 58.9(2) and 34.0(3)° with the 2-phenyl substituent in (I) and (II), respectively, excluding π interactions between them. In Z-isomer (I),

the methyl substituent on sulfur lies in the plane of the ethenyl group [torsion angle C—S—C=C 3.0(5)°], but this is not the case in E-isomer (II), which reveals a torsion angle C—S—C=C of 156.1(8)°.

Comment

The hydrometallation of non-symmetrically substituted alkynes can afford four possible isomeric ethenyl metal complexes. Especially in the case of alkynes $R'C\equiv CYR_n$ with Lewis basic heteroatom groups ($YR_n = NR_2, PR_2, OR, SR$; $R/R' = \text{alkyl, aryl, H}$), it might be difficult to predict the regio- and/or stereoselectivity. In the case of the hydrostannylation of PhC $\equiv$ CSPH by Ph₃SnH, both the non-catalysed and the Pd-catalysed reactions are quite selective and yield one major isomer at about 90–95% in each case. Because the assignment of the isomers by means of NMR spectroscopy (¹H, ¹³C and ¹¹⁹Sn) is not straightforward, the structures of both isomers were determined by single-crystal X-ray diffraction. The non-catalysed reaction gave (Z)-Ph₃Sn—C(SMe)=CHPh, (I), and the Pd-catalysed reaction (E)-Ph₃Sn—C(SMe)=CHPh, (II).



The molecular structures and numbering schemes are shown in Fig. 1. Both complexes consist of discrete molecules in which the Sn atoms are tetrahedrally coordinated by three phenyl ligands and one 1-methylthio-2-phenylethenyl ligand [C—Sn—C: 106.3(1)–112.7(1)° in (I) and 105.6(3)–110.0(3)° in (II)]. The three Sn—C_{Ph} bond lengths are similar [(I): 2.131(4)–2.143(4), mean value 2.135 Å; (II): 2.135(8)–2.162(8), mean value 2.146 Å]. In the case of isomer (I), the Sn—C_{ethenyl} bond seems to be slightly longer [(I): 2.173(4); (II): 2.148(7) Å]. There are no significant differences in the Sn—C bond lengths in Ph₃Sn—CH=CH₂ [Sn—C_{Ph} 2.11(3)–2.17(4) and Sn—C_{ethenyl} 2.07(3)–2.15(3) Å; Theobald & Trimaille, 1984] and (Z)-Ph₃Sn—C(CMe₂OH)=CHMe [Sn—C_{Ph} 2.118(3)–2.129(3) and Sn—C_{ethenyl} 2.125(3) Å; Willem *et al.*, 1994].

The ethenyl skeleton (C1–C3, Sn, S) is nearly planar [maximum deviation: 0.064(3) Å for C2 in (I), and 0.027(6) Å for C1 in (II)]. As in *trans*-[Pt{(E)-C(SMe)=CHPh}Cl(PPh₃)₂] (Steinborn *et al.*, 1998), in Z-isomer (I), the methyl substituent on sulfur (C9) also lies in this plane [deviation 0.011(8) Å; torsion angle C9—S—C1=C2 3.0(5)°]; this is not the case in E-isomer (II) [deviation 0.73(2) Å; torsion angle C9—S—C1=C2 156.1(8)°]. Without obvious reason, the angles at the ethenyl C1 and C2 atoms exhibit




Fig. 1. The molecular structures of (a) Z-isomer (I) and (b) E-isomer (II) showing 30% probability displacement ellipsoids. H atoms are shown as circles of an arbitrary radius.

distinct deviations from the ideal angle of 120° expected for sp^2 -hybridized C atoms [111.6(2)–126.6(4)° for (I) and 117.6(5)–130.0(7)° for (II)]. The interplanar angles of 58.9(2) and 34.0(3)° between the 2-phenyl substituent (C3–C8) and the ethenyl group in (I) and (II), respectively, exclude π interactions.

The closest $C_{Ph} \cdots H \cdots S$ distance is intermolecular in (I) [C26 $\cdots$ Sⁱ 3.594(7) and H26 $\cdots$ Sⁱ 2.79 Å; symmetry code: (i) 1 + x, y, z] and intramolecular in (II) [C4 $\cdots$ S 3.125(10) and H4 $\cdots$ S 2.63 Å].

Experimental

(Z)-Ph₃Sn–C(SMe)=CHPh, (I), was synthesized by the reaction of stoichiometric amounts of Ph₃SnH and PhC≡CSMe in *n*-hexane at 328 K under anaerobic conditions. After 5 h, the precipitate was filtered, washed with hexane and dried *in vacuo* (yield: 69%; m.p. 379 K). Crystals suitable for X-ray diffraction studies were obtained from chloroform/*n*-pentane solution. (E)-Ph₃Sn–C(SMe)=CHPh, (II), was obtained using an analogous reaction in THF at 243 K in the presence of 1 mol% [Pd(PPh₃)₄]. After 15 min, the solvent was removed *in vacuo* affording an oily residue (purity: 95%). Crystals (m.p. 341–343 K) suitable for X-ray diffraction studies separated after standing for several weeks at 288 K.

Compound (I) – Z isomer

Crystal data

[Sn(C₆H₅)₃(C₉H₉S)]

$M_r = 499.21$

Monoclinic

$P2_1/c$

$a = 10.190(9)$ Å

$b = 19.104(9)$ Å

$c = 12.808(9)$ Å

$\beta = 110.67(6)^\circ$

$V = 2333(3)$ Å³

$Z = 4$

$D_x = 1.421$ Mg m^{−3}

D_m not measured

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 32 reflections

$\theta = 5.23$ – 16.73°

$\mu = 1.195$ mm^{−1}

$T = 298(2)$ K

Block

$0.8 \times 0.6 \times 0.4$ mm

Colourless

Data collection

Stoe Stadi-4 four-circle diffractometer

$\omega/2\theta$ scans

Absorption correction:

semi-empirical via ψ scan (North *et al.*, 1968)

$T_{min} = 0.45$, $T_{max} = 0.62$

6742 measured reflections

4101 independent reflections

3750 reflections with

$I > 2\sigma(I)$

$R_{int} = 0.109$

$\theta_{max} = 25^\circ$

$h = -12 \rightarrow 12$

$k = 0 \rightarrow 23$

$l = -15 \rightarrow 15$

1 standard reflection

frequency: 60 min

intensity decay: none

Refinement

Refinement on F^2

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.098$

$S = 1.160$

4100 reflections

289 parameters

Only H-atom U 's refined

$w = 1/[\sigma^2(F_o^2) + (0.0414P)^2 + 0.9634P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} < 0.001$

$\Delta\rho_{max} = 1.144$ e Å^{−3}

$\Delta\rho_{min} = -1.402$ e Å^{−3}

Extinction correction: none

Scattering factors from

International Tables for Crystallography (Vol. C)

Table 1. Selected geometric parameters (Å, °) for (I)

C1–C2	1.331(5)	C10–Sn	2.143(4)
C1–S	1.740(4)	C16–Sn	2.131(4)
C1–Sn	2.173(4)	C22–Sn	2.133(4)
C9–S	1.789(5)		
C2–C1–S	123.7(3)	C16–Sn–C10	106.28(13)
C2–C1–Sn	124.6(3)	C22–Sn–C10	106.7(2)
S–C1–Sn	111.6(2)	C16–Sn–C1	111.34(13)

C1—C2—C3	126.6 (4)
C1—S—C9	105.1 (2)
C16—Sn—C22	111.57 (14)

C22—Sn—C1	112.74 (14)
C10—Sn—C1	107.9 (2)

Financial support from Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie is gratefully acknowledged.

Compound (II) – *E* isomer

Crystal data

[Sn(C₆H₅)₃(C₉H₉S)]

M_r = 499.21

Triclinic

P $\bar{1}$

a = 9.5512 (8) Å

b = 11.0433 (8) Å

c = 12.3793 (10) Å

α = 83.380 (8)°

β = 81.737 (8)°

γ = 67.018 (7)°

V = 1187.1 (2) Å³

Z = 2

D_x = 1.397 Mg m^{−3}

D_m not measured

Data collection

Stoe Stadi-4 four-circle diffractometer

$\omega/2\theta$ scans

Absorption correction:

semi-empirical via ψ scan (North *et al.*, 1968)

T_{min} = 0.62, *T_{max}* = 0.79

3732 measured reflections

3732 independent reflections

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 32 reflections

θ = 10.0–13.5°

μ = 1.174 mm^{−1}

T = 293 (2) K

Prism

0.4 × 0.3 × 0.2 mm

Colourless

3070 reflections with

I > 2σ(*I*)

$\theta_{\max}$ = 25°

h = −11 → 8

k = −12 → 13

l = 0 → 14

1 standard reflection

frequency: 60 min

intensity decay: none

Refinement

Refinement on *F*²

R [*F*² > 2σ(*F*²)] = 0.051

wR (*F*²) = 0.189

S = 1.239

3721 reflections

287 parameters

Only H-atom *U*'s refined

w = 1/[σ²(*F_o*) + (0.1046*P*)²]
where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} < 0.001

Δρ_{max} = 0.766 e Å^{−3}

Δρ_{min} = −0.841 e Å^{−3}

Extinction correction:

SHELXL93

Extinction coefficient:

0.030 (4)

Scattering factors from

International Tables for Crystallography (Vol. C)

Table 2. Selected geometric parameters (Å, °) for (II)

C1—C2	1.345 (11)	C10—Sn	2.162 (8)
C1—S	1.761 (7)	C16—Sn	2.135 (8)
C1—Sn	2.148 (7)	C22—Sn	2.141 (7)
C9—S	1.769 (10)		
C2—C1—S	120.7 (6)	C16—Sn—C1	108.1 (3)
C2—C1—Sn	117.6 (5)	C22—Sn—C1	110.6 (3)
S—C1—Sn	121.6 (4)	C16—Sn—C10	105.6 (3)
C1—C2—C3	130.0 (7)	C22—Sn—C10	110.9 (3)
C1—S—C9	105.1 (5)	C1—Sn—C10	111.6 (3)
C16—Sn—C22	110.0 (3)		

For both compounds, data collection: *STADI4* (Stoe & Cie, 1996a); cell refinement: *STADI4*; data reduction: *X-RED* (Stoe & Cie, 1996b); program(s) used to solve structures: *SHELXS86* (Sheldrick, 1990); program(s) used to refine structures: *SHELXL93* (Sheldrick, 1993); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996); software used to prepare material for publication: *SHELXL93*.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: NA1382). Services for accessing these data are described at the back of the journal.

References

- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Steinborn, D., Becke, S., Bruhn, C. & Heinemann, F. W. (1998). *J. Organomet. Chem.* **556**, 189–196.
- Stoe & Cie (1996a). *STADI4. Diffractometer Control Program*. Version 1.05c. Stoe & Cie, Darmstadt, Germany.
- Stoe & Cie (1996b). *X-RED. Data Reduction Program*. Version 1.05. Stoe & Cie, Darmstadt, Germany.
- Theobald, F. & Trimaille, B. (1984). *J. Organomet. Chem.* **267**, 143–149.
- Willem, R., Delmott, A., De Borger, I., Biesemans, M., Gielen, M., Kayser, F. & Tiekink, E. R. T. (1994). *J. Organomet. Chem.* **480**, 255–259.

Acta Cryst. (1999). **C55**, 365–368

A dimeric form of the nickel(II) thio-cyanate complex of bis(3-aminopropyl)-methylamine

ALOK K. MUKHERJEE,^a NIRMALYA P. NAYAK,^a ARUNENDU MONDAL^b AND NIRMALENDU RAY CHAUDHURI^b

^aDepartment of Physics, Jadavpur University, Calcutta 700 032, India, and ^bDepartment of Inorganic Chemistry, Indian Association for the Cultivation of Science, Calcutta 700 032, India. E-mail: akm@juphys.ernet.in

(Received 20 May 1998; accepted 2 November 1998)

Abstract

The structure of bis[*N,N*-bis(3-aminopropyl)methylamine]-1κ³*N*,2κ³*N*-di-μ-thiocyanato-1:2κ²*N*:S;1:2κ²*S*:*N*-dithiocyanato-1κ*N*,2κ*N*-dinickel(II), [Ni₂(NCS)₄(C₇H₁₉N₃)₂] or [Ni₂(μ-SCN)₂(medpt)₂(NCS)₂], where medpt is bis(3-aminopropyl)methylamine, consists of two crystallographically independent dimer types differing in their orientations. In each dimer, the two Ni^{II} ions are bridged by two SCN[−] ligands in an end-to-end fashion. The coordination polyhedra about the Ni^{II} atoms are distorted octahedra consisting of three N atoms of the medpt lig-