

Natural Product Synthesis

International Edition: DOI: 10.1002/anie.201604744
German Edition: DOI: 10.1002/ange.201604744


The Enantioselective Total Synthesis of Exochomine

Alison X. Gao[†], Tomoaki Hamada[†], and Scott A. Snyder^{*}

Dedicated to Professor K. C. Nicolaou on the occasion of his 70th birthday

Abstract: Molecules that possess fully substituted chiral centers are often challenging to construct, particularly if those centers connect two seemingly different halves or include a nitrogen atom. Herein, we describe an efficient approach to a molecule that combines both challenges in a single center in the form of exochomine. Failures in direct coupling led to a design fueled by highly specific reaction conditions for several steps and the development of an improved protocol for 1,4-reduction in a hindered context where numerous side reactions were possible. These chemoselective solutions should have value to other problems. Challenges in obtaining matching spectral data for the synthesized natural product are also discussed.

One of the key driving forces for new synthetic tactics and reactions are fully substituted chiral centers. Indeed, the efficient preparation of such centers often necessitates unique designs and novel reactivity considerations.^[1] Those challenges, however, are arguably magnified in those molecules where nature has placed such fully substituted centers at the connection point of two seemingly different halves. Compounds such as azadirachtin (**1**, Scheme 1),^[2] haplophytine (**2**),^[3] aplyviolene (**3**),^[4] and bionectin A (**4**)^[5] are representative cases, where years of effort and creative solutions often based on novel methods were required to generate those centers marked with a blue asterisk, noting that direct coupling of the two halves has typically proven impossible to achieve. Similarly high challenge also exists with α -tertiary amine centers,^[6] such as that found in acutumine (**5**).^[7] We have recently embarked on a program exploring molecules that combine both of these challenges within single structures,^[8] compounds that include the defensive alkaloids produced by ladybugs in the form of psyllorine A (**6**),^[9] the chilicorines (**7** and **8**),^[10] and exochomine (**9**).^[11,12] As one reflection of that challenge, direct couplings of seemingly appropriate monomers,^[8] such as the Prins-type addition^[13] shown between **10** and **11**, have failed in our hands to deliver



Scheme 1. The challenge of fully substituted chiral center formation, particularly in cases where two seemingly disparate halves are merged (**1–4**) and when those centers contain nitrogen (**5**), a problem combined in the coccinellid alkaloids (**6–9**).

the needed products, necessitating the development of alternate solutions. Herein, we document a route which can enantioselectively deliver exochomine (**9**) in 16 steps, highlighting a series of substrate specific steps and unique reaction conditions whose success in such a hindered context should be relevant for other synthesis problems.

[*] A. X. Gao,^[†] Dr. T. Hamada,^[†] Prof. Dr. S. A. Snyder
Dept. of Chemistry, The Scripps Research Institute
130 Scripps Way, Jupiter, FL 33458 (USA)

Prof. Dr. S. A. Snyder
Dept. of Chemistry, University of Chicago
5735 S. Ellis Avenue, Chicago, IL 60637 (USA)
E-mail: sasnyder@uchicago.edu
Homepage: <http://snyder-group.uchicago.edu/>

[†] These authors contributed equally to this work.


Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under <http://dx.doi.org/10.1002/anie.201604744>.

to most efforts towards similarly complex tricyclic amines.^[25] The relative stereochemistry of **23** was verified by X-ray diffraction of a benzoate derivative of its corresponding alcohol (see SI).

The synthesis of the pyrrole-based coupling partner commenced with heterocycle formation using inexpensive, commercially available L-glutamic acid ethyl ester **24** (Scheme 3) and 2,5-dimethoxytetrahydrofuran **25**. A subsequent HBr/AcOH-promoted Friedel–Crafts cyclization smoothly gave the desired acyl pyrrole,^[26] and was followed by dithiolane protection with 1,2-ethanedithiol as promoted by Zn(OTf)₂^[27] to give **26** in 37% yield overall for these 3 steps. Homologation to ester **27** was achieved in a further 3-step sequence involving DIBAL-H reduction of the ester to an aldehyde, Wittig olefination, and reduction using the copper hydride formed by premixing Cu(OAc)₂·H₂O, BDP, and poly(methylhydrosiloxane).^[28] The use of standard heterogeneous hydrogenation was precluded in the final step as it also impacted the dithiolane group, noting that this protecting group proved critical as other alternatives, such as ketals or protected alcohols, were too fragile. Of note, although the sequence proceeded through a potentially epimerizable aldehyde, the enantiopurity was largely retained, being determined to be 97% *ee* by chiral HPLC, and **27** could readily be prepared (> 4 g synthesized). Finally, **27** could also be converted into cyclized product **28** in 51% yield through exposure to BBr₃ in CH₂Cl₂ at 0°C for 20 min. This latter material had been previously prepared racemically by Meinwald in 5 steps,^[12] our overall route, though 2 steps longer, was asymmetric.


With azaphenylene aldehyde **23** in hand, as well as two potential coupling partners (**27** and **28**), studies were undertaken to bring the two halves together. Using these pieces and a range of other compounds (not shown), that process initially proved challenging, with approaches based on metathesis, Horner–Wadsworth–Emmons olefinations,^[29] and Nozaki–Hiyama–Kishi couplings^[30] all failing.

Our first success was registered with a titanium-mediated aldol condensation^[31] between ketone **28** and aldehyde **23** which delivered **30**, albeit in low and variable yields due principally to the instability of ketone **28** under the reaction conditions. Pleasingly, after much optimization, we found that a milder coupling could be achieved by using the more conformationally flexible ester **27**. In the event, LDA-mediated enolization and aldol addition with **23** (0.9 equiv) in THF at –78°C for 2 h provided **29** (any warming led to retro-aldol reactions); this intermediate was then treated with *p*-TsOH in situ, with subsequent solvent evaporation, resuspension in benzene, and heating at 60°C for 4 h effecting dehydration and Friedel–Crafts cyclization in one pot, delivering **30** in 32% yield. No intermediates were characterizable, so the exact order of steps is unknown, and though this outcome is not numerically high, 3 steps were achieved overall, with the likely culprit for throughput being the Friedel–Crafts reaction given the observed yield (51%) in the conversion of **27** into **28**. Also worth noting is that the same conditions (as well as other protocols) did not lead to coupling products when **28** was the enolate partner, likely because



Scheme 3. Completion of the total synthesis of exochomine: a) **25** (1.05 equiv), ClCH₂CH₂Cl, 65 °C, 4 h, 80%; b) HBr (2 equiv), AcOH, 25 °C, 3 h, 60%; c) 1,2-ethanedithiol (1.3 equiv), Zn(OTf)₂ (1.3 equiv), CH₂Cl₂, 25 °C, 12 h, 78%; d) DIBAL-H (1.05 equiv), CH₂Cl₂, –78 °C, 30 min; e) PPh₃CHCO₂Me (1.1 equiv), CH₂Cl₂, 0 → 25 °C, 12 h, 81% over 2 steps; f) Cu(OAc)₂·H₂O (5 mol%), BDP (0.5 mol%), PMHS (2.0 equiv), *t*-BuOH (2.0 equiv), toluene, 25 °C, 8 h, 90%; g) BBr₃ (2.5 equiv), CH₂Cl₂, 0 °C, 20 min, 51%; h) LDA (1.5 equiv), **23** (0.9 equiv), THF, –78 °C, 2 h; *p*TsOH·H₂O (5.0 equiv), –78 → 25 °C; evaporate; benzene, 60 °C, 4 h, 32%; i) TiCl₄ (2.0 equiv), Et₃N (2.1 equiv), CH₂Cl₂, –78 °C, 2 h, 19%; j) Mn(dpm)₃ (2.2 equiv), MeSi(OEt)₂H (5.0 equiv), *i*-PrOH/CH₂Cl₂ (8:1), 25 °C, 16 h, 48%, 10:1 d.r.; k) MeLi (2.0 equiv), THF, –78 → –30 °C, 30 min, *p*TsOH·H₂O (5 equiv), –30 → 25 °C, 4% **9** and 65% protected **9**; l) PhI(OAc)₂ (1.3 equiv), CH₃CN/CH₂Cl₂/H₂O (5:1:1), –10 °C, 10 min, 31%, 23% from **31**; p) HCl (1.0 M in Et₂O, 1.5 equiv), EtOAc, 25 °C, 30 min, 70%. Ts = tosyl; BDP = 1,2-bis(diphenylphosphino)benzene; PMHS = poly(methylhydrosiloxane).

retro-aldol pathways might be favored due to its more rigid framework.

Table 1: Key optimization studies for the selective 1,4-reduction of enone **30**.

Entry	Conditions	Product	Yield of 31 [%] ^[a]
1	Cu(OAc) ₂ ·H ₂ O, BDP, PMHS	n.r.	0
2	[CuH(PPh ₃) ₆]	n.r.	0
3	Cu(OAc) ₂ ·H ₂ O, P(O <i>i</i> -Pr) ₃ , MeSi(OEt) ₂ H	31 + 1,6-reduction	< 20 ^[b]
4	SmI ₂ , <i>t</i> -BuOH, THF	33 + 31 + 1,6-reduction	< 20 ^[b]
5	Pd(PPh ₃) ₄ , ZnCl ₂ , <i>n</i> -Bu ₃ SnH	n.r.	0
6	catecholborane	33	0
7	DIBAL-H	33	0
8	RedAl, CuBr, BF ₃ ·OEt ₂	n.r.	0
9	CuBr·Me ₂ S, <i>t</i> -BuLi, DIBAL-H, THF/HMPA	n.r.	0
10	NBSH, Et ₃ N	n.r.	0
11	TsNHNH ₂ , NaOAc, THF/H ₂ O	n.r.	0
12	Mn(dpm) ₃ , PhSiH ₃ , TBHP	31 + deprotection	25
13	Mn(dpm) ₃ , PhSiH ₃	31	40
14	Mn(dpm) ₃ , MeSi(OEt) ₂ H	31	52

[a] A very small amount of isomer at the newly formed chiral center was also observed in these reductions, but was separated and is not part of these yield calculations; [b] Isolated yield, but includes some minor inseparable contaminants.

From here, only a few operations remained to complete the target, with the next, selective conjugate enone reduction of **30** to give **31**, proving highly challenging to effect as the dithiolane group, benzylic ketone, and acyl pyrrole all proved prone to reduction (leading to products such as **33**). As shown in Table 1, copper hydride-based approaches gave sluggish reactions and, when products were observed, typically the major adduct was 1,6-reduction across the pyrrole ring (entries 1–3). Radical-based methods such as SmI₂ also afforded predominantly 1,6-reduction (entry 4), while a variety of other standard protocols for conjugate enone reduction did not provide promising results (entries 5–11). Only Mn^{III}-mediated conditions as reported by Shenvi^[32] (entry 12) and Magnus^[33] (entry 13) provided encouraging selectivity and yield, with the use of the same hydride source from entry 3, that is, MeSi(OEt)₂H, providing the best overall outcome for this transformation in 52% isolated yield (entry 14). The identity and absolute configuration of **31** was confirmed by X-ray analysis.

Finally, a one-pot methyl addition, elimination, and ketone deprotection from **31** afforded a small amount of exochomine (**9**) alongside its protected form in 4% and 65% yield, respectively. The use of PhI(OAc)₂ in a 5:1:1 mixture of CH₃CN/CH₂Cl₂/H₂O converted any remaining protected intermediate into the final compound (i.e. **9**).^[34] As NMR data for the free base form of the target was not available, synthetic exochomine was converted into its corresponding HCl salt (i.e. **32**) and purified by preparative TLC. However, upon comparison of both its ¹H and ¹³C NMR data to that reported by the isolation team, including experiments which

sought to alter the acid content by titration with HCl as well as eliminate water content in the sample,^[35] there were always subtle discrepancies from the isolation data without specific patterning. Pleasingly, however, X-ray diffraction of a suitably grown crystal of synthetic **32** obtained from slow evaporation in CH₂Cl₂/EtOAc was able to confirm its structure. We are not completely sure of the source of the variances in the respective NMR spectra between the synthetic and natural material, but believe that water content is the likely culprit and should be considered as an additional point of concern for such compounds;^[36] counterion impurities could also be at play as the exact procedure by which the isolation team obtained their HCl salt from the crude extract is not defined.^[37]

In conclusion, we have completed the first enantioselective total synthesis of the coccinellid alkaloid exochomine (**9**) in 16 linear steps. The key α -tertiary

amine center linking the two disparate halves was constructed using a highly diastereoselective Strecker reaction from an iminium precursor, with the key linking C–C bond formed convergently using an aldol reaction from two advanced intermediates in a cascade where a Friedel–Crafts cyclization proceeded in the same pot. Flexibility in coupling partners was key to its success. In addition, selective 1,4-reduction conditions based on the combination of Mn^{III} and a unique hydride source succeeded on a reactive and sterically encumbered system where other tools failed, and should be of use for other challenging systems. Current efforts seek to deploy the strategies and methodological advances from this synthesis to other challenging targets.

Acknowledgements

We thank Dr. HaJeung Park (Scripps) for a preliminary X-ray structure of a derivative of **23** and Dr. Alexander Filatov (Chicago) for X-ray structures of **31** and **32**. The latter was obtained at Argonne National Labs; ChemMatCARS Sector 15 is principally supported by the Divisions of Chemistry (CHE) and Materials Research (DMR), National Science Foundation, under grant number NSF/CHE-1346572. Use of the Advanced Photon Source, an Office of Science User Facility operated for the U.S. Department of Energy (DOE) Office of Science by Argonne National Laboratory, was supported by the U.S. DOE under Contract No. DE-AC02-06CH11357. We also thank Dr. Xiangming Kong (Scripps) for NMR assistance and Dr. Jun-Yong Choi (Scripps) for help

with calculations. Financial support came from TSRI, the University of Chicago, and Daiichi-Sankyo (to T.H.).

Keywords: 1,4-reduction · alkaloid · azaphenylene · coccinellid · total synthesis

- [1] For general reviews on forming all carbon quaternary centers, see: a) K. Fuji, *Chem. Rev.* **1993**, *93*, 2037; b) K. W. Quasdorf, L. E. Overman, *Nature* **2014**, *516*, 181.
- [2] G. E. Veitch, E. Beckmann, B. J. Burke, A. Boyer, S. L. Maslen, S. V. Ley, *Angew. Chem. Int. Ed.* **2007**, *46*, 7629; *Angew. Chem.* **2007**, *119*, 7773.
- [3] a) K. C. Nicolaou, S. M. Dalby, U. Majumder, *J. Am. Chem. Soc.* **2008**, *130*, 14942; b) K. C. Nicolaou, S. M. Dalby, S. Li, T. Suzuki, D. Y.-K. Chen, *Angew. Chem. Int. Ed.* **2009**, *48*, 7616; *Angew. Chem.* **2009**, *121*, 7752; c) H. Ueda, H. Satoh, K. Matsumoto, K. Sugimoto, T. Fukuyama, H. Tokuyama, *Angew. Chem. Int. Ed.* **2009**, *48*, 7600; *Angew. Chem.* **2009**, *121*, 7736.
- [4] a) M. J. Schnermann, L. E. Overman, *J. Am. Chem. Soc.* **2011**, *133*, 16425; b) M. J. Schnermann, L. E. Overman, *Angew. Chem. Int. Ed.* **2012**, *51*, 9576; *Angew. Chem.* **2012**, *124*, 9714.
- [5] A. Coste, J. Kim, T. C. Adams, M. Movassaghi, *Chem. Sci.* **2013**, *4*, 3191.
- [6] a) F. Li, S. S. Tartakoff, S. L. Castle, *J. Am. Chem. Soc.* **2009**, *131*, 6674; b) S. M. King, N. A. Calandra, S. B. Herzon, *Angew. Chem. Int. Ed.* **2013**, *52*, 3642; *Angew. Chem.* **2013**, *125*, 3730.
- [7] For general reviews on forming α -tertiary amines, see: a) J. Clayden, M. Donnard, J. Lefranc, D. J. Tetlow, *Chem. Commun.* **2011**, *47*, 4624; b) A. Hager, N. Vrieling, D. Hager, J. Lefranc, D. Trauner, *Nat. Prod. Rep.* **2016**, *33*, 491; For a few recent cases, see: c) T. Nishikawa, D. Urabe, M. Isobe, *Angew. Chem. Int. Ed.* **2004**, *43*, 4782; *Angew. Chem.* **2004**, *116*, 4886; d) B. Trost, N. Cramer, H. Bernsmann, *J. Am. Chem. Soc.* **2007**, *129*, 3086; e) T. J. Greshock, A. W. Grubbs, R. M. Williams, *Tetrahedron* **2007**, *63*, 6124; f) A. Hinman, J. Du Bois, *J. Am. Chem. Soc.* **2003**, *125*, 11510.
- [8] T. C. Sherwood, A. H. Trotta, S. A. Snyder, *J. Am. Chem. Soc.* **2014**, *136*, 9743.
- [9] a) F. C. Schröder, T. Tolasch, *Tetrahedron* **1998**, *54*, 12243; b) B. Lebrun, J.-C. Braekman, D. Daloz, P. Kalushkov, J. M. Pasteels, *Tetrahedron Lett.* **1999**, *40*, 8115.
- [10] a) K. D. McCormick, A. B. Attygalle, S.-C. Xu, A. Svatoš, J. Meinwald, M. A. Houck, C. L. Blankespoor, T. Eisner, *Tetrahedron* **1994**, *50*, 2365; b) X. Shi, A. B. Attygalle, J. Meinwald, *Tetrahedron* **1995**, *51*, 8711; c) Q. Hunag, A. B. Attygalle, J. Meinwald, M. A. Houck, T. Eisner, *J. Nat. Prod.* **1998**, *61*, 508.
- [11] M. Timmermans, J.-C. Braekman, D. Daloz, J. M. Pasteels, J. Merline, J.-P. Declercq, *Tetrahedron Lett.* **1992**, *33*, 1281.
- [12] Only one report outside our efforts (Ref. [8]) exists for these targets: J. C. Shattuck, J. Meinwald, *Tetrahedron Lett.* **1997**, *38*, 347.
- [13] Intermolecular aza-Prins reaction are very rare: a) H. Kinoshita, O. J. Ingham, W. W. Ong, A. B. Beeler, J. A. Porco, *J. Am. Chem. Soc.* **2010**, *132*, 6412; For selected, recent intramolecular processes, see: b) H. M. Lee, C. Hieto-Oberhuber, M. D. Shair, *J. Am. Chem. Soc.* **2008**, *130*, 16864; c) X. Liu, M. P. McCormack, S. P. Waters, *Org. Lett.* **2012**, *14*, 5574; d) T. Nishimura, A. K. Unni, S. Yokoshima, T. Fukuyama, *J. Am. Chem. Soc.* **2013**, *135*, 3243.
- [14] a) M. G. Pizzuti, A. J. Minnaard, B. L. Feringa, *Org. Biomol. Chem.* **2008**, *6*, 3464; b) C. A. Guerrero, E. J. Sorensen, *Org. Lett.* **2011**, *13*, 5164; c) D. Stead, G. Carbone, P. O'Brien, K. R. Campos, I. Coldham, A. Sanderson, *J. Am. Chem. Soc.* **2010**, *132*, 7260; For the first case of α -alkylation using Boc as a directing group (without Cu), see: d) P. Beak, W.-K. Lee, *Tetrahedron Lett.* **1989**, *30*, 1197; e) P. Beak, W. K. Lee, *J. Org. Chem.* **1993**, *58*, 1109.
- [15] a) D. M. Brestensky, D. E. Huesland, C. McGettigan, J. M. Stryker, *Tetrahedron Lett.* **1988**, *29*, 3749; b) W. S. Mahoney, D. M. Brestensky, J. M. Stryker, *J. Am. Chem. Soc.* **1988**, *110*, 291; c) J. W. Yang, M. T. H. Fonseca, B. List, *Angew. Chem. Int. Ed.* **2004**, *43*, 6660; *Angew. Chem.* **2004**, *116*, 6829; For an NADH model, see: d) N. Baba, T. Makino, J. Oda, Y. Inouye, *Can. J. Chem.* **1980**, *58*, 387.
- [16] Inspired by the addition of arylstannanes: D. A. Black, B. A. Arndtsen, *J. Org. Chem.* **2005**, *70*, 5133.
- [17] a) M. Kira, T. Hino, H. Sakurai, *Chem. Lett.* **1991**, *277*; b) Y. Nishikawa, M. Kitajima, H. Takayama, *Org. Lett.* **2008**, *10*, 1987.
- [18] Y. Arata, M. Hanaoka, S. K. Kim, *Chem. Pharm. Bull.* **1975**, *23*, 1142.
- [19] N. R. Candeias, F. Montalbano, P. M. S. D. Cal, P. M. P. Gois, *Chem. Rev.* **2010**, *110*, 6169.
- [20] a) J. W. Yang, C. Chandler, M. Stadler, D. Kampen, B. List, *Nature* **2008**, *452*, 453; b) S. Kobayashi, H. Kiyohara, M. Yamaguchi, *J. Am. Chem. Soc.* **2011**, *133*, 708; c) R. A. Pilli, D. Russowsky, *J. Chem. Soc. Chem. Commun.* **1987**, 1053; d) S. Kobayashi, S. Nagayama, *J. Am. Chem. Soc.* **1997**, *119*, 10049.
- [21] a) J. C. Lo, J. Gui, Y. Yabe, C.-M. Pan, P. S. Baran, *Nature* **2014**, *516*, 343; b) J. Gui, C.-M. Pan, Y. Jin, T. Qin, J. C. Lo, B. J. Lee, S. H. Spergel, M. E. Mertzman, W. J. Pitts, T. E. La Cruz, M. A. Schmidt, N. Darvatkar, S. R. Natarajan, P. S. Baran, *Science* **2015**, *348*, 886.
- [22] R. H. Mueller, M. E. Thompson, R. M. DiPardo, *J. Org. Chem.* **1984**, *49*, 2217.
- [23] For reviews, see: a) R. V. Stevens, *Acc. Chem. Res.* **1984**, *17*, 289; b) H.-P. Husson, J. Royer, *Chem. Soc. Rev.* **1999**, *28*, 383; We note that our own efforts to observe any nitrile equilibration in deuterated NMR solvents were inconclusive; kinetic control is likely, but this example is somewhat different from those in the reviews listed above.
- [24] A. D. Bochevarov, E. Harder, T. F. Hughes, J. R. Greenwood, D. A. Braden, D. M. Philipp, D. Rinaldo, M. D. Halls, J. Zhang, R. A. Friesner, *Int. J. Quantum Chem.* **2013**, *113*, 2110.
- [25] N. N. Rao, J. K. Cha, *J. Am. Chem. Soc.* **2015**, *137*, 2243.
- [26] C. W. Jefford, K. Sienkiewicz, S. R. Thornton, *Helv. Chim. Acta* **1995**, *78*, 1151.
- [27] a) M. E. Kuehne, W. G. Bornmann, W. G. Earley, I. Marko, *J. Org. Chem.* **1986**, *51*, 2913; b) B. M. Trost, J. R. Parquette, *J. Org. Chem.* **1994**, *59*, 7568.
- [28] B. A. Baker, Z. V. Boskovic, B. H. Lipshutz, *Org. Lett.* **2008**, *10*, 289.
- [29] a) J. A. Bisceglia, L. R. Orelli, *Curr. Org. Chem.* **2012**, *16*, 2206; b) M. A. Blanchette, W. Choy, J. T. Davis, A. P. Essensfeld, S. Masamune, W. R. Roush, T. Sakai, *Tetrahedron Lett.* **1984**, *25*, 2183; c) T. Calogeropoulos, G. B. Hammond, D. F. Wiemer, *J. Org. Chem.* **1987**, *52*, 4185.
- [30] a) K. Takai, M. Tagashira, T. Kuroda, K. Oshima, K. Utimoto, H. Nozaki, *J. Am. Chem. Soc.* **1986**, *108*, 6048; b) H. Jin, J. Uenishi, W. J. Christ, Y. Kishi, *J. Am. Chem. Soc.* **1986**, *108*, 5644; c) Y. Kishi, *Pure Appl. Chem.* **1992**, *64*, 343.
- [31] a) C. R. Harrison, *Tetrahedron Lett.* **1987**, *28*, 4135; b) R. Annunziata, M. Cinquini, F. Cozzi, P. G. Cozzi, E. Consolandi, *Tetrahedron* **1991**, *47*, 7897.
- [32] K. Iwasaki, K. K. Wan, A. Oppedisano, S. W. M. Crossley, R. A. Shenvi, *J. Am. Chem. Soc.* **2014**, *136*, 1300.
- [33] P. Magnus, M. J. Waring, D. A. Scott, *Tetrahedron Lett.* **2000**, *41*, 9731.
- [34] CH_2Cl_2 was necessary to dissolve the substrate. Other deprotections, such as metal-mediated protocols (Hg, Tl, Ag),

alkylating agents (MeI, EtI) and oxidizing agents [IBX, PhI-(CO₂CF₃)₂], failed.

- [35] We thank Prof. J.-C. Braekman and Prof. D. Dalozé for providing copies of their original spectra of **32**; these data included sets not originally referenced that show variation, potentially due to water content, as one is noted to have water present and the other does not.
- [36] For recent alkaloid syntheses that point to similar challenges, such as needing to titrate acid to obtain matching spectra, see: a) R. A. Altman, B. L. Nilsson, L. E. Overman, J. R. de Alaniz, J. M. Rohde, V. Taupin, *J. Org. Chem.* **2010**, *75*, 7519; b) A. S. Lee, B. B. Liau, M. D. Shair, *J. Am. Chem. Soc.* **2014**, *136*, 13442.
- [37] For cases where an acid adduct was needed for matching spectra, see: a) X.-M. Zhang, Y.-Q. Tu, F.-M. Zhang, H. Shao, X. Ming, *Angew. Chem. Int. Ed.* **2011**, *50*, 3916; *Angew. Chem.* **2011**, *123*, 4002; b) H. Li, X. Wang, B. Hong, X. Lei, *J. Org. Chem.* **2013**, *78*, 800.

Received: May 15, 2016

Published online: ■ ■ ■ ■, ■ ■ ■ ■

Communications




Natural Product Synthesis

A. X. Gao, T. Hamada,
S. A. Snyder*



The Enantioselective Total Synthesis of
Exochomine



Challenging chirality: Use of a distinct synthetic strategy to address the challenges in merging two different halves bearing a chiral center at their merger point has enabled access to the alkaloid exochomine in 16 steps. Key operations include the functionalization of a hindered iminium precursor, a unique aldol-based coupling cascade, and a robust protocol for 1,4-reduction in a hindered context where numerous side-reactions can occur.