



## Communication

Subscriber access provided by University of Newcastle, Australia

# **Boronate Ester Bullvalenes**

Harshal D Patel, Thanh-Huyen Tran, Christopher J. Sumby, Lukáš F. Pašteka, and Thomas Fallon J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.9b12930 • Publication Date (Web): 10 Feb 2020 Downloaded from pubs.acs.org on February 11, 2020

## **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# **Boronate Ester Bullvalenes**

Harshal D. Patel,<sup>a</sup> Thanh-Huyen Tran,<sup>a</sup> Christopher J. Sumby,<sup>a</sup> Lukáš F. Pašteka,<sup>b</sup> and Thomas Fallon<sup>a\*</sup>

[a] Department of Chemistry, The University of Adelaide, Adelaide, SA 5005, Australia

[b] Department of Physical and Theoretical Chemistry and Laboratory for Advanced Materials, Faculty of Natural Sciences, Comenius University, Ilkovičova 6, Bratislava, Slovakia.

\*Corresponding author: thomas.fallon@adelaide.edu.au

Supporting Information Placeholder

**ABSTRACT:** Boronate ester bullvalenes are now accessible in two to four operationally simple steps. This unlocks late-stage diversification through Suzuki cross-coupling reactions to give mono-, di-, and tri-substituted bullvalenes. Moreover, a linchpin strategy enables preprogrammed installation of two different substituents. Analysis of solution phase isomer distributions and single crystal X-ray structures reveals that isomer preference in the crystal lattice is due to general shape selectivity.

In the 1960s the fluxional caged hydrocarbons were discovered and intently studied.<sup>1</sup> The canonical examples are bullvalene,<sup>2</sup> semibullvalene,<sup>3</sup> and the barbaryl cation.<sup>4</sup> Bullvalene is extraordinary as the only stable organic structure with the property of *total degeneracy*. Each of the ten positions in the molecule exchange with every other, and there are no permanent carbon-carbon bonds. For substituted bullvalenes, substituents will spontaneously explore all possible relative arrangements and exist as a dynamic ensemble of isomers (Figure 1a).

For such an intriguing molecular scaffold, the synthetic methods available to prepare and modify derivatives are surprisingly limited.<sup>5</sup> Schröder first prepared the parent hydrocarbon in 1963 through a serendipitous two-step sequence from cyclooctatetraene (COT).<sup>6</sup> While this method was scalable, it had no versatility and further substitution was only achieved through low-yielding bromination/elimination sequences (Figure 1b).<sup>7</sup>

Recently, Echavarren reported the preparation of mono- and di-substituted bullvalenes using a modular gold-catalysed oxidative cyclisation as the key step (Figure 1c).<sup>8</sup> In a series of reports, Bode demonstrated the synthesis of highly complex tetra-substituted bullvalenes. (Figure 1d).<sup>9</sup> However, this approach required relatively long synthetic sequences.

#### Figure 1a. Fluxional isomerism of bullvalene



1b. Schröder's synthesis of substituted bullvalenes



1c. Echavarren's gold-catalysed synthesis of bullvalenes



1d. Bode's approach to tetrasubstituted bullvalenes







This revival of interest in shape-shifting molecules is accompanied by a recognition of their potential in the context of dynamic covalent chemistry and molecular devices.<sup>10</sup> The foremost demonstration of these concepts is Bode's use of a tetrasubstituted bullvalene as a dynamic molecular sensor array for polyols.<sup>11</sup> However, further advances in this field demand more efficient synthetic methods.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

We recently reported a two-step approach to mono- and di-substituted bullvalenes through [6+2] cycloaddition of COT to alkynes followed by di- $\pi$ -methane rearrangement (Figure 1e).<sup>12</sup> While this represents the shortest synthetic strategy to date, the substrate scope with respect to both reaction steps was admittedly limited. We recognised the overarching need for synthetic building blocks and methodology that would allow for the modular late-stage incorporation of desired substituents through versatile and reliable chemistries. Boronate ester functionalised bullvalenes became irresistible targets.

In this communication we report the efficient synthesis of a family of mono- di- and tri-substituted boronate ester bullvalenes **1-4** (Figure 1f) and demonstrate their general utility in Suzuki cross-coupling reactions. The disubstituted bullvalene **3** incorporates a pinacol boronate ester (Bpin) as well as an *N*-methyliminodiacetic acid boronate ester ( $B_{MIDA}$ ) group. This linchpin enables selective double Suzuki cross-coupling reactions. The unparalleled versatility of this toolbox opens the way to explore shape-shifting chemical behaviour in a wide range of settings.

Suzuki cross-couplings gave access to a series of aryl substituted bullvalenes, many of which are highly crystalline. This allows for a comparative analysis of isomer preference in solution and the solid state. Solution phase population distributions were characterised through low-temperature NMR experiments and compared with the results of density functional theory (DFT) calculations. The isomer population distributions follow generally predictable lines. Single crystal X-ray analyses reveal several examples whereby a minor solution phase isomer is preferred in the crystal through what appears to "shape-selective" crystallisation phenomena be whereby resolution of a dynamic ensemble of isomers to a single isomer crystal is governed predominantly by dispersion forces and effective packing, rather than specific non-covalent interactions.

The key to our synthetic strategy is the cobalt-catalysed [6+2] cycloaddition of alkynes with COT.<sup>13</sup> This catalytic system has emerged in the past two decades as a powerful and versatile tool.<sup>14</sup> Among the pioneering contributions of Hilt are several examples of [4+2] cycloaddition reactions involving alkynyl boronate esters.<sup>15</sup> This encouraging precedent suggested that mono-Bpinacetylene 6 and bis-Bpin-acetylene 7<sup>16</sup> might be good reaction partners in [6+2] cycloadditions with cyclooctatetraene. Happily, this turned out to be true providing access to the corresponding bicyclo[4.2.2]deca-2,4,7,9-tetraenes (BDT) 9 and 10 in good yield and on gram scale (Scheme 1). COT-Bpin 8, prepared in two steps, undergoes [6+2] cycloaddition with 7 to give reliable access to the tri-Bpin-BDT 12, albeit in low yield.<sup>17</sup> Bis-Bpin-BDT **10** was modified by

condensation with *N*-methyliminodiacetic acid to give the corresponding ester **11**.<sup>18</sup>

With this range of boronate ester BDTs in hand we were delighted to find that the photochemical di- $\pi$ -methane rearrangement proceeded smoothly to give the corresponding bullvalenes in good to excellent yields. The use of inexpensive and operationally simple 365 nm LED lamps together with 9*H*-thioxanthen-9-one as a triplet sensitizer renders this transformation particularly rapid and convenient.<sup>19</sup>

Scheme 1. Synthesis of boronate ester bullvalenes<sup>a</sup>



<sup>a</sup>Reagents and conditions: (a) alkyne (1.0 eq),  $CoBr_2(dppe)$  (10 mol%), ZnI<sub>2</sub> (20 mol%), Zn (30 mol%), DCE, RT. (b) 9*H* thioxanthen-9-one (1 mol%), THF, 365nm LED lamp, RT. (c) Br<sub>2</sub> (1.0 eq), KOtBu (1.0 eq), THF, -78 °C. (d) n-BuLi (1.0 eq), MeOBpin (1.0 eq), THF, -78 °C to RT. (e) MIDA (12.0 eq), DMSO, 120 °C, 8 hours. (f) alkyne (1.0 eq), CoBr<sub>2</sub>(dppe) (50 mol%), ZnI<sub>2</sub> (100 mol%), Zn (150 mol%), DCE, RT.

With this collection of boronate ester bullvalenes in hand we set out to demonstrate their ease of use in Suzuki cross-coupling reactions.<sup>20</sup> Mono-Bpin-bullvalene **1** couples with a range of aryl bromides to give the corresponding bullvalenes **13a-g** in good to excellent yields under simple and traditional catalytic conditions (Scheme 2a). Bis-Bpin-bullvalene **2** couples with bromo arenes to give diarylbullvalenes **14a-g** under identical reaction conditions (Scheme 2b).

MIDA boronate esters have emerged as excellent boronic acid masking group leading to effective linchpin cross-coupling strategies.<sup>21</sup> In the case of Bpin-B<sub>MIDA</sub>bullvalene **3**, the reactivity of the Bpin and B<sub>MIDA</sub> groups can be cleanly differentiated with exclusive monocoupling with *p*-bromobenzotrifluoride to give **15** in excellent yield (Scheme 3a). This was followed by a second coupling with a small range of aryl bromides to give differentially substituted diarybullvalenes **16a-c** (Scheme 3b). This strategy opens up a wide range of possibilities for the targeted synthesis of fluxional molecules.





<sup>a</sup>Reagents and conditions: (a) Ar-Br (1.1 eq), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), NaOH(aq), THF, 60 °C. (b) Ar-Br (2.2 eq), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), NaOH(aq), THF, 60 °C. Single-point DFT calculations performed at B3LYP-D3BJ/Def2-TZVPPD/CPCM solvent (chloroform) in Orca.
<sup>b</sup>See ref 26. <sup>c</sup>2-iodotoluene used in place of 2-bromotoluene.

Tri-Bpin-bullvalene **4** underwent coupling with bromobenzene under somewhat more forcing Suzuki

coupling conditions, but cleanly furnished triphenylbullvalene 17 in 71% yield (Scheme 3c). Triphenylbullvalene was previously prepared by Schröder and known to display intriguing dynamic behaviour.<sup>22</sup> In solution the major populated isomer is isomer B. However, recrystallisation gives exclusively the  $C_3$  symmetric isomer A, as confirmed by single crystal X-ray analysis. Isomer A is also kinetically metastable with an experimentally determined room-temperature solution half-life of 1h.<sup>22</sup> The origin of this kinetic feature was explored in a recent study by our laboratory and traced to high barriers in the first and second generation isomerisation from isomer A.<sup>23</sup> DFT analysis and kinetic modelling of triphenylbullvalene predict an isomer A half-life of 4.5 min (see SI for full details).

#### Scheme 3. Suzuki cross-coupling of bullvalenes 3 and 4ª



<sup>*a*</sup>Reagents and conditions: (a) p-bromobenzotrifluoride (1.1 eq), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), Ag<sub>2</sub>CO<sub>3</sub> (1.5 eq), THF, 60 °C. (b) Ar-Br (1.1 eq), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), NaOH(aq), THF, 60 °C. (c) Ph-Br (3.3 eq), Pd(PPh<sub>3</sub>)<sub>4</sub> (10 mol%), NaOH(aq), THF, 80 °C. Single-point DFT calculations performed at B3LYP-D3BJ/Def2-TZVPPD/CPCM solvent (chloroform) in Orca.

The solution phase isomer population distributions of most of the bullvalenes prepared herein was experimentally determined using low temperature NMR

measurements.<sup>24</sup> Population distributions follow predictable patterns. Substituents generally prefer the olefinic positions, with a slight preference for the positions adjacent to the apical bridgehead. A DFT survey of relative stabilities was conducted employing our previously reported methods.<sup>12</sup> The experimental and computational ratios are expressed as A:B, A:B:C, or A:(B,B'):C patterns (Schemes 2-4). The results are in excellent agreement with experiment for monosubstituted bullvalenes. For di- and tri-substituted bullvalenes agreement with experiment is qualitative, but calculations tend to somewhat overestimate the stability of isomer A. This is likely an artefact of the intramolecular dispersion interaction between the substituents.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28 29 30

31 32

33 34 35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

Diarylbullvalenes exist as dynamic ensembles of 15 isomers (homogeneous substituents) or 30 isomers (heterogeneous substituents), respectively. However, most of their time is spent within the A:B:C set of favoured isomers. Network analysis of diphenylbullvalene 14a exemplifies this character (Figure 2). The network graph represents nodes as isomers and edges as transitions structures; all colour coded according to calculated relative energy. The A:B:C set of isomers are closely interconnected through a low energy isomerisation path and will be in rapid equilibrium at ambient temperature.



**Figure 2.** Network analysis of diphenylbullvalene **14a**. Singlepoint DFT calculations performed at B3LYP-D3BJ/Def2-TZVPPD/CPCM solvent (chloroform) in Orca.

McGonigal recently reported a study on the crystallisation behaviour of substituted *barbaralanes* – shape-shifting hydrocarbons that exist as bistable dynamic ensembles.<sup>25</sup> Detailed analysis using a range of experimental and computational techniques led to the conclusion that crystallisation into a minor barbaralane solution phase isomer is due to general shape-selectivity rather than specific non-covalent interactions. Similar phenomena seem to play out herein for aryl substituted bullvalenes.

Single-crystal X-ray structures were obtained for bullvalenes 1, 2, 13c, 13d, 13e, 13f, 14a, 14c, 14d, 14g, 16a, 16c, 17, and 18 (selection shown in Figure 3a). For the monosubstituted bullvalenes, all compounds except 18 crystallise as the major solution phase isomer A.<sup>26</sup> Similarly, most disubstituted bullvalenes crystallise in their major solution phase isomer. However, for bis-Bpinbullvalene 2, aryl disubstituted variants 14c, 14g, 16a, and the trisubstituted 17, the inherent thermodynamic preference is overturned, and crystallisation gives a minor solution isomer. To guard against the possibility that single crystal structures might not reflect the bulk solid phase isomer preference, we recorded powder X-ray diffraction (PXRD) data for compounds 2, 14c, 14d, 16a, and 16c. Experimental data closely matches patterns simulated from the single crystal structures (see SI for full details).



**Figure 3.** (a) Single crystal X-ray structures of diarylbullvalenes. (b) Hirshfeld surface analysis. Single crystal X-ray structures of diarybullvalenes (grey - C, white - H, blue - N, red - O, and yellow - F). Hirshfield surface analysis for **16c** and **16a** with selected contacts highlighted. Regions of close contact significantly closer than the sum of their respective van der Waals radii are shown in red.

Diarylbullvalenes 16a and 16c provide an intriguing comparison. They are structurally very similar, differing only in the exchange of a phenyl group with a pyrimidine group, and would be expected to adopt nearly identical shapes. 16c follows the solution preference and recrystallises into the V-shaped isomer A in which the pyrimidine group of one molecule intercalates within the cleft of an adjacent molecule. The two molecules involved in this solid state dimer appear to be stabilised by weak C-H··· $\pi$  (D<sub>H- $\pi$ (centroid)</sub> = 2.62 Å) and offset  $\pi$ stacking interactions<sup>27</sup> ( $D_{\pi(centroid)-\pi(centroid)} = 4.00$  Å). The latter is reasonable as pyrimidine has a reasonably large dipole moment and in the A-type crystal, these dipoles are in an antiparallel orientation adding an additional electrostatic stabilization to the  $\pi$ -stacking. In contrast, 16a (as well as 14c) crystallise into the more elongated

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41 42

43 44

45

46

47

48

49 50

51

52

53 54

55

56

57 58 59

60

isomer B, contrary to the solution preference. In this case the broad cleft of one molecule encompasses the bullvalene core structure of an adjacent molecule. Curiously, no dominating intermolecular interactions appear to account for the stability of this crystal form despite its deviation from the preferred solution isomer; a bullvalene C-H…F contact ( $D_{H-F} = 2.48$  Å) appears to be the most significant contact.

Disubstituted bullvalenes 2, 14a, 14d, and 14g all crystallise in the V-shaped isomer A, for which only 14a (phenyl substituents) and 14g (pyrimidine) adopt the intercalated dimer arrangement seen for 16c. Presumably isomer A in 2 and 14d are prevented from close approach due to steric bulk and lack of an acidic CH for a weak C-H $\cdots \pi$  contact, respectively. Diphenylbullvalene 14a exhibits polymorphism between our single crystal structure, and that previously recorded by Echavarren.<sup>28</sup> Both structures are in the V-shaped isomer A, yet differ in the relative orientation of one of the phenyl groups. Consequently, our structure of 14a adopts the intercalated dimer arrangement whereas Echavarren's does not.

The intriguing crystallisation behaviour of substituted bullvalenes prompted us to holistically explore the packing arrangements computationally. For all di- and trisubstituted bullvalene crystal structures, the Hirshfeld surfaces<sup>29,30</sup> were calculated. This analysis highlights regions where intermolecular distances are significantly closer than the sum of their respective van der Waals radii. The results are illustrated for compounds 16c and **16a** (Figure 3b). Aside from the interactions noted above. only weak intermolecular interactions govern the packing. This conclusion is supported by DFT modelling (CE-B3LYP\*\*) of the intermolecular interaction energies for clusters of bullvalenes 2, 14a, 14c, 14d, 14g, 16a, 16c, and 17.<sup>30</sup> In all cases, the total interaction lattice energy is dominated by the dispersion energy contribution (see SI for full details). Spontaneous resolution of these ensembles in the crystal lattice appears to be governed by general shape-selectivity, which can override the inherent thermodynamic preferences observed in solution.

Scheme 4. Miscellaneous reactions of bullvalene 1<sup>a</sup>



<sup>a</sup>Reagents and conditions: (a) imidazole (2.0 eq), Cu(OAc)<sub>2</sub> (1.0 eq), B(OH)<sub>3</sub> (2.0 eq), powdered 4Å mol sieves, MeCN, 80 °C. (b) NaOH(aq),  $H_2O_2$ , THF, EtOH, 0 °C. (c) Benzaldehyde (1.2 eq) KHF<sub>2</sub> (1.2 eq), [Rh(COD)CI]<sub>2</sub> (1 mol%), dioxane, water, 80 °C. (d) 2-cyclohexen-1-one (1.1 eq) DIPEA (1.0 eq), 2-methylpropane-1,3 -diol (1.1 eq), [Rh(COD)CI]<sub>2</sub> (1 mol%), heptane, methanol, water, 80 °C. Single-point DFT calculations performed at B3LYP-D3BJ/Def2-TZVPPD/CPCM solvent (chloroform) in Orca.

Returning to the synthetic implications of this study, we wished to demonstrate the general utility of building blocks **1-4** beyond the scope of Suzuki cross-coupling reactions. Mono-Bpin-bullvalene **1** was subjected to a selection of transformations (Scheme 4). Chan-Lam cross-coupling with imidazole furnishes bullvalene **18**.<sup>31</sup> Simple oxidation with hydrogen peroxide smoothly gives bullvalone **19**.<sup>32</sup> Rhodium-catalysed 1,2- and 1,4-addition reactions to benzaldehyde and cyclohexenone gives bullvalenes **20** and **21**, respectively.<sup>33</sup> Despite their attachment to a constantly metamorphosing core structure, the reactivity of the pinacol boronate ester appears to be robust and reliable.

In summary, this work provides the first practical and general synthetic platform to prepare mono-, di-, and trisubstituted bullvalenes. Our boronate ester bullvalenes **1-4** are easy to make and have predictable reactivities. Bpin- $B_{MIDA}$ -bullvalene **3** is an effective linchpin enabling the logical synthesis of differentially disubstituted bullvalenes. We hope that these methods will spur further research and applications based on this most enigmatic hydrocarbon.

#### ASSOCIATED CONTENT

**Supporting Information**. The Supporting Information is available free of charge on the ACS Publications website. Experimental procedures and supporting figures with additional data (PDF).

#### **AUTHOR INFORMATION**

#### **Corresponding Author**

Thomas Fallon, thomas.fallon@adelaide.edu.au

### **Funding Sources**

We gratefully acknowledge the New Zealand Royal Society (Marsden Fund No. 15-MAU-154). Calculations were performed using the supercomputing infrastructure of the Computing Center of the Slovak Academy of Sciences

### REFERENCES

(1) (a) H. Hopf, Classics in Hydrocarbon Chemistry, Wiley-VCH, Weinheim, **2000**. (b) G. A. Olah, Á. Molnár, Hydrocarbon Chemistry, 2nd Edition, Wiley-VCH, Weinheim, **2003**.

(2) von E. Doering, W.; Roth, W. R. A rapidly reversible degenerate cope rearrangement: Bicyclo[5.1.0]octa-2,5-diene. *Tetrahedron* **1963**, *19*, 715–737.

(3) Zimmerman, H. E.; Grunewald, G. L. The chemistry of barrelene. III. A unique photoisomerization to semibullvalene. *J. Am. Chem. Soc.* **1966**, *88*, 183–184.

(4) (a) Barborak, J. C.; Daub, J.; Follweiler, D. M.; Schleyer, P. von R. Degenerate rearrangements of the 9-barbaralyl cation. *J. Am. Chem. Soc.* **1969**, *91*, 7760–7761. (b) Ahlberg, P.; Harris, D. L.; Winstein, S. Direct observation of the totally degenerate 9-barbaralyl cation and the bicyclo[4.3.0]nonatrienyl cation, a 1,4-bishomotropylium ion. *J. Am. Chem. Soc.* **1970**, *92*, 4454–4456.

(5) For a recent review, see: Ferrer, S.; Echavarren, A. M. Synthesis of bullvalenes: Classical approaches and recent developments. *Synthesis* **2019**, *51*, 1037–1048.

(6) (a) Schröder, G. Preparation and properties of tricyclo[3,3,2,04,6]deca-2,7,9-triene (bullvalene). *Angew. Chem. Int. Ed. Engl.* **1963**, *2*, 481–482. (b) Schröder, G. Synthese und eigenschaften von tricyclo[3.3.2.04.6]deca-2.7.9-trien (bullvalen). *Angew. Chem.* **1963**, *75*, 722–722. (c) Schröder, G. Synthese und eigenschaften von tricyclo[3.3.2.0<sup>4.6</sup>]decatrien-(2.7.9)<sup>2.3</sup>(bullvalen). *Chem. Ber.* **1964**, *97*, 3140–3149.

(7) (a) Schröder, G.; Merenyi, R.; Oth, J. F. M. Moleküle mit schneller und reversibler valenzisomerisierung.II: Synthese und eigenschaften von brombullvalen und tert-butyl-bullvalyläther. *Tetrahedron Lett.* **1964**, *5*, 773–777. (b) Oth, J. F. M.; Merényi, R.; Engel, G.; Schröder, G. Synthese und nmr-spektroskopisches verhalten zweier disubstituierter bullvalene. *Tetrahedron Lett.* **1966**, *7*, 3377– 3382.

(8) Ferrer, S.; Echavarren, A. M. Synthesis of barbaralones and bullvalenes made easy by gold catalysis. *Angew. Chem. Int. Ed.* **2016**, *55*, 11178–11182.

(9) (a) Lippert, A. R.; Keleshian, V. L.; Bode, J. W. Dynamic supramolecular complexation by shapeshifting organic molecules. *Org. Biomol. Chem.* 2009, *7*, 1529–1532. (b) Lippert, A. R. Naganawa, A. Keleshian, V. L. Bode, J. W. Synthesis of phototrappable shapeshifting molecules for adaptive guest binding. *J. Am. Chem. Soc.* 2010, *132*, 15790–15799. (c) He, M.; Bode, J. W. Racemization as a stereochemical measure of dynamics and robustness in shape-shifting organic molecules. *PNAS.* 2011, *108*, 14752–14756. (d) Larson, K. K.; He, M.; Teichert, J. F.; Naganawa, A.; Bode, J. W. Chemical sensing with shapeshifting organic molecules. *Chem. Sci.* 2012, *3*, 1825–1828. (e) He, M.; Bode, J. W. E pluribus unum: isolation, structure determination, network analysis and DFT studies of a single metastable structure from a shapeshifting mixture of 852 bullvalene structural isomers. *Org. Biomol. Chem.* 2013, *11*, 1306–1317.

(10) For recent reviews, see: (a) Rowan, S. J.; Cantrill, S. J.; Cousins, G. R. L.; Sanders, J. K. M.; Stoddart, J. F. Dynamic covalent chemistry. *Angew. Chem. Int. Ed.* 2002, *41*, 898–952. (b) Corbett, P. T.; Leclaire, J.; Vial, L.; West, K. R.; Wietor, J.-L.; Sanders, J. K. M.; Otto, S. Dynamic combinatorial chemistry. *Chem. Rev.* 2006, *106*, 3652–3711. (c) Jin, Y.; Yu, C.; Denman, R. J.; Zhang, W. Recent advances in dynamic covalent chemistry. *Chem. Soc. Rev.* 2013, *42*, 6634–6654.

(11) Teichert, J. F.; Mazunin, D.; Bode, J. W. Chemical sensing of polyols with shapeshifting boronic acids as a self-contained sensor array. *J. Am. Chem. Soc.* **2013**, *135*, 11314–11321.

(12) Yahiaoui, O.; Pašteka, L. F.; Judeel, B.; Fallon, T. Synthesis and analysis of substituted bullvalenes. *Angew. Chem. Int. Ed.* **2018**, *57*, 2570–2574.

Journal of the American Chemical Society

(13) (a) Achard, M.; Mosrin, M.; Tenaglia, A.; Buono, G. Cobalt(I)catalyzed [6+2] cycloadditions of cyclooctatetra(tri)ene with alkynes. *J. Org. Chem.* **2006**, *71*, 2907–2910. (b) D'yakonov, V. A.; Kadikova, G. N.; Dzhemileva, L. U.; Gazizullina, G. F.; Ramazanov, I. R.; Dzhemilev, U. M. *J.* Cobalt-catalyzed [6 + 2] cycloaddition of alkynes with 1,3,5,7-cyclooctatetraene as a key element in the direct construction of substituted bicyclo[4.3.1]decanes. *Org. Chem.* **2017**, *82*, 471–480.

(14) For reviews, see: (a) Hess, W.; Treutwein, J.; Hilt, G. *Synthesis* **2008**, 2008, 3537–3562. (b) Gandeepan, P.; Cheng, C.-H. Cobalt catalysis involving  $\pi$  components in organic synthesis. *Acc. Chem. Res.* **2015**, 48, 1194–1206. (c) Röse, P.; Hilt, G. Cobalt-catalysed bond formation reactions; part 2. *Synthesis* **2016**, 48, 463–492.

(15) The catalytic system used in Hilt's [4+2] cycloadditions of alkynyl boronate esters is analogous to the catalytic system reported in [6+2] cycloadditions. For selected works by Hilt see: (a) Hilt, G.; Smolko, K. I. Alkynylboronic esters as efficient dienophiles in cobalt-catalyzed diels–alder reactions. *Angew. Chemie Int. Ed.* **2003**, *42*, 2795–2797. (b) Hilt, G.; Lüers, S.; Smolko, K. I. A two-step reaction sequence for the syntheses of tetrahydronaphthalenes. *Org. Lett.* **2005**, *7*, 251–253. (c) Auvinet, A.-L.; Harrity, J. P. A.; Hilt, G. J. Ambient-temperature cobalt-catalyzed cycloaddition strategies to aromatic boronic esters. *Org. Chem.* **2010**, *75*, 3893–3896.

(16) Mono-Bpin-acetylene 6 and bis-Bpin-acetylene 7 are both commercially available. See the SI for details on their preparation.

(17) The use of alkynyl boronate esters, in particular bis-Bpinacetylene **7**, under these cobalt-catalysed reaction conditions was initially capricious and required extensive optimisation. We found that stringent purification and drying of all of the reaction components is necessary in order to suppress proto-deborylation of the substrate (see SI for full details). Proto-deborylation of bis-Bpin-acetylene **7** in a Co<sup>I</sup> catalysed [2+2+2] cycloaddition has been reported, see: Iannazzo, L.; Vollhardt, K. P. C.; Malacria, M.; Aubert, C.; Gandon, V. Alkynylboronates and -boramides in Co<sup>I</sup>- and Rh<sup>I</sup>-catalyzed [2+2+2] cycloadditions: Construction of oligoaryls through selective suzuki couplings. *Eur. J. Org. Chem.* **2011**, *2011*, 3283–3292.

(18) This reaction was selective for mono-condensation.

(19) Williams, J. D.; Nakano, M.; Gérardy, R.; Rincón, J. A.; de Frutos, Ó.; Mateos, C.; Monbaliu, J.-C. M.; Kappe, C. O. Finding the perfect match: A combined computational and experimental study toward efficient and scalable photosensitized [2 + 2] cycloadditions in flow. *Org. Process Res. Dev.* **2019**, *23*, 78–87.

(20) Stille couplings of triflate substituted bullvalene have been reported by Echavarren, see ref 8.

(21) Gillis, E. P.; Burke, M. D. Iterative cross-coupling with MIDA boronates: Towards a general platform for small molecule synthesis. *Aldrichimica Acta* **2009**, *42*, 17–27.

(22) Rebsamen, K.; Röttele, H.; Schröder, G. Mono-, di-, tri-, tetra-, penta-und hexaphenylbullvalene. *Chem. Ber.* **1993**, *126*, 1429–1433.

(23) Yahiaoui, O.; Pašteka, L. F.; Blake, C. J.; Newton, C. G.; Fallon, T. Network analysis of substituted bullvalenes. *Org. Lett.* **2019**, *21*, 9574–9578.

(24) Low temperature NMR measurements were made at -60 °C. In some cases, population analysis was precluded due to poor solubility (3, 14d), or severe signal overlap and/or restricted single bond rotation effects (14b, 15, 21).

(25) Bismillah, A. N.; Sturala, J.; Chapin, B. M.; Yufit, D. S.; Hodgkinson, P.; McGonigal, P. R. Shape-selective crystallisation of fluxional carbon cages. *Chem. Sci.* **2018**, *9*, 8631–8636.

(26) Echavarren reported the single X-ray crystal structure of **13a** (CCDC 1487024), see ref 8.

(27) Martinez, C. R.; Iverson, B. L. Rethinking the term "pistacking". Chem. Sci. 2012, 3, 2191–2201.

(28) For Echavarren's crystal structure of **14a** (CCDC 1487025), see ref 8.

(29) (a) Spackman, M.A.; McKinnon, J.J. Fingerprinting intermolecular interactions in molecular crystals. *CrystEngComm* 

59

60

, *4*, 378–392. (b) Spackman M. A.; Jayatilaka, D. Hirshfeld surface analysis. *CrystEngComm* **2009**, *11*, 19.

(30) All calculations were performed using the CrystalExplorer software package, version 17.5. http://crystalexplorer.scb.uwa.edu.au (last accessed 22/11/2019)

(31) Vantourout, J. C.; Miras, H. N.; Isidro-Llobet, A.; Sproules, S.; Watson, A. J. B. Spectroscopic studies of the chan-lam amination: A mechanism-inspired solution to boronic ester reactivity. *J. Am. Chem. Soc.* **2017**, *139*, 4769–4779.

(32) This 3-step procedure constitutes the shortest synthesis of bullvalone to date. For the original work on bullvalone, see: W. von E. Doering, W.; Ferrier, B. M.; Fossel, E. T.; Hartenstein, J. H.; Jones, M.;



Klumpp, G.; Rubin, R. M.; Saunders, M. A rational synthesis of bullvalene barbaralone and derivatives; bullvalone. *Tetrahedron* **1967**, *23*, 3943–3963.

(33) (a) Zhang, C.; Yun, J. Rh-catalyzed addition of β-carbonyl pinacol alkylboronates to aldehydes: Asymmetric synthesis of γbutyrolactones. *Org. Lett.* **2013**, *15*, 3416–3419. (b) Simmons, E. M.; Mudryk, B.; Lee, A. G.; Qiu, Y.; Razler, T. M.; Hsiao, Y. Development of a kilogram-scale process for the enantioselective synthesis of 3isopropenyl-cyclohexan-1-one via Rh/DTBM-SEGPHOS-catalyzed asymmetric hayashi addition enabled by 1,3-diol additives. *Org. Process Res. Dev.* **2017**, *21*, 1659–1667.