

Rh(III)-Catalyzed C–H Amidation of 2-Arylindoles with Dioxazolones: A Route to Indolo[1,2-*c*]quinazolines

Xiaogang Wang,^{†,||} Jin Zhang,^{*,†,||} Di Chen,^{†,‡} Bo Wang,[†] Xiufang Yang,[†] Yangmin Ma,^{*,†} and Michal Szostak^{*,†,§}


[†]College of Chemistry and Chemical Engineering, Shaanxi Key Laboratory of Chemical Additives for Industry, Shaanxi University of Science and Technology, Xi'an 710021, China

[‡]School of Food and Biological Engineering, Shaanxi University of Science and Technology, Xi'an 710021, China

[§]Department of Chemistry, Rutgers University, 73 Warren Street, Newark, New Jersey 07102, United States

S Supporting Information

ABSTRACT: Rhodium(III)-catalyzed C–H amidation of 2-arylindoles with dioxazolones for the synthesis of indolo[1,2-*c*]quinazolines is reported. The reaction is compatible with a wide range of electronically diverse 2-arylindoles and dioxazolones, providing indolo[1,2-*c*]quinazolines in high to excellent yields. Most notably, the combination of this Rh-catalyzed C–H amidation and intramolecular N–H/N–C(O) cyclization enables the most straightforward direct route to indolo[1,2-*c*]quinazolines to date. Mechanistic studies and evaluation of antitumor activity of these high value heterocycles are disclosed.



Indolo[1,2-*c*]quinazolines are an important class of indole-fused heterocycles that are widely represented in natural products, bioactive compounds, and organic photoluminescent devices (Figure 1).^{1,2} A classic method for the synthesis of





Figure 1. Examples of indolo[1,2-*c*]quinazolines.


indolo[1,2-*c*]quinazolines involves intramolecular condensation of 2-*ortho*-aminoarylindoles with carbonyl equivalents; however, this approach is inefficient, requires indole prefunctionalization, and is limited in terms of atom and step economy (Scheme 1A).^{3,4} Given our interest in amide bonds,^{5,6} we postulated that a direct route to this high value scaffold should be feasible utilizing C–H amidation of broadly available 2-arylindoles⁷ (Scheme 1B). We hypothesized that the capacity of N–H indoles to form amidorhodium⁸ species followed by C–H amidation^{9,10} together with a proximity driven electrophilic cyclization of the aromatic amide¹¹ might enable an expedient route to indolo[1,2-*c*]quinazolines. If successful, this would provide the most straightforward catalytic route to this high value heterocyclic scaffold to date.^{12–15}

Scheme 1. Previous Studies and This Work


A: Classical synthesis



B: This work: direct C–H amidation/N–H to C(O)–N cyclization



C: Previous studies: prefunctionalized substrates



Considering the synthetic importance of indolo[1,2-*c*]quinazolines, methods for the improved synthesis of this

Received: July 25, 2019

heterocycle have attracted a great deal of attention. For example, the studies by Zhang demonstrated that indolo[1,2-*c*]quinazolines can be accessed by Cu-catalyzed sequential amination (X = Br, I)/aerobic oxidative cyclization of 2-(2-haloaryl)-1*H*-indoles (Scheme 1C).¹² Fan reported Cu-catalyzed cascade amination/condensation/oxidative cyclization of 2-(2-bromoaryl)-1*H*-indoles (Scheme 1C).¹³ More recently, a Cu-catalyzed electrophilic amidation of 2-(2-amidoaryl)-1*H*-indoles was reported, wherein the Cu-catalyst acts as an O-coordinating Lewis acid to facilitate the cyclization.¹⁴ However, these methods start from ortho-substituted 2-arylindoles¹⁵ and are limited by atom economy.³ In sharp contrast, a direct synthesis of indolo[1,2-*c*]quinazolines by C–H amidation/N–H to N–C(O) condensation of 2-arylindoles would provide an attractive platform for generating a diversity of this important heterocycle.

In the past decade, group-directed C–H functionalization has been broadly developed for building valuable C–C and C–heteroatom bonds.¹⁶ Chang¹⁷ and Li¹⁸ have reported dioxazolones as electrophilic –NC(O)R sources for direct C–H amidation reactions.¹⁹ These highly reactive amidating reagents have numerous advantages that permit introducing valuable amide bonds by leveraging high affinity of the Lewis basic nitrogen atom with facile nitrenoid formation.²⁰ The amidation step could be combined with the subsequent cyclization for the synthesis of heterocycles, such as quinazolines,²¹ benzimidazoles,²² quinazolinone,²³ or quinazoline N-oxides.^{24,25} Herein, we report Rh(III)-catalyzed synthesis of indolo[1,2-*c*]quinazolines by a direct C–H amidation of 2-arylindoles with dioxazolones (Scheme 1B). The method enables the most straightforward diversity generating direct route to indolo[1,2-*c*]quinazolines and is characterized by high atom and step economy with H₂O and CO₂ formed as byproducts.

We commenced our studies by investigating the reaction between 2-phenyl-1*H*-indole (**1a**) and 3-phenyl-1,4,2-dioxazol-5-one (**2a**) to optimize the reaction conditions (Table 1). The reaction using [Cp*RhCl₂]₂ (5 mol %) and AgSbF₆ (25 mol %) as the catalytic system and CsOAc (25 mol %) as a basic additive in DCE at 140 °C afforded the desired C–H

amidation/N–H to N–C(O) cyclization product in 62% (entry 1). Among other catalytic systems, including [Cp*RhCl₂]₂, [Ru(*p*-cym)Cl₂]₂/AgSbF₆, [Cp*CoCOI₂]/AgSbF₆, and Pd(OAc)₂ (entries 2–5) only Co(III) afforded a trace of the desired product (<5%). A solvent screen revealed that DCE is optimal (entries 6–8). We could further improve the yield of (**3aa**) to 85% by using LiOAc as an additive (25 mol %) (entry 9), whereas other additives (NaOAc, KOAc, Cu(OAc)₂) were ineffective (entries 10–12). [Cp*Ir(III)Cl₂]₂ is not a competent catalyst for the reaction (recovery of the starting material). The C–H amidation does not proceed at 80 °C (recovery of the starting material).

With the optimized conditions in hand, the scope of the reaction was next investigated (Scheme 2). As shown, the scope of 2-aryl-1*H*-indole substrates is quite broad and

Scheme 2. Scope of 2-Aryl-1*H*-Indole Substrates^{a,b}




Table 1. Optimization of Reaction Conditions^a

entry	catalyst	additive	solvent	yield (%)
1	[Cp*RhCl ₂] ₂ /AgSbF ₆	CsOAc	DCE	62
2	[Cp*RhCl ₂] ₂	CsOAc	DCE	<5
3	[Ru(<i>p</i> -cym)Cl ₂] ₂ /AgSbF ₆	CsOAc	DCE	<5
4	[Cp*CoCOI ₂]/AgSbF ₆	CsOAc	DCE	<5
5	Pd(OAc) ₂	CsOAc	DCE	<5
6	[Cp*RhCl ₂] ₂ /AgSbF ₆	CsOAc	dioxane	<5
7	[Cp*RhCl ₂] ₂ /AgSbF ₆	CsOAc	toluene	12
8	[Cp*RhCl ₂] ₂ /AgSbF ₆	CsOAc	DMF	<5
9	[Cp*RhCl ₂] ₂ /AgSbF ₆	LiOAc	DCE	85
10	[Cp*RhCl ₂] ₂ /AgSbF ₆	NaOAc	DCE	44
11	[Cp*RhCl ₂] ₂ /AgSbF ₆	KOAc	DCE	<5
12	[Cp*RhCl ₂] ₂ /AgSbF ₆	Cu(OAc) ₂	DCE	<5


^aConditions: **1a** (1.0 equiv), **2a** (1.2 equiv), catalyst (5 mol %), additive (25 mol %), solvent (0.10 M), 140 °C, 24 h.

^aConditions: **1a** (1.0 equiv), **2a** (1.2 equiv), [Cp*RhCl₂]₂ (5 mol %), AgSbF₆ (25 mol %), LiOAc (25 mol %), DCE (0.10 M), 140 °C, 24 h. ^bIsolated yields. See Supporting Information (SI) for details.

accommodates a broad range of electron-donating (3ba–3da), electron-withdrawing (3ea–3ga), and halide groups (3ea) at the para position of the 2-aryl ring; however, 2-aryl-1*H*-indoles bearing electron-withdrawing groups gave lower yields (*vide infra*). It is noteworthy that a single regioisomer was formed using a meta-substituted substrate (3ha). Furthermore, the reaction was equally efficient when sterically hindered ortho-substituted 2-aryl-1*H*-indoles were used (3ia–3ja). The scope could be further extended to a naphthyl-substituted indole, furnishing a conjugated pentacyclic benzo[*g*]indolo[1,2-*c*]quinazoline (3ka). Importantly, 2-phenyl-1*H*-indoles bearing electron-donating (3ma–3na) as well as halide groups (3la, 3oa–3pa) at the 5- or 6-position of the indole ring could also be used to rapidly generate the indolo[1,2-*c*]quinazoline scaffold.

To further investigate the substrate scope, we subjected a range of dioxazolones to the reaction with 2-phenyl-1*H*-indole (Scheme 3). Pleasingly, we found that both 3-alkyl and 3-aryl

Scheme 3. Scope of 1,4,2-Dioxazol-5-one Substrates^{a,b}




substituted 1,4,2-dioxazol-5-one reagents could be used for the synthesis of indolo[1,2-*c*]quinazolines. The reactions with alkyl dioxazolones afforded the products in excellent yields (3ab–3ac). 3-Aryl-1,4,2-dioxazol-5-ones containing electron-donating (3ad–3ae), polyaromatic (3af), electron-withdrawing (3ag), and halide groups (3ah–3aj) worked well to afford functionalized indolo[1,2-*c*]quinazolines. The functional group

tolerance for an aryl bromide is noteworthy (3aj). Finally, we delighted to find that the reaction of 3,5-dimethoxyphenyl-1,4,2-dioxazol-5-one afforded 3ak in 68% yield. This product shows excellent antibacterial activity, highlighting the practical utility of the current protocol. Typically, we have not observed side reactions in this method. At the present stage of reaction development 4-nitrophenyl dioxazolone has not been tested. Future work will focus on expanding the scope of C–H amidation processes.


We conducted experiments to gain insight into the reaction mechanism (Scheme 4). Intermolecular competition studies

Scheme 4. Mechanistic Studies

A: Intermolecular competition: 2-aryl-1*H*-indoles **1**




B: Intermolecular competition: 2-aryl-1*H*-indoles **2**



C: Intermolecular competition: 1,4,2-dioxazol-2-ones **3**




D: Radical inhibition



between differently substituted 2-aryl-1*H*-indoles revealed electron-rich arenes to be inherently more reactive (Scheme 4A). Furthermore, electron-rich 2-phenyl-1*H*-indoles are inherently more reactive than their electron-deficient counterparts (Scheme 4B), while electron-deficient arenes are transferred preferentially from the dioxazolone amidation reagent (Scheme 4C). Moreover, radical inhibition studies revealed decreased yields of the amidation product (Scheme 4D). Overall, these studies are consistent with an electrophilic C–H activation mechanism, with the N–H to C(O)–N cyclization facilitated by electron-rich indoles.

A plausible reaction mechanism is shown in Scheme 5. The formation of an amidorhodium N–Rh(III) species and subsequent C–H activation gives a five-membered rhodacycle **4**.⁹ Coordination of dioxazolone and CO₂ extrusion gives a highly reactive Rh-imido intermediate **5**.¹⁸ Migratory insertion and cyclization gives the indolo[1,2-*c*]quinazoline product **3** and regenerates the catalyst. The electrophilic cyclization onto

Scheme 5. Proposed Catalytic Cycle



the amide is likely facilitated by a N–O–Rh migration to increase the amide bond electrophilicity.¹¹

The potential of the present method in the synthesis of complex indolo[1,2-*c*]quinazolines is highlighted in cytotoxic studies (Table 2). Intrigued by the unique capacity of this

Table 2. Cytotoxicity of 3aa–3ak in Human Cancer Cells^a

entry	compound	IC ₅₀ (μM)		
		PC3	A549	MCF-7
1	3aa	229.8	134.9	n/a
2	3pa	88.16	63.1	n/a
3	3ad	132.2	51.4	252
4	3ag	242.8	30.7	n/a
5	3ak	158.4	87.4	287.9
6	cisplatin	4.97	12.2	70.4

^aSelected results are shown. See SI for full details.

valuable heterocyclic scaffold to act as a privileged synthon in drug discovery, we tested the synthesized products against human prostate cancer cells (PC3), human lung cancer cells (A549), and human Michigan cancer foundation cells (MCF-7) (see SI for details).²⁶ As shown, the compounds 3pa, 3ad, 3ag, and 3ak displayed promising growth inhibition of A549 cells (3ag: IC₅₀ = 30.7 μM). The results demonstrate that indolo[1,2-*c*]quinazolines could represent a new class of cancer growth inhibitors.²⁷

In summary, we have reported Rh-catalyzed synthesis of indolo[1,2-*c*]quinazolines through direct C–H amidation and intramolecular N–H/N–C(O) cyclization of broadly available 2-aryl-1H-indoles with dioxazolones. This mild and highly efficient process permits the most straightforward platform for the synthesis of indolo[1,2-*c*]quinazolines developed to date. The reaction showed broad scope with respect to indole and dioxazolone. The reaction proceeds with high atom economy with H₂O and CO₂ formed as byproducts and avoids any substrate prefunctionalization. The potential of this method has been highlighted by the discovery of indolo[1,2-*c*]quinazolines that exhibit promising antitumor activity. Further studies on the formation of amides via C–H functionalization as well as tandem annulation are underway in our laboratory.

■ ASSOCIATED CONTENT

§ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.9b02615.

Experimental procedures and characterization data (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: zhangjin@sust.edu.cn.

*E-mail: mym63@sina.com.

*E-mail: michal.szostak@rutgers.edu.

ORCID

Jin Zhang: 0000-0002-6616-7087

Michal Szostak: 0000-0002-9650-9690

Author Contributions

^{||}X.W. and J.Z. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

J.Z. thanks the China Scholarship Council (No. 2018086-10096). Financial support was provided by the Foundation for Young Scholars of Shaanxi University of Science and Technology (No. BJ12-26). M.S. thanks Rutgers University and the NSF (CAREER CHE-1650766).

■ REFERENCES

- (a) Blackman, A. J.; Hambley, T. W.; Picker, K.; Taylor, W. C.; Thirasasana, N. Hinckdentine-A: a novel alkaloid from the marine bryozoan Hinckinsoflustra denticulate. *Tetrahedron Lett.* **1987**, 28, 5561–5562. (b) Liu, Y.; McWhorter, W. W. Synthesis of 8-desbromohinckdentine A1. *J. Am. Chem. Soc.* **2003**, 125, 4240–4252.
- (a) Rohini, R.; Shanker, K.; Reddy, P. M.; Sekhar, V. C.; Ravinder, V. 6-substituted indolo[1,2-*c*]quinazolines as new antimicrobial agents. *Arch. Pharm.* **2009**, 342, 533–540. (b) Rohini, R.; Reddy, P. M.; Shanker, K.; Hu, A.; Ravinder, V. Antimicrobial study of newly synthesized 6-substituted indolo[1,2-*c*]quinazolines. *Eur. J. Med. Chem.* **2010**, 45, 1200–1205. (c) Sharma, V.; Kumar, P.; Pathak, D. Biological importance of the indole nucleus in recent years: a comprehensive review. *J. Heterocycl. Chem.* **2010**, 47, 491–502. (d) Sravanthi, T. V.; Manju, S. L. Indoles - A promising scaffold for drug development. *Eur. J. Pharm. Sci.* **2016**, 91, 1–10. (e) Shin, C.; Kim, K.; Hoon, D.; Jang, J.; Chun, M. Aromatic compounds as electroluminescent material and/or semiconductor material for organic electroluminescent devices. WO2012173371 2012, Dec 20, 2012. (f) Lee, D. U.; Bae, J. S.; Nam, H.; Hong, S. G.; Lee, D. H.; Jeon, S. Y. Organic electroluminescent material and organic electroluminescent apparatus utilizing the electroluminescent material. KR 2010082986 2010, Jul 21, 2010.
- Joule, J. A.; Mills, K. *Heterocyclic Chemistry*; Wiley-Blackwell: Oxford, 2010.
- Li, C. J.; Trost, B. M. Green chemistry for chemical synthesis. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, 105, 13197–13202.
- (a) Shi, S.; Nolan, S. P.; Szostak, M. Well-defined palladium(II)-NHC (NHC = N-Heterocyclic Carbene) precatalysts for cross-coupling reactions of amides and esters by selective acyl CO–X (X = N, O) cleavage. *Acc. Chem. Res.* **2018**, 51, 2589–2599. (b) Liu, C.; Szostak, M. Decarbonylative cross-coupling of amides. *Org. Biomol. Chem.* **2018**, 16, 7998–8010. (c) Li, G.; Ji, C. L.; Hong, X.; Szostak, M. Highly chemoselective, transition-metal-free transamidation of

unactivated amides and direct amidation of alkyl esters by N-C/O-C cleavage. *J. Am. Chem. Soc.* **2019**, *141*, 11161–11172.

(6) For representative studies from our laboratory, see: (a) Zhang, J.; Hou, Y.; Ma, Y.; Szostak, M. Synthesis of amides by mild palladium-catalyzed aminocarbonylation of arylsilanes with amines enabled by copper(II) fluoride. *J. Org. Chem.* **2019**, *84*, 338–345. (b) Nan, J.; Hu, Y.; Chen, P.; Ma, Y. Metal-free synthesis of 2-fluoroalkylated quinolines using polyfluoroalkanoic acids as direct fluorine sources. *Org. Lett.* **2019**, *21*, 1984–1988. (c) Liu, S.; Gao, W.; Miao, Y.; Wang, M. Dinuclear zinc-azephenol catalyzed asymmetric aza-Henry reaction of *N*-Boc imines and nitroalkanes under ambient conditions. *J. Org. Chem.* **2019**, *84*, 2652–2659.

(7) Lal, S.; Snape, T. J. 2-Arylindoles: a privileged molecular scaffold with potent, broad-ranging pharmacological activity. *Curr. Med. Chem.* **2012**, *19*, 4828–4837.

(8) (a) Morimoto, K.; Hirano, K.; Satoh, T.; Miura, M. Rhodium-catalyzed oxidative coupling/cyclization of 2-phenylindoles with alkynes via C-H and N-H bond cleavages with air as the oxidant. *Org. Lett.* **2010**, *12*, 2068–2071. (b) Yamauchi, D.; Nishimura, T.; Yorimitsu, H. Asymmetric hydroarylation of vinyl ethers catalyzed by a hydroxiridium complex: azoles as effective directing groups. *Chem. Commun.* **2017**, *53*, 2760–2763.

(9) For an excellent review on C–H amination, see: Park, Y.; Kim, Y.; Chang, S. Transition metal-catalyzed C–H amination: scope, mechanism, and applications. *Chem. Rev.* **2017**, *117*, 9247–9301.

(10) For an excellent review on non-classical methods of synthesis of amides, see: de Figueiredo, R. M.; Suppo, J. S.; Campagne, J. M. Nonclassical routes for amide bond formation. *Chem. Rev.* **2016**, *116*, 12029–12122.

(11) For a review on using as amides as N–C(O) electrophiles, see: Greenberg, A.; Breneman, C. M.; Liebman, J. F. *The Amide Linkage: Structural Significance in Chemistry, Biochemistry and Materials Science*; Wiley-VCH: New York, 2003.

(12) Sang, P.; Xie, Y. J.; Zou, J. W.; Zhang, Y. H. Copper-catalyzed sequential Ullmann *N*-arylation and aerobic oxidative C–H amination: a convenient route to indolo[1,2-*c*]quinazoline derivatives. *Org. Lett.* **2012**, *14*, 3894–3897.

(13) Guo, S. H.; Tao, L.; Zhang, W. W.; Zhang, X. Y.; Fan, X. S. Regioselective synthesis of indolo[1,2-*c*]quinazolines and 11*H*-Indolo[3,2-*c*]quinolines via copper-catalyzed cascade reactions of 2-(2-bromoaryl)-1*H*-indoles with aldehydes and aqueous ammonia. *J. Org. Chem.* **2015**, *80*, 10955–10964.

(14) Guo, S.; Wang, F.; Tao, L.; Zhang, X.; Fan, X. Solvent-dependent copper-catalyzed indolyl C3-oxygenation and N1-cyclization reactions: selective synthesis of 3*H*-indol-3-ones and indolo[1,2-*c*]quinazolines. *J. Org. Chem.* **2018**, *83*, 3889–3896.

(15) (a) For the classic Kiang–Mann synthesis using acyl halides, see: Kiang, A. K.; Mann, F. G.; Prior, A. F.; Topham, A. Action of acyl cyanides on 2- and 1,2-substituted indoles. II. Derivatives of 2-*o*-aminophenylindole. *J. Chem. Soc.* **1956**, 1319–1331. (b) For an oxidative variant using aldehydes followed by oxidation, see refs 2a and b.

(16) For selected reviews on C–H activation, see: (a) *Science of Synthesis: Catalytic Transformations via C–H Activation*; Yu, J. Q., Ed.; Thieme: Stuttgart, 2015. (b) Davies, H. M. L.; Morton, D. Recent advances in C–H functionalization. *J. Org. Chem.* **2016**, *81*, 343–350. (c) Rossi, R.; Bellina, F.; Lessi, M.; Manzini, C. Cross-coupling of heteroarenes by C–H functionalization: Recent progress towards direct arylation and heteroarylation reactions involving heteroarenes containing one heteroatom. *Adv. Synth. Catal.* **2014**, *356*, 17–117. (d) Rouquet, G.; Chatani, N. Catalytic functionalization of C(sp³)-H and C(sp²)-H bonds by using bidentate directing groups. *Angew. Chem., Int. Ed.* **2013**, *52*, 11726–11743. (e) Yeung, C. S.; Dong, V. M. Catalytic dehydrogenative cross-coupling: forming carbon–carbon bonds by oxidizing two carbon–hydrogen bonds. *Chem. Rev.* **2011**, *111*, 1215–1292. (f) Lyons, T.; Sanford, M. Palladium-catalyzed ligand-directed C–H functionalization reactions. *Chem. Rev.* **2010**, *110*, 1147–1169. (g) Ackermann, L.; Vicente, R.; Kapdi, A. R. Transition-metal-catalyzed direct arylation of (hetero)arenes by C–H

bond cleavage. *Angew. Chem., Int. Ed.* **2009**, *48*, 9792–9826. (h) Gandeepan, P.; Müller, T.; Zell, D.; Cera, G.; Warratz, S.; Ackermann, L. 3d Transition Metals for C–H Activation. *Chem. Rev.* **2019**, *119*, 2192–2452. (i) Sambiagio, C.; Schönbauer, D.; Blicke, R.; Dao-Huy, T.; Pototschnig, G.; Schaaf, P.; Wiesinger, T.; Zia, M. F.; Wencel-Delord, J.; Besset, T.; Maes, B. U. W.; Schnürch, M. A comprehensive overview of directing groups applied in metal-catalysed C–H functionalisation chemistry. *Chem. Soc. Rev.* **2018**, *47*, 6603–6743.

(17) (a) Park, Y.; Park, K. T.; Kim, J. G.; Chang, S. Mechanistic studies on the Rh(III)-mediated amido transfer process leading to robust C–H amination with a new type of amidating reagent. *J. Am. Chem. Soc.* **2015**, *137*, 4534–4542. (b) Park, J.; Chang, S. Comparative catalytic activity of group 9 [Cp**M*III] complexes: cobalt-catalyzed C–H amidation of arenes with dioxazolones as amidating reagents. *Angew. Chem., Int. Ed.* **2015**, *54*, 14103–14107.

(18) Wang, H.; Tang, G. D.; Li, X. W. Rhodium(III)-catalyzed amidation of unactivated C(sp³)-H bonds. *Angew. Chem., Int. Ed.* **2015**, *54*, 13049–13052.

(19) For a mechanistic study, see: Park, Y.; Heo, J.; Baik, M. H.; Chang, S. Why is the Ir(III)-mediated amido transfer much faster than the Rh(III)-mediated reaction? A combined experimental and computational study. *J. Am. Chem. Soc.* **2016**, *138*, 14020–14029.

(20) For recent elegant intramolecular amidations using dioxazolones, see: (a) Hong, S. Y.; Park, Y.; Hwang, Y.; Kim, Y. B.; Baik, M. H.; Chang, S. Selective formation of γ -lactams via C–H amidation enabled by tailored iridium catalysts. *Science* **2018**, *359*, 1016–1021. (b) Park, Y.; Chang, S. Asymmetric formation of γ -lactams via C–H amidation enabled by chiral hydrogen-bond-donor catalysts. *Nat. Catal.* **2019**, *2*, 219–227.

(21) (a) Wang, F.; Wang, H.; Wang, Q.; Yu, S.; Li, X. Co(III)-catalyzed synthesis of quinazolines via C–H activation of *N*-sulfonylimines and benzimidates. *Org. Lett.* **2016**, *18*, 1306–1309. (b) Wang, J.; Zha, S.; Chen, K.; Zhang, F.; Song, C.; Zhu, J. Quinazoline synthesis via Rh(III)-catalyzed intermolecular C–H functionalization of benzimidates with dioxazolones. *Org. Lett.* **2016**, *18*, 2062–2065.

(22) (a) Xia, J.; Yang, X.; Li, Y.; Li, X. Iridium(III)-catalyzed synthesis of benzimidazoles via C–H activation and amidation of aniline derivatives. *Org. Lett.* **2017**, *19*, 3243–3246. (b) Chen, Y. Y.; Zhang, R.; Peng, Q. J.; Xu, L. T.; Pan, X. H. Rhodium(III)-catalyzed directed C–H amidation of *N*-nitrosoanilines and subsequent formation of 1,2-disubstituted benzimidazoles. *Chem. - Asian J.* **2017**, *12*, 2804–2808.

(23) Mishra, A.; Mukherjee, U.; Vats, T. K.; Deb, I. Ir(III)/MPAA-catalyzed mild and selective C–H amidation of *N*-sulfonyl ketimines: Access to benzosultam-fused quinazolines/quinazolinones. *J. Org. Chem.* **2018**, *83*, 3756–3767.

(24) Wang, Q.; Wang, F.; Yang, X. F.; Zhou, X. K.; Li, X. W. Rh(III)- and Zn(II)-catalyzed synthesis of quinazoline *N*-oxides via C–H amidation-cyclization of oximes. *Org. Lett.* **2016**, *18*, 6144–6147.

(25) For additional representative examples, see: (a) Wang, F.; Jin, L.; Kong, L.; Li, X. Cobalt(III)- and rhodium(III)-catalyzed C–H amidation and synthesis of 4-quinolones: C–H activation assisted by weakly coordinating and functionalizable. *Org. Lett.* **2017**, *19*, 1812–1815. (b) Mei, R.; Loup, J.; Ackermann, L. Oxazolonyl-assisted C–H amidation by cobalt(III) catalysis. *ACS Catal.* **2016**, *6*, 793–797. (c) Hermann, G. N.; Bolm, C. Mechanochemical rhodium(III)-catalyzed C–H bond amidation of arenes with dioxazolones under solventless conditions in a ball mill. *ACS Catal.* **2017**, *7*, 4592–4596. (d) Wang, H.; Lorion, M. M.; Ackermann, L. Overcoming *N*-heterocycles by cobalt catalysis. *Angew. Chem., Int. Ed.* **2016**, *55*, 10386–10390. (e) Zhou, Y.; Engl, O. D.; Bandar, J. S.; Chant, E. D.; Buchwald, S. L. CuH-catalyzed asymmetric hydroamidation of vinylarenes. *Angew. Chem., Int. Ed.* **2018**, *57*, 6672–6675. (f) Halskov, K. S.; Roth, H. S.; Ellman, J. A. Synthesis of [5,6]-bicyclic heterocycles with a ring-junction nitrogen atom: Rhodium(III)-catalyzed C–H functionalization of alkenyl azoles. *Angew. Chem., Int. Ed.* **2017**, *56*,

9183–9187. For representative direct intramolecular amidations, see:

(g) Wu, X.; Zhao, Y.; Ge, H. Direct aerobic carbonylation of C(sp^2)-H and C(sp^3)-H bonds through Ni/Cu synergistic catalysis with DMF as the carbonyl source. *J. Am. Chem. Soc.* **2015**, *137*, 4924–2927.

(h) Wu, X.; Yang, K.; Zhao, Y.; Sun, H.; Li, G.; Ge, H. Cobalt-catalysed site-selective intra- and intermolecular dehydrogenative amination of unactivated sp^3 carbons. *Nat. Commun.* **2015**, *6*, 6462.

(26) Berridge, M. V.; Herst, P. M.; Tan, A. S. Tetrazolium dyes as tools in cell biology: new insights into their cellular reduction. *Biotechnol. Annu. Rev.* **2005**, *11*, 127–152.

(27) Dadashpour, S.; Emami, S. Indole in the target-based design of anticancer agents: A versatile scaffold with diverse mechanisms. *Eur. J. Med. Chem.* **2018**, *150*, 9–29.