

Reductive Arylation of Arylidene Malonates Using Photoredox Catalysis


Rick C. Betori and Karl A. Scheidt*

Department of Chemistry, Center for Molecular Innovation and Drug Discovery, Northwestern University, 2145 Sheridan Road, Evanston, Illinois 60208, United States

Supporting Information

ABSTRACT: A strategy with arylidene malonates provides access to β -umpolung single-electron species. Reported here is the utilization of these operators in intermolecular radical–radical arylations, while avoiding conjugate addition/dimerization reactivity that is commonly encountered in enone-based photoredox chemistry. This reactivity relies on tertiary amines that serve to both activate the arylidene malonate for single-electron reduction by a proton-coupled electron transfer mechanism as well as serve as a terminal reductant. This photoredox catalysis pathway demonstrates the versatility of stabilized radicals for unique bond-forming reactions.

KEYWORDS: photoredox catalysis, arylation, proton coupled electron transfer, arylidene malonates, β -umpolung




The study of underexplored reactive intermediates has driven advances in carbon–carbon and carbon–heteroatom bond-forming reactions, leading to powerful transformations that afford construction of both simple and structurally complex molecular architectures.¹ Among established methods, disconnections utilizing inverse polarity concepts, termed umpolung, have been a significant focus and represent an appealing route for the preparation of chiral molecules.² N-heterocyclic carbenes (NHCs) have emerged as unique catalysts for umpolung reactivity, where NHC-homoenolate equivalents have been thoroughly explored as a means of polarity-reversed reactivity of conjugate acceptors.³

More recently, the development of photoredox catalysis has yielded new opportunities for the preparation of unconventional operators under mild conditions.⁴ Lowest unoccupied molecular orbital (LUMO)-lowering photoredox cooperative catalysis, such as proton-coupled electron transfer (PCET)⁵ and Lewis acid/photoredox cooperative photoredox,⁶ has facilitated the development of d_1 (ketyl) and d_3 (enone) umpolung synthons previously inaccessible for single-electron reduction by traditional visible light-absorbing photocatalysts (Figure 1A). Recent reports utilizing bi-functional Lewis acid/visible light catalysis⁷ and covalent interactions to facilitate bathochromic activation and subsequent redox chemistry have expanded this emerging field to access new chemical reactivity.⁸


As part of our ongoing program to generate new opportunities in β -umpolung reactivity, we recently reported the exploitation of arylidene malonates as substrates in photoredox/Lewis acid-cooperative catalysis to afford radical–radical cross-couplings, radical dimerizations, transfer hydrogenations, and reductive annulations.⁹ These studies focused on forming intermediates to access previously unexplored chemical reactivities, namely, nondimerization/cycloaddition reactions often seen with enone-derived radicals

Previous Work

A) Single-Electron Approaches to Umpolung Reactivity




Proton Coupled Electron Transfer for Umpolung Reactivity




B) Cyanoarenes in Photoredox Arylation Chemistry


α -Arylation of Amines




Direct Allylic Arylation



β -Arylation of Saturated Carbonyls



Umpolung Arylation of Carbonyls



This Work

C) β -Umpolung Arylation of Arylidene Malonates




Figure 1. Photoredox umpolung and arylations strategies.

(Figure 1A).^{6a–c,10} We demonstrated that arylidene malonates [$E_{1/2}^{\text{red}} = -1.57$ V for **1a** vs saturated calomel electrode (SCE)]¹¹ demonstrate substantial capacity in LUMO-lowering catalysis as shown by a dramatic change in reduction potential

Received: August 23, 2019

Revised: September 27, 2019

upon complexation with a Lewis acid ($E_{1/2}$ red = -0.37 V vs SCE, **1a** + 100 mol % $\text{Sc}(\text{OTf})_3$). In particular, increased resonance stabilization of the β -radical enolate intermediate provides a versatile species that demonstrates reactivity divergent from conventional β -enones.

We postulated that this stable β -radical enol would be able to undergo cross-coupling reactions with cyanoarene-derived radicals, producing β -arylation products from the arylidene malonate. Photoredox arylation using cyanoarenes has attracted significant interest primarily because of the inherent value of direct arene functionalization. In particular, MacMillan and co-workers have demonstrated α -amino arylations,¹² arylations of allylic sp^3 C–H bonds,¹³ and β -arylation of saturated aldehydes and ketones.¹⁴ Of note is the facile access of these synthons using photoredox chemistry, enabling their broad applicability (Figure 1B).¹⁵ Herein, we report the utilization of these β -radical intermediates through a photoredox reductive arylation of arylidene malonates with cyanoarenes to provide densely functionalized diarylmalonates (Figure 1C).

To facilitate rapid reaction development, we utilized high-throughput experimentation (HTE) to allow for the execution of a large number of experiments to be conducted in parallel while requiring less effort and cost per experiment when compared to traditional means of experimentation (Scheme 1, see Supporting Information for details).¹⁶ Variables that were considered for optimization were as follows: photocatalyst, terminal reductant, Lewis acid, and solvent, where equivalents of cyanoarene (2.0 equiv), terminal reductant (2.5 equiv),


photocatalyst concentration (3 mol %), and reaction concentration (0.1 M) were held constant throughout. Gratifyingly, the diaryl product **2a** was formed with DPAIPN and 2.5 equiv of NEt_3 in CH_3CN in 85% yield on 5 μmol scale (see the Supporting Information for yields of all reactions in 96-well plate), where these results were validated on an initial scale up to 0.2 mmol to provide **2a** in 85% yield (see Supporting Information for follow up optimization on isolable scales).

Observable trends demonstrate that a photocatalyst must be both a capable oxidant and reductant (DPAIPN, $\text{Ir}(\text{ppy})_2\text{dtbpy}$), where catalysts that are strong oxidants/mild reductants (Ph-MesAcr) provided no reactivity, and strong reductants/mild oxidants ($\text{Ir}(\text{ppy})_3$) resulted in dimerization of **1a**.^{9a} Evaluations of terminal reductants demonstrated that tertiary amines were superior, where the Hantzsch ester (HE) was only able to afford **2a** in trace quantities. This highlights the differences between tertiary amines and other terminal reductants. Tertiary amines can, upon initial single-electron reduction, result in the oxidized nitrogen atom forming a 2-center/3- e^- interaction¹⁷ or, after a [1,2]-H shift, serving as a hydrogen-bond donor.¹⁸ This results in the oxidized amine serving as both the terminal reductant and the Lewis acidic intermediate necessary for activation of **1a**.¹⁹ It is likely that the [1,2]-H-shift to form a Brønsted acid is necessary, as substitution of triethylamine for triphenylamine, a tertiary amine unable to undergo a [1,2]-H-shift provided minimal product either as the sole terminal reductant or used alongside $\text{NEt}_3/\text{NHET}_3\text{Cl}$ (see Supporting Information).²⁰ Because of the dual terminal reductant/Lewis acid role of NEt_3 , inclusion of an exogenous Lewis acid proved deleterious due to a more complex reaction profile. A series of control experiments demonstrated that the reaction did not take place in the absence of photocatalyst or light (see the Supporting Information for details). Moreover, we were pleased to find that the optimized reaction conditions were not sensitive to operational variations in reaction conditions, where differences in temperature ($\pm 15^\circ\text{C}$), concentration (0.05–0.3 M), H_2O level (addition of desiccant—10 equiv H_2O), light intensity, and O_2 level all resulted in minimal changes in the observed yield (Scheme 2, see Supporting Information).²¹


With these optimized conditions, we investigated a variety of substrates (Table 1). Generally, desired products were obtained in good to excellent yields. Variations of the arylidene malonate were well tolerated, where electron-rich and electron-poor containing compounds were accessed in high yields (Table 1, **2a–2m**).

Diversity could be introduced into the dicarbonyl moiety to tolerate a variety of diesters (Table 1, **2n–2p**). A variety of heteroaromatic systems were highly efficient in this reaction, as pyridine, indole, furan, and pyrrole-containing arylidene malonates afforded the diaryl species (Table 1, **2q–2t**). Unfortunately, alkylidene malonates are not successful with these conditions (**1u**), presumably because of the increased reduction potential relative to their arylidene malonate counterparts (see the Supporting Information). Moreover, substrates designed to afford quaternary centers were also unsuccessful (**1v**). Unsubstituted pyridines and quinolones only provided transfer hydrogenation products (**1w–1aa**). Substrates derived from Meldrum's acid showed no conversion under the optimized conditions (**1ab**), presumably because of an inability of **1ab** to coordinate the NEt_3 radical cation or NHET_3^+ to engage in PCET. Lastly, benzylidenepentane-2,4-

Scheme 1. Reaction Optimization Using HTE^a



^aReactions conducted on 5 μmol scale. Reactions were irradiated with Lumidox 456 nm light-emitting diodes (LEDs) for 18 h. Yields determined by ultra-performance liquid chromatography–mass spectrometry (MS) using naphthalene as internal standard. See Supporting Information.


Scheme 2. Reaction-Condition Sensitivity Assessment^a

^aReaction conditions: constant: **1** (0.2 mmol), 4-CN-pyridine (0.4 mmol), NEt₃ (0.5 mmol) DPAIPN (3 mol %). Reaction was irradiated with Kessil PR160 456 nm LEDs for 18 h. Yields determined by gas chromatography MS using biphenyl as internal standard. See [Supporting Information](#).

dione (**1ac**) provided a complex reaction mixture likely because of ketyl radical formation, leading to undesired side reactivity. Arene variation allowed for differentially substituted pyridines, pyrimidines, pyrroles, and indoles ([Table 2](#)). Attempts to utilize 1,4-dicyanobenzene (1,4-DCB) provided no product, likely because of SET from DPAIPN to 1,4-DCB ($E_{1/2}$ red = −1.52 V vs SCE for DPAIPN radical anion) being endergonic ($E_{1/2}$ red = 1.67 V vs SCE for 1,4-DCB).

While this process affords similar connectivity compared to transition-metal catalyzed conjugate arylations²² and organometallic reagent conjugate additions,²³ it is noteworthy that the use of heteroaryl conjugate acceptors or heteroaryl nucleophiles is problematic, likely because of Lewis basic functionality interacting with the metal catalysts.²⁴


While the use of arylidene malonates as conjugate acceptors in Friedel–Crafts arylation reactions is a well-established paradigm,²⁵ nearly all of the products prepared using this method would not be possible using established methods, highlighting the value of this method. To compare against alternative approaches to afford diarylmalonates, we attempted to prepare **2a** through a Knoevenagel/hydrogenation or organometallic addition into **1a** and a Lewis-acid-catalyzed dehydration coupling from diarylmethanol **5a** using established procedures ([Scheme 3](#)).²⁶ Additionally, we tried a variety of

Table 1. Arylidene Malonate Reaction Scope^a

^aReaction conditions: **1** (0.2 mmol), 4-CN-pyridine (0.4 mmol), NEt₃ (0.5 mmol) DPAIPN (3 mol %), CH₃CN (2.0 mL) was irradiated with Kessil PR160 456 nm LEDs for 18 h. Reported yields are determined after isolation by chromatography. See [Supporting Information](#).


different organometallic and transition-metal catalyzed procedures for conjugate addition into the arylidene malonate. Lastly, a series of dehydrogenative coupling reactions from diarylmalonate **6a** were also attempted.²⁷ Unfortunately, all conditions screened (see [Supporting Information](#) for details) were unsuccessful either leading to no conversion of starting material or decomposition under the reaction conditions, highlighting the utility of this process.

To demonstrate utility of this methodology on a multigram scale, we linearly scaled the reaction 1000-fold from the initial screening conditions to access **2a** in 85% yield. Additionally, >90% of the DPAIPN photocatalyst was recovered via column chromatography and was able to again reproduce the title

Table 2. Arene Reaction Scope^a

^aReaction conditions: 1a (0.2 mmol), arene (0.4 mmol), NEt₃ (0.5 mmol) DPAIPN (3 mol %), CH₃CN (2.0 mL) was irradiated with Kessil PR160 456 nm LEDs for 18 h. Reported yields are determined after isolation by chromatography. See the [Supporting Information](#).

Scheme 3. Comparison to Known Methods




reaction on gram scale without loss of yield (Scheme 4A). We then evaluated a tandem one-pot Knoevenagel/arylation process starting from benzaldehyde and dimethyl malonate, where we were pleased to find that 2a could be accessed in excellent yield after purification. Moreover, we found this process could be telescoped further by concentration and redissolution of the crude arylation reaction mixture in 1:9 H₂O/dimethyl sulfoxide (DMSO) with LiCl to provide the Krapcho product 7a in 64% yield over the three-transformation process (Scheme 4B,C).


To study the mechanism of this process, we employed fluorescence-quenching techniques with 1a and 4-CN pyridine as model substrates (Scheme 5A).²⁸ Stern–Volmer analysis demonstrated that neither 1a nor 4-CN pyridine quench the excited state of DPAIPN [$E_{1/2}$ (DPAIPN*/DPAIPN^{•+}) = 1.28 V vs SCE] in acetonitrile at 25 °C. However, addition of NEt₃ resulted in a large decrease in the measured fluorescence, indicating that this reaction proceeds through a reductive quenching mechanism of DPAIPN [$E_{1/2}$ (DPAIPN*/DPAIPN^{•-}) = 1.10 V vs SCE] by NEt₃ ($E_{1/2}$ ox = 0.83 V vs

Scheme 4. Gram-Scale Reaction and Telescope Process


A) Gram Scale Reaction and Recovery/Reusability of Photocatalyst



B) Single-Flask Knoevenagel/Arylation




C) Single-Flask Knoevenagel/Arylation/Krapcho



SCE).²⁹ Notably, as both 1a and 4-CN pyridine are activated for radical–radical coupling through single-electron reductions, this likely indicates that the DPAIPN photocatalyst is going through two separate redox cycles, both of which are initiated by reductive quenching by NEt₃. This observation is corroborated by a lack of complete conversion of 1a when fewer than 2 equivalents of NEt₃ were employed (see [Supporting Information](#) for details).

To determine if this process proceeds through a closed photoredox cycle, we measured the quantum yield (Φ = 0.28).³⁰ To verify that nonproductive relaxation pathways such as phosphorescence or fluorescence were not impacting the observed quantum yield measurement, we conducted simple quenching experiments. Briefly, DPAIPN was irradiated and the absolute fluorescence was measured. Subsequently, 83 equiv of NEt₃ was added to a solution of DPAIPN (reflective of equivalents of NEt₃ under standard reaction conditions), irradiated, and the absolute fluorescence was measured. Minimal fluorescence was observed, indicating near-complete quenching of the DPAIPN photocatalyst by NEt₃ (quenching fraction, Q = 0.94, Scheme 5B). Based on our proposed mechanism that outlines the need for two photons required for one molecule of the product being formed, an expected quantum yield of <0.5 would be indicative of the reaction potentially proceeding through a closed catalytic cycle. To evaluate if isotope effects impacted the rate of 1a-radical formation, we synthesized 1a-d₁ and compared the rate of consumption of 1a to 1a-d₁ using initial rates. A slight preference for 1a-d₁ was measured (isolated: k_H/k_D = 0.92 \pm 0.03, competition k_H/k_D = 0.88 \pm 0.03). This rate of consumption is likely due to the change in hybridization from sp² to sp³ upon reduction. Because of the in-plane bend of an sp² carbon being much stiffer than the out-of-plane bend (where the in-plane and out-of-plane bends for sp³ carbons are degenerate in energy), this results in a significant difference in the zero-point energy of the two species, producing an inverse secondary kinetic isotope effect (Scheme 5C).³¹

Scheme 5. Mechanistic Experiments




As aforementioned, 1,4-DCB is unreactive under these reaction conditions, which was surprising because of the potential difference between 1,4-DCB and 4-CN pyridine ($E_{1/2}$ red = -1.67 V vs SCE for 1,4-DCB and $E_{1/2}$ red = -1.87 V vs

SCE for 4-CN-pyridine).³² However, it is noteworthy that NHET_3^+ , which is likely being formed under the reaction conditions, ($\text{pK}_a = 9.00$ in DMSO)³³ is sufficiently acidic to protonate 4-CN-pyridine, therefore allowing for a shift in the reduction potential through Brønsted acid activation and resulting in SET instead occurring on the 4-CN-pyridinium ion.³⁴ To investigate this, we conducted cyclic voltammetry (CV) experiments with stoichiometric concentrations of NHET_3Cl with 4-CN-pyridine, and found there was a notable shift in reduction potential for the 4-CN-pyridinium species ($E_{1/2}$ red = -1.51 V), allowing for SET from the DPAIPN radical anion to be slightly exergonic. Contrastingly, 1,4-DCB showed minimal change in reduction potential upon titrating NHET_3Cl ($E_{1/2}$ red = -1.60 V vs SCE) which would result in SET from the DPAIPN radical anion being endergonic, thereby rationalizing the difference in reactivity observed (Scheme 5D). No shift in reduction potential was observed with either substrate upon addition of NEt_3 (see Supporting Information for details). These Brønsted acid activation observations were corroborated by UV-vis studies, where 4-CN-pyridine demonstrates both a noticeable change in the absorption profile as well as an increase in absorption based on stoichiometric addition of NHET_3Cl throughout the 250–300 nm range. However, 1,4-DCB shows a minimal change in the UV absorption profile.³⁵ No UV-vis change is observed with either substrate upon addition of NEt_3 (see Supporting Information for details). In addition, UV-vis studies with **1a** demonstrated no shift upon addition of NEt_3 , but both a change in the absorption profile and an absorbance increase with NHET_3Cl throughout the 250–320 nm range.

A reasonable reaction pathway that accounts for the observed data above begins with irradiation with visible light that results in the formation of the excited DPAIPN photocatalyst, a capable oxidant (Scheme 6). The reductive quenching of the excited state of DPAIPN [$E_{1/2}$ (DPAIPN*/

Scheme 6. Proposed Reaction Pathway



DPAIPN^{•-}) = 1.10 V vs SCE]³⁶ by SET with NEt₃ ($E_{1/2}$ ox = 0.83 V vs SCE) provides a strongly reducing DPAIPN radical anion ($E_{1/2}$ red = -1.52 V vs SCE). The NEt₃ radical cation exists in equilibrium via a 1,2 H-shift with the α -amino radical cation and can be deprotonated by an additional molecule of NEt₃ to form the α -amino radical and NHet₃⁺. Either the NEt₃ radical cation or NHet₃⁺ can engage in PCET with **1** to produce the nucleophilic β -radical and regenerate the ground state DPAIPN catalyst.

A second redox cycle in which NEt₃ or the previously generated α -amino radical can reductively quench DPAIPN to provide the DPAIPN radical anion and either the NEt₃ radical cation or the iminium ion. While it is thermodynamically feasible for the α -amino radical to be the active species for reductive quenching of DPAIPN,³⁷ it is unlikely that this occurs primarily because of the higher relative concentration of NEt₃. The resulting DPAIPN radical anion then undergoes SET with the 4-CN pyridinium cation to form the corresponding arene radical, followed by radical–radical cross-coupling to afford the desired reduction product after cyanide anion elimination and deprotonation.

We have developed a photoredox catalytic manifold that generates stabilized radical species from arylidene malonates. This reactive intermediate undergoes radical–radical cross-coupling with cyanoarene derived arene radicals to afford diaryl malonates in excellent yield. This platform sets the stage for further development of β -umpolung reactivity via photoredox catalysis currently underway in our laboratory.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acscatal.9b03608.

Experimental procedures, characterization of products, and spectroscopic data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: scheidt@northwestern.edu.

ORCID

Karl A. Scheidt: 0000-0003-4856-3569

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the National Institute of General Medical Sciences (R01GM073072, R01GM131431 to K.A.S. and T32GM105538 to R.C.B.) for financial support. R.C.B. was supported in part by the Chicago Cancer Baseball Charities at the Lurie Comprehensive Cancer Center of Northwestern University. The authors thank Mark Maskeri for plate visualization assistance and Joshua Zhu for CV measurements.

■ REFERENCES

(1) (a) Noyori, R. Pursuing practical elegance in chemical synthesis. *Chem. Commun.* **2005**, 1807–1811. (b) Hanessian, S. The Enterprise of Synthesis: From Concept to Practice. *J. Org. Chem.* **2012**, *77*, 6657–6688. (2) (a) Seebach, D. Methods of Reactivity Umpolung. *Angew. Chem., Int. Ed.* **1979**, *18*, 239–258. (b) Hoppe, D.; Hense, T. Enantioselective Synthesis with Lithium/(–)-Sparteine Carbanion Pairs. *Angew. Chem., Int. Ed.* **1997**, *36*, 2282–2316. (c) Reissig, H.-U.;

Zimmer, R. Donor–Acceptor-Substituted Cyclopropane Derivatives and Their Application in Organic Synthesis. *Chem. Rev.* **2003**, *103*, 1151–1196. (d) Johnson, J. S. Catalyzed Reactions of Acyl Anion Equivalents. *Angew. Chem., Int. Ed.* **2004**, *43*, 1326–1328. (e) Smith, A. B.; Adams, C. M. Evolution of Dithiane-Based Strategies for the Construction of Architecturally Complex Natural Products. *Acc. Chem. Res.* **2004**, *37*, 365–377. (f) Ballini, R.; Bosica, G.; Fiorini, D.; Palmieri, A.; Petrini, M. Conjugate Additions of Nitroalkanes to Electron-Poor Alkenes: Recent Results. *Chem. Rev.* **2005**, *105*, 933–972. (g) Ye, L.-W.; Zhou, J.; Tang, Y. Phosphine-triggered synthesis of functionalized cyclic compounds. *Chem. Soc. Rev.* **2008**, *37*, 1140–1152. (h) Guo, F.; Clift, M. D.; Thomson, R. J. Oxidative Coupling of Enolates, Enol Silanes, and Enamines: Methods and Natural Product Synthesis. *Eur. J. Org. Chem.* **2012**, *2012*, 4881–4896.

(3) (a) Enders, D.; Niemeier, O.; Henseler, A. Organocatalysis by N-Heterocyclic Carbenes. *Chem. Rev.* **2007**, *107*, 5606–5655. (b) Nair, V.; Vellalath, S.; Babu, B. P. Recent advances in carbon–carbon bond-forming reactions involving homoenolates generated by NHC catalysis. *Chem. Soc. Rev.* **2008**, *37*, 2691–2698. (c) Nair, V.; Menon, R. S.; Biju, A. T.; Sinu, C. R.; Paul, R. R.; Jose, A.; Sreekumar, V. Employing homoenolates generated by NHC catalysis in carbon–carbon bond-forming reactions: state of the art. *Chem. Soc. Rev.* **2011**, *40*, 5336–5346. (d) Bugaut, X.; Glorius, F. Organocatalytic umpolung: N-heterocyclic carbenes and beyond. *Chem. Soc. Rev.* **2012**, *41*, 3511–3522. (e) Flanagan, D. M.; Romanov-Mikhailidis, F.; White, N. A.; Rovis, T. Organocatalytic Reactions Enabled by N-Heterocyclic Carbenes. *Chem. Rev.* **2015**, *115*, 9307–9387. (f) Menon, R. S.; Biju, A. T.; Nair, V. Recent advances in employing homoenolates generated by N-heterocyclic carbene (NHC) catalysis in carbon–carbon bond-forming reactions. *Chem. Soc. Rev.* **2015**, *44*, 5040–5052.

(4) (a) Terrett, J. A.; Clift, M. D.; MacMillan, D. W. C. Direct β -Alkylation of Aldehydes via Photoredox Organocatalysis. *J. Am. Chem. Soc.* **2014**, *136*, 6858–6861. (b) Jeffrey, J. L.; Petronijević, F. R.; MacMillan, D. W. C. Selective Radical–Radical Cross-Couplings: Design of a Formal β -Mannich Reaction. *J. Am. Chem. Soc.* **2015**, *137*, 8404–8407. (c) Fava, E.; Millet, A.; Nakajima, M.; Loescher, S.; Rueping, M. Reductive Umpolung of Carbonyl Derivatives with Visible-Light Photoredox Catalysis: Direct Access to Vicinal Diamines and Amino Alcohols via α -Amino Radicals and Ketyl Radicals. *Angew. Chem., Int. Ed.* **2016**, *55*, 6776–6779. (d) Fuentes de Arriba, A. L.; Urbitsch, F.; Dixon, D. J. Umpolung synthesis of branched α -functionalized amines from imines via photocatalytic three-component reductive coupling reactions. *Chem. Commun.* **2016**, *52*, 14434–14437. (e) Wang, R.; Ma, M.; Gong, X.; Panetti, G. B.; Fan, X.; Walsh, P. J. Visible-Light-Mediated Umpolung Reactivity of Imines: Ketimine Reductions with Cy2NMe and Water. *Org. Lett.* **2018**, *20*, 2433–2436. (f) Xu, W.; Ma, J.; Yuan, X.-A.; Dai, J.; Xie, J.; Zhu, C. Synergistic Catalysis for the Umpolung Trifluoromethylthiolation of Tertiary Ethers. *Angew. Chem., Int. Ed.* **2018**, *57*, 10357–10361.

(5) (a) Tarantino, K. T.; Liu, P.; Knowles, R. R. Catalytic Ketyl-Olefin Cyclizations Enabled by Proton-Coupled Electron Transfer. *J. Am. Chem. Soc.* **2013**, *135*, 10022–10025. (b) Knowles, R.; Yayla, H. Proton-Coupled Electron Transfer in Organic Synthesis: Novel Homolytic Bond Activations and Catalytic Asymmetric Reactions with Free Radicals. *Synlett* **2014**, *25*, 2819–2826. (c) Miller, D. C.; Tarantino, K. T.; Knowles, R. R. Proton-Coupled Electron Transfer in Organic Synthesis: Fundamentals, Applications, and Opportunities. *Top. Curr. Chem.* **2016**, *374*, 30. (d) Gentry, E. C.; Knowles, R. R. Synthetic Applications of Proton-Coupled Electron Transfer. *Acc. Chem. Res.* **2016**, *49*, 1546–1556.

(6) (a) Ischay, M. A.; Anzovino, M. E.; Du, J.; Yoon, T. P. Efficient Visible Light Photocatalysis of [2+2] Enone Cycloadditions. *J. Am. Chem. Soc.* **2008**, *130*, 12886–12887. (b) Du, J.; Yoon, T. P. Crossed Intermolecular [2+2] Cycloadditions of Acyclic Enones via Visible Light Photocatalysis. *J. Am. Chem. Soc.* **2009**, *131*, 14604–14605. (c) Lu, Z.; Shen, M.; Yoon, T. P. [3+2] Cycloadditions of Aryl Cyclopropyl Ketones by Visible Light Photocatalysis. *J. Am. Chem. Soc.* **2011**, *133*, 1162–1164. (d) Yoon, T. P. Photochemical

Stereocontrol Using Tandem Photoredox-Chiral Lewis Acid Catalysis. *Acc. Chem. Res.* **2016**, *49*, 2307–2315. (e) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* **2016**, *116*, 10035–10074.

(7) (a) Tan, Y.; Yuan, W.; Gong, L.; Meggers, E. Aerobic Asymmetric Dehydrogenative Cross-Coupling between Two C sp³–H Groups Catalyzed by a Chiral-at-Metal Rhodium Complex. *Angew. Chem., Int. Ed.* **2015**, *54*, 13045–13048. (b) Huo, H.; Harms, K.; Meggers, E. Catalytic, Enantioselective Addition of Alkyl Radicals to Alkenes via Visible-Light-Activated Photoredox Catalysis with a Chiral Rhodium Complex. *J. Am. Chem. Soc.* **2016**, *138*, 6936–6939. (c) Zhou, Z.; Li, Y.; Han, B.; Gong, L.; Meggers, E. Enantioselective catalytic β -amination through proton-coupled electron transfer followed by stereocontrolled radical–radical coupling. *Chem. Sci.* **2017**, *8*, 5757–5763. (d) Zhang, L.; Meggers, E. Steering Asymmetric Lewis Acid Catalysis Exclusively with Octahedral Metal-Centered Chirality. *Acc. Chem. Res.* **2017**, *50*, 320–330. (e) de Assis, F. F.; Huang, X.; Akiyama, M.; Pilli, R. A.; Meggers, E. Visible-Light-Activated Catalytic Enantioselective β -Alkylation of α,β -Unsaturated 2-Acyl Imidazoles Using Hantzsch Esters as Radical Reservoirs. *J. Org. Chem.* **2018**, *83*, 10922–10932. (f) Huang, X.; Meggers, E. Asymmetric Photocatalysis with Bis-cyclometalated Rhodium Complexes. *Acc. Chem. Res.* **2019**, *52*, 833–847.

(8) (a) Dell'Amico, L.; Fernández-Alvarez, V. M.; Maseras, F.; Melchiorre, P. Light-Driven Enantioselective Organocatalytic β -Benzoylation of Enals. *Angew. Chem., Int. Ed.* **2017**, *56*, 3304–3308. (b) Silvi, M.; Verrier, C.; Rey, Y. P.; Buzzetti, L.; Melchiorre, P. Visible-light excitation of iminium ions enables the enantioselective catalytic β -alkylation of enals. *Nat. Chem.* **2017**, *9*, 868. (c) Bonilla, P.; Rey, Y. P.; Holden, C. M.; Melchiorre, P. Photo-Organocatalytic Enantioselective Radical Cascade Reactions of Unactivated Olefins. *Angew. Chem., Int. Ed.* **2018**, *57*, 12819–12823. (d) Mazzearella, D.; Crisenza, G. E. M.; Melchiorre, P. Asymmetric Photocatalytic C–H Functionalization of Toluene and Derivatives. *J. Am. Chem. Soc.* **2018**, *140*, 8439–8443. (e) Verrier, C.; Alandini, N.; Pezzetta, C.; Moliterno, M.; Buzzetti, L.; Hepburn, H. B.; Vega-Peñalosa, A.; Silvi, M.; Melchiorre, P. Direct Stereoselective Installation of Alkyl Fragments at the β -Carbon of Enals via Excited Iminium Ion Catalysis. *ACS Catal.* **2018**, *8*, 1062–1066.

(9) (a) McDonald, B. R.; Scheidt, K. A. Intermolecular Reductive Couplings of Arylidene Malonates via Lewis Acid/Photoredox Cooperative Catalysis. *Org. Lett.* **2018**, *20*, 6877–6881. (b) Betori, R. C.; McDonald, B. R.; Scheidt, K. A. Reductive annulations of arylidene malonates with unsaturated electrophiles using photoredox/Lewis acid cooperative catalysis. *Chem. Sci.* **2019**, *10*, 3353–3359.

(10) (a) Zhao, G.; Yang, C.; Guo, L.; Sun, H.; Lin, R.; Xia, W. Reactivity Insight into Reductive Coupling and Aldol Cyclization of Chalcones by Visible Light Photocatalysis. *J. Org. Chem.* **2012**, *77*, 6302–6306. (b) Tyson, E. L.; Farney, E. P.; Yoon, T. P. Photocatalytic [2 + 2] Cycloadditions of Enones with Cleavable Redox Auxiliaries. *Org. Lett.* **2012**, *14*, 1110–1113. (c) Hurtley, A. E.; Cismesia, M. A.; Ischay, M. A.; Yoon, T. P. Visible light photocatalysis of radical anion hetero-Diels–Alder cycloadditions. *Tetrahedron* **2011**, *67*, 4442–4448. (d) Du, J.; Espelt, L. R.; Guzei, I. A.; Yoon, T. P. Photocatalytic reductive cyclizations of enones: Divergent reactivity of photogenerated radical and radical anion intermediates. *Chem. Sci.* **2011**, *2*, 2115–2119. (e) Ischay, M. A.; Lu, Z.; Yoon, T. P. [2+2] Cycloadditions by Oxidative Visible Light Photocatalysis. *J. Am. Chem. Soc.* **2010**, *132*, 8572–8574.

(11) Gang, D.; Jun, Y.; Xiao-ping, Y.; Hui-jun, X. Hydride transfer in the reduction of substituted benzylidene malonic diesters by coenzyme NAD(P)H model. *Tetrahedron* **1990**, *46*, 5967–5974.

(12) McNally, A.; Prier, C. K.; MacMillan, D. W. C. Discovery of an α -Amino C–H Arylation Reaction Using the Strategy of Accelerated Serendipity. *Science* **2011**, *334*, 1114.

(13) Cuthbertson, J. D.; MacMillan, D. W. C. The direct arylation of allylic sp³ C–H bonds via organic and photoredox catalysis. *Nature* **2015**, *519*, 74.

(14) Petronijević, F. R.; Nappi, M.; MacMillan, D. W. C. Direct β -Functionalization of Cyclic Ketones with Aryl Ketones via the Merger of Photoredox and Organocatalysis. *J. Am. Chem. Soc.* **2013**, *135*, 18323–18326.

(15) (a) Chen, M.; Zhao, X.; Yang, C.; Xia, W. Visible-Light-Triggered Directly Reductive Arylation of Carbonyl/Imine Derivatives through Photocatalytic PCET. *Org. Lett.* **2017**, *19*, 3807–3810. (b) Zuo, Z.; MacMillan, D. W. C. Decarboxylative Arylation of α -Amino Acids via Photoredox Catalysis: A One-Step Conversion of Biomass to Drug Pharmacophore. *J. Am. Chem. Soc.* **2014**, *136*, 5257–5260. (c) Qvortrup, K.; Rankic, D. A.; MacMillan, D. W. C. A General Strategy for Organocatalytic Activation of C–H Bonds via Photoredox Catalysis: Direct Arylation of Benzylic Ethers. *J. Am. Chem. Soc.* **2014**, *136*, 626–629.

(16) (a) Krska, S. W.; DiRocco, D. A.; Dreher, S. D.; Shevlin, M. The Evolution of Chemical High-Throughput Experimentation To Address Challenging Problems in Pharmaceutical Synthesis. *Acc. Chem. Res.* **2017**, *50*, 2976–2985. (b) Shevlin, M. Practical High-Throughput Experimentation for Chemists. *ACS Med. Chem. Lett.* **2017**, *8*, 601–607. (c) Isbrandt, E. S.; Sullivan, R. J.; Newman, S. G. High Throughput Strategies for the Discovery and Optimization of Catalytic Reactions. *Angew. Chem., Int. Ed.* **2019**, *58*, 7180–7191. (d) Mennen, S. M.; Alhambra, C.; Allen, C. L.; Barberis, M.; Berritt, S.; Brandt, T. A.; Campbell, A. D.; Castanon, J.; Cherney, A. H.; Christensen, M.; Damon, D. B.; Eugenio de Diego, J.; García-Cerrada, S.; García-Losada, P.; Haro, R.; Janey, J. M.; Leitch, D. C.; Li, L.; Liu, F.; Lobben, P. C.; MacMillan, D. W. C.; Magano, J.; McInturff, E. L.; Monfette, S.; Post, R. J.; Schultz, D. M.; Sitter, B. J.; Stevens, J. M.; Strambeanu, I. I.; Twilton, J.; Wang, K.; Zajac, M. A. The Evolution of High-Throughput Experimentation in Pharmaceutical Development, and Perspectives on the Future. *Org. Process Res. Dev.* **2019**, *23*, 1213–1242.

(17) Humbel, S.; Côte, I.; Hoffmann, N.; Bouquant, J. Three-Electron Binding between Carbonyl-like Compounds and Ammonia Radical Cation. Comparison with the Hydrogen Bonded Complex. *J. Am. Chem. Soc.* **1999**, *121*, 5507–5512.

(18) Akalay, D.; Dürner, G.; Bats, J. W.; Bolte, M.; Göbel, M. W. Synthesis of C2-Symmetric Bisamidines: A New Type of Chiral Metal-Free Lewis Acid Analogue Interacting with Carbonyl Groups. *J. Org. Chem.* **2007**, *72*, 5618–5624.

(19) (a) Nakajima, M.; Fava, E.; Loescher, S.; Jiang, Z.; Rueping, M. Photoredox-Catalyzed Reductive Coupling of Aldehydes, Ketones, and Imines with Visible Light. *Angew. Chem., Int. Ed.* **2015**, *54*, 8828–8832. (b) Fava, E.; Nakajima, M.; Nguyen, A. L. P.; Rueping, M. Photoredox-Catalyzed Ketyl–Olefin Coupling for the Synthesis of Substituted Chromanols. *J. Org. Chem.* **2016**, *81*, 6959–6964.

(20) Robinson, J.; Osteryoung, R. A. An investigation into the electrochemical oxidation of some aromatic amines in the room-temperature molten salt system aluminum chloride-butylpyridinium chloride. *J. Am. Chem. Soc.* **1980**, *102*, 4415–4420.

(21) Pitzer, L.; Schäfers, F.; Glorius, F. Rapid Assessment of the Reaction-Condition-Based Sensitivity of Chemical Transformations. *Angew. Chem., Int. Ed.* **2019**, *58*, 8572–8576.

(22) (a) Sörgel, S.; Tokunaga, N.; Sasaki, K.; Okamoto, K.; Hayashi, T. Rhodium/Chiral Diene-Catalyzed Asymmetric 1,4-Addition of Arylboronic Acids to Arylmethylene Cyanoacetates. *Org. Lett.* **2008**, *10*, 589–592. (b) Takatsu, K.; Shintani, R.; Hayashi, T. Copper-Catalyzed 1,4-Addition of Organoboronates to Alkylidene Cyanoacetates: Mechanistic Insight and Application to Asymmetric Catalysis. *Angew. Chem., Int. Ed.* **2011**, *50*, 5548–5552.

(23) (a) Macdonald, D. I.; Durst, T. A synthesis of trans-2-arylbenzocyclobuten-1-ols. *Tetrahedron Lett.* **1986**, *27*, 2235–2238. (b) Cahiez, G.; Venegas, P.; Tucker, C. E.; Majid, T. N.; Knochel, P. Addition of polyfunctional and pure (E or Z) alkenylcopper and arylcopper compounds to alkylidenemalonates. *J. Chem. Soc., Chem. Commun.* **1992**, 1406–1408. (c) Almansa, C.; Gómez, L. A.; Cavalcanti, F. L.; de Arriba, A. F.; Carceller, R.; García-Rafanell, J.; Forn, J. Diphenylpropionic Acids as New AT1 Selective Angiotensin II Antagonists. *J. Med. Chem.* **1996**, *39*, 2197–2206. (d) Curran, D.

- P.; Gabarda, A. E. Formation of cyclopropanes by homolytic substitution reactions of 3-iodopropyl radicals: Preparative and rate studies. *Tetrahedron* **1999**, *55*, 3327–3336. (e) Salomone, A.; Capriati, V.; Florio, S.; Luisi, R. Michael Addition of Ortho-Lithiated Aryloxiranes to α,β -Unsaturated Malonates: Synthesis of Tetrahydroindolenofuranones. *Org. Lett.* **2008**, *10*, 1947–1950. (f) De La Rosa, M. M. W.; Samano, V. Modulators of indoleamine 2,3-dioxygenase. Eur. Pat. Appl. 3558966, June 28, 2018. (g) Feenstra, R.; Stoit, A.; Terpstra, J.; Pras-Raves, M.; McCreary, A.; Van Vliet, B.; Hesselink, M.; Kruse, C.; Van Scharrenburg, G. Phenylpiperazine derivatives with a combination of partial dopamine-D2 receptor agonism and serotonin reuptake inhibition. U.S. Patent 20,070,072,870 A1, June 08, 2006.
- (24) (a) Hayashi, T.; Yamasaki, K. Rhodium-Catalyzed Asymmetric 1,4-Addition and Its Related Asymmetric Reactions. *Chem. Rev.* **2003**, *103*, 2829–2844. (b) Jerphagnon, T.; Pizzuti, M. G.; Minnaard, A. J.; Feringa, B. L. Recent advances in enantioselective copper-catalyzed 1,4-addition. *Chem. Soc. Rev.* **2009**, *38*, 1039–1075. (c) Edwards, H. J.; Hargrave, J. D.; Penrose, S. D.; Frost, C. G. Synthetic applications of rhodium catalyzed conjugate addition. *Chem. Soc. Rev.* **2010**, *39*, 2093–2105. (d) Heravi, M. M.; Dehghani, M.; Zadsirjan, V. Rh-catalyzed asymmetric 1,4-addition reactions to α,β -unsaturated carbonyl and related compounds: an update. *Tetrahedron: Asymmetry* **2016**, *27*, 513–588.
- (25) (a) Zhou, J.; Ye, M.-C.; Huang, Z.-Z.; Tang, Y. Controllable Enantioselective Friedel–Crafts Reaction between Indoles and Alkylidene Malonates Catalyzed by Pseudo-C3-Symmetric Trisoxazoline Copper(II) Complexes. *J. Org. Chem.* **2004**, *69*, 1309–1320. (b) Sun, Y.-J.; Li, N.; Zheng, Z.-B.; Liu, L.; Yu, Y.-B.; Qin, Z.-H.; Fu, B. Highly Enantioselective Friedel–Crafts Reaction of Indole with Alkylidenemalonates Catalyzed by Heteroarylidenemalonate-Derived Bis(oxazoline) Copper(II) Complexes. *Adv. Synth. Catal.* **2009**, *351*, 3113–3117. (c) Sun, X.-L.; Zhou, Y.-Y.; Zhu, B.-H.; Zheng, J.-C.; Zhou, J.-L.; Tang, Y. Modification of Pseudo-C3-Symmetric Trisoxazoline and Its Application to the Friedel–Crafts Alkylation of Indoles and Pyrrole with Alkylidene Malonates. *Synlett* **2011**, 935–938. (d) Liu, L.; Li, J.; Wang, M.; Du, F.; Qin, Z.; Fu, B. Synthesis of heteroarylidenemalonate derived bis(thiazolines) and their application in the catalyzed Friedel–Crafts reaction. *Tetrahedron: Asymmetry* **2011**, *22*, 550–557.
- (26) (a) Ramanathan, R.; Daniel, P. S.; Deborah, J.; Edmund, B. J.; Brian, N. A.; Gale, W. M.; David, D. A.; Rong, X.; Edwards, D. R.; Joseph, S. Ultraviolet Light Absorbers. WO 03016292 A1, 2001. (b) Kuwano, R.; Kusano, H. Palladium-catalyzed Nucleophilic Substitution of Diarylmethyl Carbonates with Malonate Carbanions. *Chem. Lett.* **2007**, *36*, 528–529. (c) Baba, A.; Babu, S.; Yasuda, M.; Tsukahara, Y.; Yamauchi, T.; Wada, Y. Microwave-Irradiated Transition-Metal Catalysis: Rapid and Efficient Dehydrative Carbon–Carbon Coupling of Alcohols with Active Methylenes. *Synthesis* **2008**, 1717–1724. (d) Onaka, M.; Wang, J.; Masui, Y. Efficient Nucleophilic Substitution of α -Aryl Alcohols with 1,3-Dicarbonyl Compounds Catalyzed by Tin Ion-Exchanged Montmorillonite. *Synlett* **2010**, 2493–2497. (e) Bonda, C. A.; Hu, S.; Zhang, Q. J.; Zhan, Z. Compositions, Apparatus, Systems, and Methods for Resolving Electronic Excited States. WO 2013US54408, 2013. (f) Nitti, A.; Villafiorita-Montealeone, F.; Pacini, A.; Botta, C.; Virgili, T.; Forni, A.; Cariati, E.; Boiocchi, M.; Pasini, D. Structure–activity relationship for the solid state emission of a new family of “push–pull” π -extended chromophores. *Faraday Discuss.* **2017**, *196*, 143–161. (g) Konishi, A.; Okada, Y.; Nakano, M.; Sugisaki, K.; Sato, K.; Takui, T.; Yasuda, M. Synthesis and Characterization of Dibenzo[a,f]pentalene: Harmonization of the Antiaromatic and Singlet Biradical Character. *J. Am. Chem. Soc.* **2017**, *139*, 15284–15287. (h) Konishi, A.; Okada, Y.; Kishi, R.; Nakano, M.; Yasuda, M. Enhancement of Antiaromatic Character via Additional Benzoannulation into Dibenzo[a,f]pentalene: Syntheses and Properties of Benzo[a]naphtho[2,1-f]pentalene and Dinaphtho[2,1-a,f]pentalene. *J. Am. Chem. Soc.* **2019**, *141*, 560–571.
- (27) (a) Yang, K.; Song, Q. Fe-Catalyzed Double Cross-Dehydrogenative Coupling of 1,3-Dicarbonyl Compounds and Arylmethanes. *Org. Lett.* **2015**, *17*, 548–551. (b) Li, Z.; Cao, L.; Li, C.-J. FeCl₂-Catalyzed Selective C–C Bond Formation by Oxidative Activation of a Benzylic C–H Bond. *Angew. Chem., Int. Ed.* **2007**, *46*, 6505–6507. (c) Shi, J.-L.; Luo, Q.; Yu, W.; Wang, B.; Shi, Z.-J.; Wang, J. Fe(II)-Catalyzed alkenylation of benzylic C–H bonds with diazo compounds. *Chem. Commun.* **2019**, *55*, 4047–4050.
- (28) Buzzetti, L.; Crisenza, G. E. M.; Melchiorre, P. Mechanistic Studies in Photocatalysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 3730–3747.
- (29) Nicewicz, D.; Roth, H.; Romero, N. Experimental and Calculated Electrochemical Potentials of Common Organic Molecules for Applications to Single-Electron Redox Chemistry. *Synlett* **2015**, *27*, 714–723.
- (30) Cismesia, M. A.; Yoon, T. P. Characterizing chain processes in visible light photoredox catalysis. *Chem. Sci.* **2015**, *6*, 5426–5434.
- (31) Meyer, M. P. New Applications of Isotope Effects in the Determination of Organic Reaction Mechanisms. In *Advances in Physical Organic Chemistry*; Williams, I. H., Williams, N. H., Eds.; Academic Press, 2012; Vol. 46, Chapter 2, pp 57–120.
- (32) McDevitt, P.; Vittimberga, B. M. The electron transfer reactions of cyano substituted pyridines and quinolines with thermally generated diphenyl ketyl. *J. Heterocycl. Chem.* **1990**, *27*, 1903–1908.
- (33) (a) Bordwell, F. G. Equilibrium acidities in dimethyl sulfoxide solution. *Acc. Chem. Res.* **1988**, *21*, 456–463. (b) Kolthoff, I. M.; Chantooni, M. K.; Bhowmik, S. Dissociation constants of uncharged and monovalent cation acids in dimethyl sulfoxide. *J. Am. Chem. Soc.* **1968**, *90*, 23–28. (c) Crampton, M. R.; Robotham, I. A. Acidities of Some Substituted Ammonium Ions in Dimethyl Sulfoxide. *J. Chem. Res., Synop.* **1997**, 22–23.
- (34) (a) Kitamura, N.; Nambu, Y.; Endo, T. Redox behavior of pyridinium salts and their polymers having radical stabilizing groups. *J. Polym. Sci., Part A: Polym. Chem.* **1990**, *28*, 3137–3143. (b) Yan, Y.; Zeitler, E. L.; Gu, J.; Hu, Y.; Bocarsly, A. B. Electrochemistry of Aqueous Pyridinium: Exploration of a Key Aspect of Electrocatalytic Reduction of CO₂ to Methanol. *J. Am. Chem. Soc.* **2013**, *135*, 14020–14023.
- (35) Tang, Z.; Wu, X.; Guo, B.; Zhang, L.; Jia, D. Preparation of butadiene–styrene–vinyl pyridine rubber–graphene oxide hybrids through co-coagulation process and in situ interface tailoring. *J. Mater. Chem.* **2012**, *22*, 7492–7501.
- (36) Luo, J.; Zhang, J. Donor–Acceptor Fluorophores for Visible-Light-Promoted Organic Synthesis: Photoredox/Ni Dual Catalytic C(sp³)–C(sp²) Cross-Coupling. *ACS Catal.* **2016**, *6*, 873–877.
- (37) (a) Wayner, D. D. M.; Dannenberg, J. J.; Griller, D. Oxidation potentials of α -aminoalkyl radicals: bond dissociation energies for related radical cations. *Chem. Phys. Lett.* **1986**, *131*, 189–191. (b) Ames, J. R.; Brandänge, S.; Rodriguez, B.; Castagnoli, N.; Ryan, M. D.; Kovacic, P. Cyclic voltammetry with cyclic iminium ions: Implications for charge transfer with biomolecules (metabolites of nicotine, phencyclidine, and spermine). *Bioorg. Chem.* **1986**, *14*, 228–241. (c) Nakajima, K.; Miyake, Y.; Nishibayashi, Y. Synthetic Utilization of α -Aminoalkyl Radicals and Related Species in Visible Light Photoredox Catalysis. *Acc. Chem. Res.* **2016**, *49*, 1946–1956.