Mechanistic Studies on the Rearrangement of 1-Alkenyl-2alkynylcyclopropanes: From Allylic Gold(I) Cations to Stable Carbocations**

Eva Tudela, Jairo González, Rubén Vicente, Javier Santamaría, Miguel A. Rodríguez, and Alfredo Ballesteros*

Abstract: An allylic gold(I) cation, proposed as key intermediate in the gold-promoted rearrangement of 1,5-enynes bearing a fixed conformation, has been detected and characterized by NMR spectroscopy. Moreover, its participation in the overall transformation was confirmed. Computational studies indicate that the gold-catalyzed transformation occurs through an uncommon rearrangement. Additionally, this study led us to isolate and characterize a stable homoantiaromatic carbocation.

Gold-catalyzed cycloisomerization of 1,*n*-enynes represents a paradigm in gold catalysis.^[1] From these substrates, a great variety of products with an increased molecular complexity can be easily accessed by skeletal rearrangements. These transformations involve the nucleophilic attack of the alkene onto the gold-activated alkyne, thus leading to an intermediate which can be depicted with multiple resonance structures (Scheme 1 a).^[2] Thus, the reaction outcome has been related to the contribution of these resonance structures, which can be strongly influenced by the enyne structure, catalyst, or reaction conditions. Furthermore, the high reactivity of the proposed intermediates makes their trapping a remarkably challenging task. In this sense, NMR studies have shown that gold(I) coordinates to both the alkene and alkyne to give an equilibrium mixture (Scheme 1b),^[3] while unique experimental evidence of a gold(I) intermediate has been recently reported by Widenhoefer and co-workers in the cycloisomerization of a 1.6-envne (Scheme 1 c).^[4]

We have recently studied the reactivity of a specific class of 1,5-enynes, that is, the alkynylcyclopropanes 1, which

[*] Dr. E. Tudela, J. González, Dr. R. Vicente, Dr. J. Santamaría, Prof. Dr. A. Ballesteros Instituto de Química Organometálica "Enrique Moles" and Departamento de Química Orgánica e Inorgánica, Universidad de Oviedo, c/Julián Clavería 8, 33007, Oviedo (Spain) E-mail: abg@uniovi.es Prof. Dr. M. A. Rodríguez

Departamento de Química, Centro de Investigación en Síntesis Química, Universidad de la Rioja, 26006 Logroño (Spain)

[**] We are grateful to the MINECO (Spain) (Projects CTQ-2010-20517-C02, CTQ2013-41511-P and CTQ2011-24800) for financial support. R.V. is a Ramón y Cajal fellow. We thank Dr. I. Merino and Dr. E. Rubio for the assistance in NMR experiments, Dr. A. L. Suárez-Sobrino for the X-Ray analysis, and Prof. Dr. J. Barluenga and Prof. Dr. M. Tomás for their suggestions.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201407055.



Scheme 1. a) Resonance structures in gold(I)-promoted enyne cycloisomerizations. b) Coordination of gold(I) to 1,*n*-enynes. c) NMR evidence of gold(I) intermediates in cycloisomerizations.



Scheme 2. Gold(I)-catalyzed rearrangement of the enyne **1** into the cyclohexadienes **2** and **3** and previously proposed mechanism.

undergoes a gold-catalyzed rearrangement to the alkynylcyclohexadienes **2** and **3** (Scheme 2).^[5] At that stage, a tentative pathway for the formation of **2** and **3** involving a 3-*exo-dig* attack via the intermediates **I** and **I'**, respectively, was suggested. The formation of **2** and **3** was justified assuming an isomerization of **1** through a reversible 6-*endo-dig* attack of the olefin involving intermediate **II**, a formal allylic gold(I) cation. To gain further proof on the existence of intermediates such as **II**, as well as to unravel the reaction pathway, we carried out experimental and computational studies, the results of which are described herein.

Preliminary indications on the participation of **II** were gained by its trapping with alcohols as nucleophiles,^[5] and led to the isolation of the compounds **4** and **5** (see Scheme 3)

Angew. Chem. Int. Ed. 2014, 53, 1-5

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angewandte Communications



Scheme 3. Gold-catalyzed selective reactions of 1a with MeOH(D). [a] exo/endo $\approx 2:1$.

depending to the size of the alcohol.^[6] Moreover, we have also found that the outcome in the reaction of the alkynylcyclopropane **1a** and MeOH can be correlated with the percent buried volume ((V_{bur})) of the gold complexes (Scheme 3).^[7] Thus, while the use of $[Au(PPh_3)]^+$ led to **4a**, the bulkier catalyst $[Au(JohnPhos)]^+$ gave rise to **5a**. Besides, the NHCstabilized gold catalyst $[Au(IPr)]^+$ selectively afforded **4a**. Experiments using MeOD gave rise to either [D]-**4a** or [D]-**5a**, thus providing additional experimental support. These experiments clearly point out the existence of the allylic gold(I) cation structure **II**.

Then, we attempted to detect **II** (Scheme 4). Treatment of **1b** with stoichiometric amounts of $[Au(PPh)_3Cl]/AgOTf$ in



Scheme 4. Gold-mediated conversion of **1b** into **6a–c** and selected NMR data. The chemical shifts are given in ppm. [a] Mixture of isomers. $PMP=4-MeOC_6H_4$, Tf=trifluoromethanesulfonyl.

CDCl₃ at -50 °C immediately produced a bright blue solution, thus indicating the formation of a new species. The NMR analysis evidenced the formation of the new symmetrical complex **6a**. Thus, C3 was identified at $\delta = 168.5$ ppm (²J_{P-C} = 108 Hz), while signals corresponding to C2–C4 were found at $\delta = 202.0$ ppm (selected signals shown in Scheme 4). Unfortunately, **6a** proved unstable and decomposed gradually during the measurement. We then accomplished the study

using [Au(IPr)][NTf₂], and were also able to generate the gold complex 6b. The compound 6b showed a superior stability and it could be characterized by one- and twodimensional NMR spectroscopy (see the Supporting Information). Signals corresponding to C3 and C2-C4 were located at $\delta = 165.7$ and 201.8 ppm, respectively. Moreover, the complex 6c, derived from the use of [Au(JohnPhos)(MeCN)]- $[SbF_6]$, was also prepared and characterized by NMR spectroscopy. A mixture of two isomers at -20°C was observed, with signals at $\delta = 169.1/168.1$ (C3) and 199.8/ 199.6 ppm (C2-C4) in the ¹³C NMR (³¹P decoupled) spectra.^[8] Chemical shifts of C2, C3, and C4 in complexes 6a-c indicate that the positive charge is distributed along the η^1 -allyl gold moiety.^[9] Besides, the complexes $\mathbf{6a-c}$ display a structure which might be potentially destabilized by π -electron delocalization with certain homoantiaromatic character.^[10] Indeed, the paramagnetic ring current associated to the homoconjugation is reflected in the low-field shifts ($\delta = 15-20$ ppm) of bridged methylene signals observed for C8 in 6a-c with respect to **4b** ($R^1 = R^2 = p$ -MeOC₆H₄; $\delta = 41.8, 42.8$ ppm, *exo/* endo), obtained as described in Scheme 3.^[11]

The participation of the allylic gold(I) cations 6 in the catalytic processes was further proved (Scheme 5). Thus,



Scheme 5. Catalytic competence of gold complex 6c.

treatment of the cyclopropanes 1a and 1b with approximately 2 mol% of 6c in CDCl₃ at 70 °C afforded the expected cyclohexadienes 2a and 2b, respectively. Moreover, when the reaction was performed in the presence of MeOH, the corresponding compounds 5a and 5b were isolated.

The characterization η^1 -allylic gold(I) cationic species (6) constitutes the first experimental evidence on the formation of such complexes. This result is not only relevant to the present transformation. These elusive species have been proposed in other gold(I)-catalyzed reactions but were never characterized.^[12-14]

Although X-ray analysis of the complexes **6** failed because of their decomposition, a new species which was identified as the carbocation **7** was detected in these trials (Scheme 6). Indeed, **7** was obtained by treatment of **1b** with trifluoromethane sulfonic acid (TfOH, 1.0 equiv) in CH₂Cl₂ at -20 °C. Remarkably, TfOH proved unable to catalyze the



Scheme 6. Synthesis of carbocation 7. ($PMP = 4 - MeO - C_6H_4$)

www.angewandte.org

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angew. Chem. Int. Ed. 2014, 53, 1-5

These are not the final page numbers!

rearrangement of **1** into **2**, and **7** could not be converted into **2b** under the many reaction conditions attempted. These results indicated the need of gold to accomplish the transformation of **1** into **2** and **3**. Besides, **7** showed high stability as it could be fully characterized by NMR and X-Ray analysis at room temperature.^[15] Homoconjugation in **7** is also significant considering the low-field shift in signals corresponding to H8-H8' ($\delta = 2.6, 2.9$ ppm).^[16] The compound **7** represents a rare example of a stable homoantiaromatic species.^[9,10]

We also carried out a computational study (B3PW91-D3/ cc-pVDZ-PP//B3PW91/cc-pVDZ-PP level) on the mechanism of the rearrangement of 1c ($R^1 = R^2 = Ph$) into 2c, which is depicted in Figure 1 (see the Supporting Information for



Figure 1. Calculated relative free-energy profile for the rearrangement of 1 c into 2 c. (Carbon numbers were assigned as indicated for complexes 6; numbers within parentheses indicate ΔG in kcal mol⁻¹; [Au] = AuPH₃).

details). Starting from the species 1c-[Au], alkyne coordination to gold facilitates the 6-*endo-dig* π_{C-C} attack to give the transition-state **TS1** (+ 6.2 kcalmol⁻¹). From **TS1**, the intrinsic reaction coordinate (IRC) leads to **TS2** (-0.9 kcalmol⁻¹), which showed an elongation on the C1–C7 bond (1.370 to 1.446 Å) and the imminent formation of the C1–C2 bond (2.217 to 1.590 Å). Since the IRC calculation connects two transition states, there must be a valley-ridge inflection (VRI) or a bifurcation along the IRC.^[17] **TS2** can evolve towards the structures **A** (-13.0 kcalmol⁻¹) or **B** (-17.3 kcalmol⁻¹), which resemble the previously observed complexes **6**. Interestingly, we found an energetically plausible reaction pathway which connects **B** with the final structure **2c**-[Au]. Thus, a rare skeleton rearrangement from **B** to the more stable structure C (-18.4 kcal mol⁻¹) occurs to overcome the highest energy barrier (23.5 kcalmol⁻¹) through **TS3** (+6.2 kcal mol⁻¹).^[18] This step involves a double 1,3-migration as two bonds are being cleaved and two bonds are being created. Finally, the transition-state **TS4** $(-3.8 \text{ kcal mol}^{-1})$ connects **C** with 2c-[Au] (-13.1 kcalmol⁻¹). Thus, this computational study provides a plausible reaction pathway for the conversion of 1 into 2 and 3 through the common intermediate B, whose participation has been experimentally proven by spectroscopic observation. The relative energies found for **B** going to either **TS1** or **TS3** (23.5 kcalmol⁻¹, in both cases) indicate that an equilibrium between C and A is feasible, as it was proposed to explain the formation of isomers.^[5] Conversely, the earlier postulated 3-exo-dig nucleophilic attack (Scheme 2) seems to be less likely as the calculated structure for the corresponding transition state is higher in energy than **TS1** (+17.8 versus +6.2; see the Supporting Information for the complete calculated pathway).

In summary, this study allows us to propose a revised mechanism for the gold-catalyzed rearrangement of envnes (1) into the cyclohexadienes 2 and 3. While examples on the observation of gold intermediates is rare, we were able to detect and, for the first time, characterize by NMR spectroscopy the η^1 -allylic gold(I) cationic intermediates 6. Remarkably, structures like 6 are not only relevant in 1,n-enyne cycloisomerizations. In this sense, η^1 -allylic gold(I) cationic species have been proposed in the racemization of allenes and their further functionalizations, yet they were never experimentally characterized.^[19] Additionally, the computational study provides a reaction pathway connecting $1 \rightarrow 6 \rightarrow 2/3$. This study supports the role of 6 in the overall transformation, which undergoes an unusual rearrangement by a double 1,3migration. Moreover, we were also able to isolate and fully characterize compound 7, a rare example of stable homoantiaromatic carbocation.

Received: July 9, 2014 Published online:

Keywords: allylic compounds · carbocations · gold · reactive intermediates · rearrangements

For reviews on gold-catalyzed cycloisomerizations, see: a) L. Zhang, J. Sun, S. A. Kozmin, Adv. Synth. Catal. 2006, 348, 2271– 2296; b) E. Jiménez-Nuñez, A. M. Echavarren, Chem. Rev. 2008, 108, 3326-3350; c) V. Michelet, P. Y. Toullec, J.-P. Genet, Angew. Chem. Int. Ed. 2008, 47, 4268-4315; Angew. Chem. 2008, 120, 4338-4386; d) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria, A. Simonneau, Chem. Rev. 2011, 111, 1954-1993; For general recent reviews on gold catalysis, see: e) N. Krause, C. Winter, Chem. Rev. 2011, 111, 1994-2009; f) A. Corma, A. Leyva-Pérez, M. J. Sabater, Chem. Rev. 2011, 111, 1657-1712; g) H. Huang, Y. Zhou, H. Liu, Beilstein J. Org. Chem. 2011, 7, 897-936; h) Modern Gold Catalyzed Synthesis (Eds.: A. S. K. Hashmi, F. D. Toste), Wiley-VCH, Weinheim, 2012.

 ^[2] a) A. Fürstner, Acc. Chem. Res. 2014, 47, 925–938; b) C. Obradors, A. M. Echavarren, Chem. Commun. 2014, 50, 16–28; c) C. Obradors, A. M. Echavarren, Acc. Chem. Res. 2014, 47, 902–912.

^{© 2014} Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angewandte Communications

- [3] M. García-Mota, N. Cabello, F. Maseras, A. M. Echavarren, J. Pérez-Ramírez, N. López, *ChemPhysChem* 2008, 9, 1624–1629.
- [4] R. E. M. Brooner, T. J. Brown, R. A. Widenhoefer, Angew. Chem. Int. Ed. 2013, 52, 6259–6261; Angew. Chem. 2013, 125, 6379–6381.
- [5] J. Barluenga, E. Tudela, R. Vicente, A. Ballesteros, M. Tomás, *Angew. Chem. Int. Ed.* 2011, 50, 2107–2110; *Angew. Chem.* 2011, 123, 2155–2158.
- [6] Bicycles like **4a** were obtained when using MeOH, while bulkier alcohols afforded tricycles like **5a**.
- [7] H. Clavier, S. P. Nolan, Chem. Commun. 2010, 46, 841-861.
- [8] The formation of two isomers likely originates from the steric congestion between the allylic cation fragment and the phosphine ligand. Attempts to reach coalescence signals at higher temperatures proved unsuccessful because of the degradation of the complex 6c.
- [9] For a representative example of chemical shifts in the ¹³C NMR spectra of benzylic carbocations, see: H. C. Brown, D. P. Kelly, M. Periasamy, *Proc. Natl. Acad. Sci. USA* **1980**, 77, 6956–6960.
- [10] Homoaromaticity is a term coined by Winstein: a) S. Winstein, J. Am. Chem. Soc. 1959, 81, 6524-6525; For a review focused on this topic, see: b) R. V. Williams, Chem. Rev. 2001, 101, 1185-1204.
- [11] For two examples of homoantiaromatic cations related to the complexes 6, see: a) H. Hart, M. Kuzuya, *J. Am. Chem. Soc.* **1975**, 97, 2459–2468; b) H. Volz, J.-H. Shin, *J. Org. Chem.* **2006**, *71*, 2220–2226.
- [12] For theoretical studies related to allylic gold(I) cations, see:
 a) B. W. Gung, D. T. Craft, L. N. Bailey, K. Kirschbaum, *Chem. Eur. J.* 2010, *16*, 639–644;
 b) D. Benítez, E. Tkatchouk, A. Z. Gonzalez, W. A. Goddard III, F. D. Toste, *Org. Lett.* 2009, *11*, 4798–4801;
 c) B. Trillo, F. López, S. Montserrat, G. Ujaque, L. Castedo, A. Lledós, J. L. Mascareñas, *Chem. Eur. J.* 2009, *15*, 3336–3339;
 d) V. Gandon, G. Lemiere, A. Hours, L. Fenster-

bank, M. Malacria, Angew. Chem. Int. Ed. 2008, 47, 7534–7538; Angew. Chem. 2008, 120, 7644–7648.

- [13] For an example of related gold carbodicarbene complexes, see: A. Fürstner, M. Alcarazo, R. Goddard, C. W. Lehmann, *Angew. Chem. Int. Ed.* 2008, 47, 3210–3214; *Angew. Chem.* 2008, 120, 3254–3258.
- [14] For the isolation of η²-allene-gold complexes, see: a) T. J. Brown, A. Sugie, M. G. Dickens, R. A. Widenhoefer, *Organometallics* 2010, 29, 4207–4209; b) T. J. Brown, A. Sugie, M. G. D. Leed, R. A. Widenhoefer, *Chem. Eur. J.* 2012, *18*, 6959–6971.
- [15] CCDC 1011686 (7) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www. ccdc.cam.ac.uk/data_request/cif.
- [16] Theoretical calculations provides shift values of $\delta = 2.5$ -3.0 ppm for H8,H8' in **7**. In the absence of the double bond, these signals were calculated to appear at $\delta = 1.9$ -2.0 ppm. see the Supporting Information for details.
- [17] a) O. Castaño, L.-M. Frutos, R. Palmeiro, R. Notario, J.-L. Andrés, R. Gomperts, L. Blancafort, M. A. Robb, Angew. Chem. Int. Ed. 2000, 39, 2095–2097; Angew. Chem. 2000, 112, 2168–2170; b) R. Palmeiro, L. M. Frutos, O. Castaño, Int. J. Quantum Chem. 2002, 86, 422–425; For selected examples of gold-catalyzed transformations involving an energetic bifurcation see: c) L. Ye, Y. Wang, D. H. Aue, L. Zhang, J. Am. Chem. Soc. 2012, 134, 31–34; d) M. M. Hansmann, M. Rudolph, F. Rominger, A. S. K. Hashmi, Angew. Chem. Int. Ed. 2013, 52, 2593–2598; Angew. Chem. 2013, 125, 2653–2659.
- [18] For a recent example of a theoretical study on uncommon 1,3migration in gold-catalyzed rearrangements, see: Q. Zhou, Y. Li, *J. Am. Chem. Soc.* 2014, *136*, 1505–1513.
- [19] R. J. Harris, K. Nakafuku, R. A. Widenhoefer, *Chem. Eur. J.* 2014, 20, 12245–12254. See also Ref. [12,14].

Communications

Reactive Intermediates

E. Tudela, J. González, R. Vicente, J. Santamaría, M. A. Rodríguez, A. Ballesteros* _____ ####-#####

Mechanistic Studies on the Rearrangement of 1-Alkenyl-2alkynylcyclopropanes: From Allylic Gold(I) Cations to Stable Carbocations



No longer elusive: The allylic gold(I) cation **2**, which has been proposed as an intermediate in the rearrangement of alkynylcyclopropanes (**1**) into alkynyl-cyclohexadienes (**3**), has been detected and characterized by NMR spectroscopy.

Participation of **2** was supported experimentally and theoretically, and through these studies, a stable homo-antiaromatic carbocation was isolated and characterized.