This is an open access article published under a Creative Commons Attribution (CC-BY) License, which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.



pubs.acs.org/OrgLett

Letter

Electrophilic Reactivities of Vinyl p-Quinone Methides

Andreas Eitzinger,[§] Robert J. Mayer,[§] Nathalie Hampel, Peter Mayer, Mario Waser, and Armin R. Ofial*

Cite This: https://dx.doi.org/10.1021/acs.orglett.0c00338





T he interest in vinyl *p*-quinone methides $(pVQMs)^{1-4}$ increased recently because it was shown that applying a 1,6-addition/cyclization strategy in reactions of *pVQMs* with sulfonium ylides,⁵ carbanions,⁶ or ammonium ylides^{7a} gave rise to vinyl cyclopropanes that rearranged to chiral spirocyclopentenes. Hence, *pVQMs* are versatile building blocks for the stereocontrolled synthesis of complex molecules.⁵⁻⁷ The further development of *pVQM*-based organic synthesis could clearly benefit from the knowledge of their electrophilic reactivity to define scope and limitations of their reactions with nucleophiles.¹⁻³



The electrophilicity of 6-aryl-substituted *p*-quinone methides (pQMs) had been studied by Mayr and co-workers⁸ who analyzed the second-order rate constants of the reactions of nucleophiles with *p*QMs according to the linear free energy relationship eq 1:⁹

$$\log k(20 \ ^{\circ}\text{C}) = s_{\text{N}}(N+E) \tag{1}$$

In this work, we set out to characterize the electrophilic reactivity E of pVQMs **1a-d** (Figure 1) by studying the kinetics of their reactions with the carbanions **2a-d** as reference nucleophiles in DMSO at 20 °C. In this way, pVQMs are integrated into Mayr's reactivity scales, which allows chemists to reliably predict the scope of their reactions with

$\int pVQM R \lambda_{max}$ (nm)			reference nucleophiles				
	1a 1b 1c 1d	OMe H CI NO ₂	432 405 408 429	$\odot \langle ^{NO_2} 2a \rangle$	N = 21.54 s _N = 0.62	⊜ CO₂Et CO₂Et	b $N = 20.22$ $s_N = 0.65$
8	L _R	- 2		⊖⟨CO ₂ Et 2c CN	N = 19.62 s _N = 0.67	⊖ CN CN 2	N = 19.36 $s_{\rm N} = 0.67$

Figure 1. *p*VQMs **1a**–**d** and reference nucleophiles **2a**–**d** used for the determination of their electrophilicities *E*. Nucleophilicity parameters *N* and *s*_N (in DMSO) were obtained from previous literature.^{8a,11}

structurally diverse nucleophiles when exploring novel organic syntheses. $^{10} \ \,$

The *p*VQMs **1a**–**d** were synthesized according to literature procedures and characterized by spectroscopic and electrochemical methods (Supporting Information). Single crystal Xray crystallography (Figure 2) revealed that the conjugated π systems in **1a**–**d** are slightly bent. The *p*VQMs **1a**–**d** are dyes with λ_{max} between 405 and 432 nm (in DMSO) and molar absorption coefficients in the range of 5 × 10⁴ M⁻¹ cm⁻¹ (Figure 1), which enabled us to follow their reactions with the colorless nucleophiles **2a**–**d** by photometry.

When solutions of the colored pVQMs 1 in DMSO (or d_6 -DMSO) were treated with the potassium salts of nucleophiles 2, a rapid fading of the color of 1 was observed. As described in Scheme 1, the reaction mixtures were then either analyzed by NMR methods or worked-up to isolate the Michael adducts. Mixtures of the regioisomers 3 and 4 were obtained via 1,6-and 1,8-additions of 2a, 2c, and 2d to pVQMs 1, which are ambident electrophiles. Only 2b underwent selective 1,8-additions to 1a-d, and the exclusive formation of regioisomers 4 could be detected in the crude reaction mixtures. Subsequent acidic workup of the reaction mixtures yielded the isolated products in good to excellent yields.

In the kinetic experiments the presence of a Brønsted acid is required to ensure fast protonation of the initial Michael adducts.^{8a} Solutions of the corresponding CH acids 2-H in DMSO were therefore only partially deprotonated by 0.5 equiv of KOtBu to generate DMSO stock solutions of the carbanions 2 as 1:1 mixtures with the CH acids 2-H. The reaction kinetics were determined by employing stopped-flow UV/vis photometry to follow the fading of the colored pVQMs 1 in their reactions with the colorless carbanions 2. By using a large

Received: January 24, 2020



pubs.acs.org/OrgLett



Figure 2. Single crystal X-ray structures of the pVQMs 1a-d. Thermal ellipsoids are shown at a 50% probability level. Bottom: Side views on 1a-d. The blue lines indicate the planes through the carbon atoms of the quinone moieties.

Scheme 1. Products of the Reactions of 1 with 2 in DMSO						
Ar 1a-d tBu	Nu (1–2 equiv) DMSO, r.t.	Ar 3 1,6-addition	+	Ar 4 (Bu 1,8-addition	3u H	
₽VQM	Nu	3	:	4 .	Yield ^a	
1a	2a -K	60 (d.r.: 1.8/1) (3aa)	:	40 (d.r.: 1.1/1) (4aa)	84%	
1a	2b -K	<1	:	>99 (4ab -K)	b	
1b	2b-K	<1	:	>99 (4bb)	90%	
1b	2b -K	<1	1	>99 (4bb)	77% ^c	
1b	2b -K	<1	1	>99 (4bb -K)	b	
1c	2b -K	<1	1	>99 (4cb -K)	b	
1d	2b -K	<1	:	>99 (4db -K)	b	
1b	2c -K	77 (3bc -K)	:	23 (4bc -K)	d	
		>95 (d.r.: 1/1) (3bc)	:	<5	75%	
1a	2d -K	40 (3ad)	:	60 (4ad)	74%	
1b	2d-K	42 (3bd)	1	58 (4bd)	63%	
1c	2d -K	40 (3cd)	1	60 (4cd)	90%	
1d	2d -K	70 (3dd)	:	30 (4dd)	99%	

^{*a*}Yields of isolated products after chromatographic workup. ^{*b*}Reaction performed in *d*₆-DMSO; the initially formed potassium phenolates **4Xb**-K were directly analyzed by NMR spectroscopic methods. ^{*c*}Reaction at 1 mmol scale. ^{*d*}Reaction performed in *d*₆-DMSO; the mixture of potassium salts **3bc**-K and **4bc**-K (both with deprotonated malononitrile moiety) was directly analyzed by NMR spectroscopic methods.

excess of the carbanions over the electrophiles, the resulting absorbance decays followed first-order kinetics. First-order rate constants k_{obs} were calculated by least-squares fitting of the single-exponential $A_t = A_0 \exp(-k_{obs}t) + C$ to the experimentally observed time-dependent absorbances (Figure 3a). Second-order rate constants k_2^{exptl} were subsequently obtained as the slopes of the linear correlations of k_{obs} with the concentrations of the carbanions [2] (Figure 3b; analogous correlations for all other electrophile–nucleophile combinations studied in this work are shown in the Supporting Information). Table 1 gathers the measured k_2^{exptl} values for the investigated reactions of pVQMs 1 with the carbanionic reference nucleophiles 2.

Next, we used eq 1 to perform a least-squares analysis, which allowed us to determine the electrophilicity parameters *E* for the *p*VQMs **1a**-**d** from k_2^{exptl} and the known nucleophilicity



Figure 3. (A) Decay of the absorbance A of **1c** ($c = 1.75 \times 10^{-5}$ M) at 408 nm in the reaction (DMSO, 20 °C) with **2b** ($c = 6.00 \times 10^{-4}$ M). (B) The slope of the linear correlation of k_{obs} with the concentration of **2b** yields the second-order rate constant k_2 .

Table 1. Second-Order Rate Constants for the Reactions of 1 with the Reference Nucleophiles 2 in DMSO at 20 $^{\circ}$ C

1	2	$k_2^{\text{exptl}} (\mathrm{M}^{-1} \mathrm{s}^{-1})$	$k_2^{\text{eq 1,a}} (\mathrm{M}^{-1} \mathrm{s}^{-1})$	$k_2^{\text{exptl}}/k_2^{\text{eq 1}}$			
1a	2a	5.72×10^{2}	3.59×10^{2}	1.6			
	2b	7.88×10^{1}	6.62×10^{1}	1.2			
	2c	3.09×10^{1}	2.98×10^{1}	1.0			
	2d	1.06×10^{1}	2.00×10^{1}	0.53			
E(1a) = -17.42							
1b	2a	7.11×10^{2}	6.49×10^2	1.1			
	2b	1.65×10^{2}	1.23×10^{2}	1.3			
	2c	6.16×10^{1}	5.65×10^{1}	1.1			
	2d	2.39×10^{1}	3.79×10^{1}	0.63			
E(1b) = -17.00							
1c	2a	1.14×10^{3}	8.21×10^{2}	1.4			
	2b	2.28×10^{2}	1.58×10^{2}	1.5			
	2c	7.44×10^{1}	7.30×10^{1}	1.0			
	2d	2.47×10^{1}	4.88×10^{1}	0.51			
$E(\mathbf{1c}) = -16.84$							
1d	2a	3.44×10^{3}	1.91×10^{3}	1.8			
	2b	5.92×10^{2}	3.82×10^2	1.5			
	2c	1.66×10^{2}	1.82×10^{2}	0.91			
	2d	5.08×10^{1}	1.22×10^{2}	0.42			
E(1d) = -16.25							

^{*a*}Second-order rate constant k_2 by applying eq 1.

parameters N (and s_N) of the reference nucleophiles (Table 1 and Figure S1, Supporting Information).

If compared to the analogously substituted pQMs the electrophilicity of pVQMs 1 is reduced by 1–2 orders of magnitude (Figure 4).⁸ Moreover, electronic substituent effects



Figure 4. Comparison of electrophilicities *E* of *p*VQMs 1 with those of analogously substituted pQMs.⁸ ^a Gray values are interpolated on the basis of the Hammett correlation described in ref 8b.

have a stronger impact on the electrophilicity of pQMs than on analogous π -extended pVQMs: While a change from a methoxy- to a nitro-substituent in pQMs increases their electrophilicity E by 1.7 units,^{8b} the same change in the series of pVQMs results in an increase of E by only 0.9 units.¹² This might be rationalized by the observed deviations from planarity in the solid state structures (Figure 2), which weaken the conjugation and thus attenuate the substituent effects.¹³

Quantum-chemical calculations were performed to gain a deeper understanding of the ambident reactivity of pVQMs. We calculated the Gibbs activation and reaction energies for the addition of nucleophiles **2b** and **2d** to the electrophile **1b** at the M06-2X/6-31+G(d,p) level considering solvation by the SMD solvation model for DMSO (Figure 5).¹⁴ In line with our experimental results and previous reports on the formation of regioisomeric mixtures upon concomitant attack of different types of nucleophiles at 1,6- and 1,8-positions of simple vinyl *p*-quinone methides,^{1c,3} the calculations show that the barriers for 1,6- and 1,8-addition differ only by 4–8 kJ mol⁻¹. For a



Figure 5. Reaction paths for additions of the nucleophiles **2b** (Acc = CO_2Et) and **2d** (Acc = CN) to *p*VQM **1b** (calculated at the SMD(DMSO)/M06-2X/6-31+G(d,p) level of theory).

given combination of 1 and 2, also both intermediates P1(1,6) and P1(1,8) are formed with similar Gibbs reaction energies.

Depending on the acidity of the $(Acc)_2CH$ moiety, the initially formed phenolate group in the adduct P1 might be protonated to yield the corresponding phenol P2. In line with NMR spectroscopic studies of the reactions (Supporting Information), the proton transfer is unfavored for 2b (pK_{aH} 18.7 for $(EtO_2C)_2MeC^{-})^{15}$ and the phenolate form P1(1,8) persists as detectable species in the reaction mixture (pK_{aH} 17.7 for 2,6-*tert*-butyl-4-methylphenolate).¹⁶ In additions of 2d (pK_{aH} 12.4 for ($NC)_2MeC^{-}$)¹⁷ to pVQMs, proton transfer from C–H to O–H occurs to yield a phenol. Owing to the energetic similarity of the competing reaction paths, the observed regiochemistry (1,6- vs 1,8-attack) for the attack of nucleophiles at pVQMs does not follow a clear pattern but seems to depend on subtle effects, which are introduced by the nature of the nucleophile.

Nevertheless, the determined electrophilicity parameters E for 1a-d can be used to rationalize reported reactions and, more intriguingly, to predict new reactions. In Figure 6, the



Figure 6. Ranking of *p*VQMs 1a-d in the Mayr reactivity scales (nucleophilicities *N* in DMSO if not mentioned otherwise).¹⁰

electrophilicity and nucleophilicity scales are arranged such that (E + N) = -3. Reaction partners on the same horizontal level react (somewhat dependent on the s_N parameter) with second-order rate constants of 10^{-3} to 10^{-2} M⁻¹ s⁻¹ at 20 °C. Accordingly, reactions of *p*VQMs 1 with sulfonium ylides, such as 12, and α -bromo malonate (*N* determined for the chloroderivative 6) have been described in the literature.^{5,6} Nucleophiles located at levels below that of the *p*VQMs can be expected to react even more rapidly.

Based on the prediction that reactions of **1** with nucleophiles of N > 13 should occur at 20 °C,¹⁸ we studied the reactions of *p*VQMs **1** with carbanions (**11** and **15**), the pyridinium ylide **14**, the heteroatom nucleophiles MeS⁻ and pyrrolidine (7), and the hydride donor NaBH₄ (**5**). For all combinations, the reaction products could be isolated in good to excellent yields without further optimization (Table 2).

As found in the initial product studies (Scheme 1), different regioisomers were also observed for the reactions of 1a-d with the nucleophiles in Table 2: While 1,6-addition was the preferred reaction mode for NaBH₄ (5), highly nucleophilic carbanions (11 and 15), and the pyridinium ylide 14, products of 1,8-attack were observed for 7 and NaSMe. We rationalize the formation of the butadienyl-substituted phenol 17 (Table 2, entry 2) by a cyclopropanation/ring opening sequence as previously observed for reactions of pQMs with α -halotosylmethyl anions.^{19,20} Interestingly, the reaction of the

Table 2. Scope of pVQM (1) Reactions with Nucleophiles



pyridinium ylide 14 with the pVQM 1a gave the pyridinium bromide 18 (Table 2, entry 3), which is in contrast to reactions of ammonium ylides with pVQMs which furnish spirocyclic products.⁷

In conclusion, we have characterized the Mayr electrophilicities *E* of the vinyl *p*-quinone methides 1a-d by analyzing the kinetics and products of their reactions with carbanions in DMSO. In agreement with earlier findings on the regioselectivities of nucleophile additions to 2,6-dimethoxy-4-(2propenylidene)-2,5-cyclohexadien-1-one and eugenol-derived vinylic p-quinone methides,^{1c,3} the pVQMs 1 are ambident electrophiles that have similar 1,6- and 1,8-reactivities. While the results of our experiments do not allow us to predict the regiochemistry of the nucleophilic attack at pVQMs, the determined Mayr E parameters reliably reflect the general electrophilic reactivity of these electron-deficient π -systems. Application of the electrophilicity parameters E in eq 1 not only rationalizes reported reactions but also empowers chemists to systematically predict novel combinations of pVQMs with nucleophiles. We demonstrated that uncatalyzed reactions of 1a-d with different types of C-, N-, S-, and Hnucleophiles with N > 14 are feasible at ambient temperature²¹ and lead to novel types of conjugate 1,6- and 1,8-adducts of pVQMs.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.orglett.0c00338.

Synthetic procedures, analytical data, X-ray structure determinations, details on kinetic measurements and

quantum-chemical calculations, and copies of NMR spectra (PDF)

Coordinates of optimized structures (ZIP)

Accession Codes

CCDC 1973228–1973231 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Armin R. Ofial – Department Chemie, Ludwig-Maximilians-Universität München, 81377 München, Germany; orcid.org/0000-0002-9600-2793; Email: ofial@lmu.de

Authors

- Andreas Eitzinger Institute of Organic Chemistry, Johannes Kepler University Linz, 4040 Linz, Austria
- Robert J. Mayer Department Chemie, Ludwig-Maximilians-Universität München, 81377 München, Germany
- Nathalie Hampel Department Chemie, Ludwig-Maximilians-Universität München, 81377 München, Germany
- Peter Mayer Department Chemie, Ludwig-Maximilians-Universität München, 81377 München, Germany
- Mario Waser Institute of Organic Chemistry, Johannes Kepler University Linz, 4040 Linz, Austria; Occid.org/0000-0002-8421-8642

Complete contact information is available at: https://pubs.acs.org/10.1021/acs.orglett.0c00338

Author Contributions

[§]A.E. and R.J.M. contributed equally.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Financial support through the Austrian Science Funds (FWF): Project No. P30237 (financed by the Austrian National Foundation for Research, Technology and Development and Research Department of the State of Upper Austria) is gratefully acknowledged. The NMR spectrometers at JKU Linz were acquired in collaboration with the University of South Bohemia (CZ) with financial support from the European Union through the EFRE INTERREG IV ETC-AT-CZ program (project M00146, "RERI-uasb"). RJ.M. thanks the Fonds der Chemischen Industrie for a Kekulé fellowship.

REFERENCES

(1) (a) Filar, L. J.; Winstein, S. Preparation and Behavior of Simple Quinone Methides. *Tetrahedron Lett.* 1960, 1 (25), 9–16.
(b) Zanarotti, A. Preparation and Reactivity of 2,6-Dimethoxy-4-allylidene-2,5-cyclohexadien-1-one (Vinyl Quinone Methide) A Novel Synthesis of Sinapyl Alcohol. *Tetrahedron Lett.* 1982, 23, 3815–3818.
(c) Zanarotti, A. Synthesis and Reactivity of Vinyl Quinone Methides. *J. Org. Chem.* 1985, *50*, 941–945.

(2) Hemmingson, J. A.; Leary, G. The Chemistry of Reactive Lignin Intermediates. Part II. Addition Reactions of Vinyl-substituted Quinone Methides in Aqueous Solution. J. Chem. Soc., Perkin Trans. 2 1975, 2, 1584–1587. (3) (a) Bolton, J. L.; Comeau, E.; Vukomanovic, V. The influence of 4-alkyl substituents on the formation and reactivity of 2-methoxyquinone methides: evidence that extended π -conjugation dramatically stabilizes the quinone methide formed from eugenol. *Chem.-Biol. Interact.* **1995**, *95*, 279–290. (b) Krol, E. S.; Bolton, J. L. Oxidation of 4-alkylphenols and catechols by tyrosinase: ortho-substituents alter the mechanism of quinoid formation. *Chem.-Biol. Interact.* **1997**, *104*, 11–27.

(4) Reichl, K. D.; Smith, M. J.; Song, M. K.; Johnson, R. P.; Porco, J. A., Jr. Biomimetic Total Synthesis of (\pm) -Griffipavixanthone via a Cationic Cycloaddition–Cyclization Cascade. J. Am. Chem. Soc. **2017**, 139, 14053–14056.

(5) Zhang, X.-Z.; Deng, Y.-H.; Gan, K.-J.; Yan, X.; Yu, K.-Y.; Wang, F.-X.; Fan, C.-A. Tandem Spirocyclopropanation/Rearrangement Reaction of Vinyl *p*-Quinone Methides with Sulfonium Salts: Synthesis of Spirocyclopentenyl *p*-Dienones. *Org. Lett.* **2017**, *19*, 1752–1755.

(6) Yuan, Z.; Gai, K.; Wu, Y.; Wu, J.; Lin, A.; Yao, H. Tandem 1,6addition/cyclization/vinylcyclopropane rearrangement at low temperature under metal-free conditions: an approach to spiro[4.5]cyclohexadienones. *Chem. Commun.* **2017**, *53*, 3485–3488.

(7) (a) Roiser, L.; Zielke, K.; Waser, M. Formal (4 + 1)-Cyclization of Ammonium Ylides with Vinylogous *para*-Quinone Methides. *Synthesis* **2018**, *50*, 4047–4054. (b) Eitzinger, A.; Winter, M.; Schörgenhumer, J.; Waser, M. Quaternary $\beta^{2,2}$ -amino acid derivatives by asymmetric addition of isoxazolidin-5-ones to para-quinone methides. *Chem. Commun.* **2020**, *56*, 579–582.

(8) (a) Lucius, R.; Loos, R.; Mayr, H. Kinetic Studies of Carbocation-Carbanion Combinations: Key to a General Concept of Polar Organic Reactivity. *Angew. Chem., Int. Ed.* **2002**, *41*, 91–95. (b) Richter, D.; Hampel, N.; Singer, T.; Ofial, A. R.; Mayr, H. Synthesis and Characterization of Novel Quinone Methides: Reference Electrophiles for the Construction of Nucleophilicity Scales. *Eur. J. Org. Chem.* **2009**, 3203–3211.

(9) (a) Mayr, H.; Patz, M. Scales of Nucleophilicity and Electrophilicity: A System for Ordering Polar Organic and Organometallic Reactions. *Angew. Chem., Int. Ed. Engl.* **1994**, 33, 938–957. (b) Mayr, H.; Ofial, A. R. A Quantitative Approach to Polar Organic Reactivity. *SAR QSAR Environ. Res.* **2015**, *26*, 619–646. (c) Mayr, H. Reactivity Scales for Quantifying Polar Organic Reactivity: The Benzhydrylium Methodology. *Tetrahedron* **2015**, *71*, 5095–5111.

(10) Mayr's reactivity parameters E_i , N_i , and s_{NJ} , which have been determined for >300 electrophiles and >1200 nucleophiles, can be retrieved from the freely accessible database at www.cup.lmu.de/oc/mayr/DBintro.html.

(11) Bug, T.; Lemek, T.; Mayr, H. Nucleophilicities of Nitroalkyl Anions. J. Org. Chem. 2004, 69, 7565–7576.

(12) For an analogous pattern when comparing the electrophilicities of benzhydrylium ions (Ar_2CH^+) with those of 1,3-diarylallylium ions ($ArCH=CH-CH^+-Ar$), see: Troshin, K.; Schindele, C.; Mayr, H. Electrophilicities of Symmetrically Substituted 1,3-Diarylallyl Cations. *J. Org. Chem.* **2011**, *76*, 9391–9408.

(13) See Supporting Information (Figure S2) for a correlation of electrophilicities E(1) with Hammett substituent parameters σ_p .

(14) For details and full citations of the computational methods, see the Supporting Information.

(15) Bordwell, F. G.; Harrelson, J. A., Jr.; Satish, A. V. Oxidation potentials of carbanions and homolytic bond dissociation energies of their conjugate acids. *J. Org. Chem.* **1989**, *54*, 3101–3105.

(16) Bordwell, F. G.; Zhang, X.-M. Acidities and homolytic bond dissociation enthalpies of 4-substituted-2,6-di-*tert*-butylphenols. J. Phys. Org. Chem. **1995**, 8, 529–535.

(17) Bordwell, F. G.; Bartmess, J. E.; Hautala, J. A. Alkyl effects on equilibrium acidities of carbon acids in protic and dipolar aprotic media and the gas phase. *J. Org. Chem.* **1978**, *43*, 3095–3101.

(18) The synthetic potential of the weakly reactive (N < 13) nucleophiles 22, 23, and 24 has been assessed in reactions with the pVQM 1b (E = -17.0). NMR spectroscopic analysis showed that, in CD₂Cl₂ solutions at room temperature, 1b reacted neither with 22

nor with **23** within 4 days. The reaction of **1b** with imidazole (**24**, *N* = 11.47 and $s_N = 0.79$ in MeCN) is predicted to be sluggish at 20 °C ($k_2^{eq 1} = 4.3 \times 10^{-5} \text{ M}^{-1} \text{ s}^{-1}$). Accordingly, we observed only a low degree of conversion (<20%) for **1b** at the end of a 4 days observation period after mixing it with 2 equiv of **24** in CD₃CN.

(19) Li, Z.; Chen, Q.; Mayer, P.; Mayr, H. Nucleophilicity Parameters of Arylsulfonyl-Substituted Halomethyl Anions. J. Org. Chem. 2017, 82, 2011–2017.

(20) Groszek, G.; Błażej, S.; Brud, A.; Świerczyński, D.; Lemek, T. Reactions of carbanions derived from α -substituted-methyl tolyl sulfones with quinone methides as Michael acceptors. *Tetrahedron* **2006**, *62*, 2622–2630.

(21) For an example of a Lewis acid catalyzed reaction of **1b**, see: Hao, Y.-J.; Hu, X.-S.; Yu, J.-S.; Zhou, F.; Zhou, Y.; Zhou, J. An efficient Fe(III)-catalyzed 1,6-conjugate addition of *para*-quinone methides with fluorinated silyl enol ethers toward $\beta_i\beta$ -diaryl α fluorinated ketones. *Tetrahedron* **2018**, 74, 7395–7398.