

Article

## Strongly directing substituents in the radical arylation of substituted benzenes

Josefa Hofmann, Timothy Clark, and Markus Rolf Heinrich

*J. Org. Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.6b01840 • Publication Date (Web): 19 Sep 2016

Downloaded from <http://pubs.acs.org> on September 23, 2016

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

# Strongly directing substituents in the radical arylation of substituted benzenes


Josefa Hofmann,<sup>a</sup> Timothy Clark,<sup>b</sup> Markus R. Heinrich<sup>a,\*</sup>

<sup>a</sup>Department of Chemistry and Pharmacy, Pharmaceutical Chemistry, Friedrich-Alexander-Universität Erlangen-Nürnberg, Schuhstraße 19, 91052 Erlangen, Germany

<sup>b</sup>Computer-Chemie-Centrum and Interdisciplinary Center for Molecular Materials, Friedrich-Alexander-Universität Erlangen-Nürnberg, Nögelsbachstraße 25, 91052 Erlangen, Germany.

markus.heinrich@fau.de

RECEIVED DATE (to be automatically inserted after your manuscript is accepted if required according to the journal that you are submitting your paper to)



## ABSTRACT:

Although the general interest in radical arylation reactions has grown rapidly in recent years, poor regioselectivities and the need to use a large excess of the radical-accepting arene have hindered their application to substituted benzenes. We now describe experimental and

computational investigations into the substituent effects that lead to regioselective addition based on the recent discovery of anilines as outstanding substrates for radical arylations.

## Introduction

The Gomberg-Bachmann reaction, which dates back to 1924,<sup>1</sup> represents one of the fundamental transformations in aryl radical chemistry.<sup>2,3</sup> Starting from aryl diazonium salts as classical radical sources,<sup>4</sup> and arenes or heteroarenes acting as radical acceptors, a broad range of biaryls can be obtained in a reaction<sup>5</sup> whose result is comparable to an aryl-C-H activation.<sup>6</sup> Recent versions of the Gomberg-Bachmann reaction rely on a photocatalytic conversion of diazonium salts,<sup>7</sup> on aryl radical generation from chloro-, bromo- or iodobenzenes in the presence of strong bases,<sup>8</sup> on arylhydrazines,<sup>9</sup> or on the *in situ* diazotization of anilines.<sup>10</sup>

Although Gomberg-Bachmann reactions could, on this basis, be very attractive due to the cheap and readily available starting materials, their application, also in the newly developed versions, has remained mostly limited to unsubstituted benzene and selected heteroarenes.<sup>7-10</sup> Explanations for this are both the relatively slow addition of aryl radicals to benzene and its derivatives,<sup>11</sup> which requires the use of the radical acceptor in large excess, and the low regioselectivities observed for most substituted benzenes.<sup>5,12</sup> Heteroarenes, on the other hand, and hereby especially electron-rich furans and pyrroles, show increased reactivity towards aryl radicals and far higher regioselectivities.<sup>3e,7,8,13</sup>

The recent extension of the Gomberg-Bachmann reaction to anilines provided motivation to investigate the suitability of substituted benzenes as aryl radical acceptors in more detail.<sup>14</sup> Both in these reactions, and in consecutively developed arylations with arylhydrazines,<sup>15</sup> anilines showed outstanding properties as aryl radical acceptors in two ways: Firstly, the aryl radical addition to anilines was found to proceed about one order of magnitude faster than to

classically favored substrates such as nitrobenzenes or anisoles,<sup>15a</sup> and secondly, comparably high regioselectivities were obtained.<sup>14,15</sup>


Little attention has been paid in the literature over the last decades to how substituents affect the regioselectivity in radical arylations of benzene derivatives with the last comprehensive review dating back to 1973.<sup>12</sup> Early studies had shown that a variation of substituents on the attacking phenyl radical does influence the substrate selectivity in competition experiments with nitrobenzene and benzene, but not the regioselectivity of the substitution at nitrobenzene.<sup>12,16</sup> Such observations led to a discussion of “polarity of free radicals”.<sup>17</sup> All substituents on the radical acceptor were considered to be weakly activating,<sup>12</sup> and comparably high regioselectivities were only obtained for particular combinations of donor-substituted phenyl radicals with acceptor-substituted benzenes, or *vice versa*.<sup>18</sup> As these studies have shown that it is particularly difficult to achieve regioselectivity with aryl radicals such as the 4-chlorophenyl radical,<sup>19</sup> this radical was chosen for the experiments reported in this communication. Based on experimental results, we will give insight into the fundamental question as to whether kinetic or thermodynamic factors play a decisive role in the addition step. So far, the low selectivity of phenyl radicals in arylations of substituted benzenes has been attributed to a “little development of the new bond at the transition state”,<sup>12</sup> so that “the stability of the product radicals is also of importance”.<sup>20</sup>

## Results and Discussion

Three recently developed protocols, starting either from 4-chlorophenyldiazonium chloride (**1**) or 4-chlorophenylhydrazine (**2**), were chosen to collect the required experimental data, as these procedures allow in particular the radical arylation of anilines (Scheme 1).<sup>14a,15a,15b</sup> All reactions were carried out under air atmosphere to ensure that rearomatization does not become an influential factor.<sup>21</sup> Moreover, the aromatic substrates **3-7** were not used as solvents, but in a lower excess of 10-20 equivalents. In this way, the results also give an



impression on the synthetic applicability and the relative reactivity of the arenes can be evaluated, as the aryl radicals are not trapped quantitatively.

**Scheme 1.** Reaction conditions A, B and C used for radical arylations.







Tables 1-3 summarize the yields and product ratios obtained with anilines **3** (Table 1), phenols **4** and phenylethers **5** (Table 2), and with benzonitriles **6** and nitrobenzene (**7a**) (Table 3). The conditions A, B and C refer to the reactions shown in Scheme 1. If available, further data from literature (conditions D-I) have been added for comparison of regioselectivities.

**Table 1.** Arylation of aniline (**3a**) and 4-fluoroaniline (**3b**)

biphenyl	isomers	conditions			
		A	B	C	D <sup>a</sup>
 <b>8a</b> (from <b>3a</b> )	2 ( <b>8a</b> )	80%	54%	60%	62%
	3 ( <b>8a'</b> )	76	76	85	75
	4 ( <b>8a''</b> )	--	--	--	--
 <b>8b</b> (from <b>3b</b> )	2 ( <b>8b</b> )	68%	53%	55%	61%
	3 ( <b>8b'</b> )	100	100	100	100
	4 ( <b>8b''</b> )	--	--	--	--




<sup>a</sup>Conditions D: See ref. 15c.

**Table 2.** Arylation of phenol (**4a**), 4-fluorophenol (**4b**), anisole (**5a**) and 4-fluoroanisole (**5b**).

biphenyl	isomers	conditions					
		A	B	C	E <sup>a</sup>	F <sup>b</sup>	G <sup>c</sup>
 <b>9a</b> (from <b>4a</b> )	□		57%	58%	62%		
	2 ( <b>9a</b> )		70	76	68		
	3 ( <b>9a'</b> )		--	--	--		
	4 ( <b>9a''</b> )		30	24	32		
 <b>9b</b> (from <b>4b</b> )			49%	40%	56%		
	2 ( <b>9b</b> )		100	100	100		
	3 ( <b>9b'</b> )		--	--	--		
 <b>10a</b> (from <b>5a</b> )		31%	34%	35%		59%	48%
	2 ( <b>10a</b> )		64	73	73	68	75
	3 ( <b>10a'</b> )		17	12	12	17	12
	4 ( <b>10a''</b> )		19	15	15	15	13
 <b>10b</b> (from <b>5b</b> )		36%	31%	36%			
	2 ( <b>10b</b> )		72	88	89		
	3 ( <b>10b'</b> )		28	12	11		

<sup>a</sup>Conditions E: Reaction of 4-cyanophenyl radical with phenolate or 4-bromophenolate, see ref. 22. <sup>b</sup>Conditions F: Reaction of 4-methylphenyl radical with anisole, see ref. 8c. <sup>c</sup>Conditions G: Reaction of 3-chlorophenyl radical with anisole, see ref. 5.

**Table 3.** Arylation of benzonitrile (**6a**), 4-fluorobenzonitrile (**6b**) and nitrobenzene (**7a**).

biphenyl	isomers	conditions				
		A	B	C	H <sup>a</sup>	I <sup>b</sup>
 <b>11a</b> (from <b>6a</b> )		49%	29%	62%	58%	
	2 ( <b>11a</b> )	48	62	52	51	
	3 ( <b>11a'</b> )	16	11	15	14	
	4 ( <b>11a''</b> )	36	27	33	35	
 <b>11b</b> (from <b>6b</b> )		20%	13%	49%		
	2 ( <b>11b</b> )	61	67	77		
	3 ( <b>11b'</b> )	39	33	23		
 <b>12a</b> (from <b>7a</b> )		36%	52%	38%		30%
	2 ( <b>12a</b> )	44	42	41		55
	3 ( <b>12a'</b> )	14	16	18		4
	4 ( <b>12a''</b> )	42	42	41		41

<sup>a</sup>Conditions H: See ref. 5; <sup>b</sup>Conditions I: Reaction of phenyl radical with nitrobenzene, see refs. 23,24.

A comparison of the average yields obtained with the mono-substituted radical acceptors **3a**, **4a**, **5a**, **6a** and **7a** under all available conditions gave the following trend in overall reactivity: Ph-NH<sub>2</sub> > Ph-O<sup>-</sup> > PhCN > PhNO<sub>2</sub> ≈ PhOMe, which is in agreement with earlier experimental studies.<sup>15a</sup> Note that, under the reaction conditions B, C and E, phenol (**4a**) and 4-fluorophenol (**4b**) are present as phenolates.


In terms of regioselectivity, only the anilines **3a-c** (Table 1) and phenolates of **4a,b** (Table 2, entries 1 and 2) were able to suppress the formation of *meta*-isomers (3-isomers), independent of the presence of a substituent in *para*-position. Arylation of the anisoles **5a,b** (Table 2, entries 3 and 4), benzonitriles **6a,b** and nitrobenzene (**7a**), in contrast, led to all basically possible regioisomers. Remarkably, the effect of the electron-donating methoxy group differs from those of the electron-withdrawing cyano and nitro groups, as anisole (**5a**) showed a preference for the 2-isomer (2/3/4 = 71:14:15), whereas benzonitrile (**6a**) and nitrobenzene (**7a**) were attacked more frequently in 4-position via average selectivities of 2/3/4 = 53:14:33 and 2/3/4 = 45:13:42, respectively.<sup>25</sup> This effect may be related to a three-electron interaction between the radical and the oxygen atom of the methoxy group in the transition state. The importance of the 4-position in arylations of electron-deficient arenes also became obvious from the significant decrease in yield caused by blocking this position by a fluorine substituent (Table 3, entries 1 and 2). Arylation experiments with 4-fluoronitrobenzene are not included since they are complicated by competing nucleophilic substitution.

In the next step, density-functional theory (DFT) calculations were carried out to determine whether the experimentally observed substituent effects in radical arylations could be predicted computationally. It was of particular interest in this respect to evaluate the influence of the transition state, **TS** compared to the stabilization of the cyclohexadienyl radical adduct **CA** (Scheme 2).

All calculations were performed with Gaussian09, revision C.01.<sup>26</sup> Geometries were optimized using the B3LYP functional,<sup>27,28</sup> and the aug-cc-pVDZ basis set.<sup>29,30</sup> Structures were shown to be minima or transition states by calculating the normal vibrations within the harmonic approximation. Energies relative to the separated reactants are reported including vibrational zero-point energy taken from these calculations. B3LYP/aug-cc-pVDZ calculations have been shown to underestimate radical activation energies by several kcal


mol<sup>-1</sup> in comparison to coupled-cluster reference calculations,<sup>31</sup> although we expect the trends to be correct for such closely related reactions.

**Scheme 2.** Reaction course.



Early calculations by James and Stuart<sup>32</sup> on the hydrogen atom addition to benzene gave an activation energy of ca. 4 kcal mol<sup>-1</sup> and suggested that the addition step is exothermic by -28 kcal mol<sup>-1</sup>. From these values, the activation energy for the attack of a phenyl radical onto benzene was estimated to > 3 kcal mol<sup>-1</sup> and the related reaction energy to -18 kcal mol<sup>-1</sup>.<sup>33</sup> The results from our calculations are summarized in Figure 1 and Tables 4 and 5.

**Figure 1.** Schematic reaction profiles for the arylation of aniline and benzene with the 4-chlorophenyl radical.




The data obtained for the arylation of aniline (**3a**) and benzene with the 4-chlorophenyl radical revealed that the radical attack on electron-rich aromatic systems proceeds via a pre-reaction complex that has not been considered previously (Figure 2). One of the two more polarized C-H bonds in *ortho* position to the radical center points to the  $\pi$ -system of the radical accepting aniline. As shown by the activation energies summarized in Table 4, this



complex plays an exceptional role in the arylation of phenolate ions. It is likely that both specific solvation and counterion-effects are important for these reactions, but considering explicit solvent molecules and/or counterions would be outside the scope of this article.

**Figure 2.** Pre-reaction complex formed on attack of the 4-chlorophenyl radical on aniline (left) and on the *ortho*-position of the phenolate anion (right). Distances are given in Å.



**Table 4.** Calculated activation barriers for the attack of a 4-chlorophenyl radical.

directing substituent R <sup>1</sup> =	activation barrier <sup>a</sup> for an attack in		
	2-position	3-position	4-position
3a: NH <sub>2</sub>	3.3	5.0	4.1
4a: O <sup>-</sup>	-9.8 <sup>b</sup>	-2.3	-9.2
5a: OCH <sub>3</sub>	3.7	5.4	4.7
6a: CN	4.5	4.8	4.6
7a: NO <sub>2</sub>	3.8	5.6	4.3
benzene: H		5.3	

<sup>a</sup>Relative to the separated reactants (B3LYP/aug-cc-pVDZ + zero-point energy). <sup>b</sup>A pre-reaction complex with a binding energy of -9.8 kcal mol<sup>-1</sup> is formed (see Figure 2). The activation barrier from this complex to the 2-isomer is essentially zero.

**Table 5.** Calculated heats of reaction for the attack of a 4-chlorophenyl radical.

radical acceptor R <sup>1</sup> = , R <sup>2</sup> =	heats of the addition for an attack in		
	2-position	3-position	4-position
3a: NH <sub>2</sub> , H	-22.8	-19.3	-21.0
3b: NH <sub>2</sub> , F	-24.0	-20.6	
4a: O <sup>-</sup> , H	-35.0 <sup>b</sup>	-22.0	-27.7
4b: O <sup>-</sup> , F	-37.8 <sup>b</sup>	-22.3	
5a: OCH <sub>3</sub> , H	-22.3	-18.9	-19.9
5b: OCH <sub>3</sub> , F	-23.1	-19.6	
6a: CN, H	-22.1	-20.3	-20.6
6b: CN, F	-21.9	-19.9	
7a: NO <sub>2</sub> , H	-22.6	-19.2	-23.5
benzene: H		-19.7	


<sup>a</sup>Relative (B3LYP/aug-cc-pVDZ + zero-point energy) heats of addition and e to the separated reactants. <sup>b</sup>A pre-reaction complex with a binding energy of -9.8 kcal mol<sup>-1</sup> is formed. The activation barrier from this complex to the *ortho*-product is essentially zero.

The activation energies (relative to the separated reactants) for the neutral systems lie in range from 3.3 to 5.6 kcal mol<sup>-1</sup>, consistent with the estimated value of > 3 kcal mol<sup>-1</sup> (Table 4). The

calculated heats of reaction lie in the range of  $-18.9$  to  $-24.0$  kcal mol<sup>-1</sup>, again consistent with the experimental estimate of  $-18$  kcal mol<sup>-1</sup> for the addition of the phenyl radical to benzene.<sup>33,34</sup> The introduction of the fluorine atom in 4-position generally led to a weak relative stabilization in the range of  $-0.3$  to  $-1.3$  kcal mol<sup>-1</sup>. A comparison of the heats of reaction to those of benzene ( $-19.7$  kcal mol<sup>-1</sup>) provided support for the long-standing assumption that all substituents on the radical acceptor are weakly activating, independent of their electron-donating or withdrawing character. The activating effect is thereby much more pronounced for 2- and 4-positions than for 3-positions.


Figure 3 shows a plot of the calculated heats of reaction against the activation energies. The correlation between the two quantities is good, so that we can conclude that the radical addition is compatible with the Bell-Evans-Polanyi principle.<sup>35</sup> However, the 3- and 4-nitro and 2-cyano cases deviate significantly and have been omitted from the correlation (indicated in grey in Figure 3). Quite generally, the  $\pi$ -acceptor substituents lie above the correlation line (i.e. the activation energies are higher than would be expected from the heat of reaction). Nevertheless, we can conclude that the differences in calculated activation barriers for the radical addition to substituted benzenes are largely controlled by the stabilization of the product radical without significant specific kinetic effects. The slope of the calculated regression line is approximately 0.5, suggesting that radical stabilization effects by the substituents are approximately 50% developed in the transition states. The  $\pi$ -acceptors are less well able to stabilize the transition states than the other substituents.

**Figure 3.** Calculated activation energies plotted vs. the calculated heats of reaction (all kcal mol<sup>-1</sup>) for the addition of the 4-chlorophenyl radical to neutral monosubstituted benzenes.<sup>a</sup>



<sup>a</sup>The regression line and the correlation equation do not include the two data-points plotted in grey, which represent significant outliers, and are both associated with strong  $\pi$ -acceptors, as discussed in the text.

**Figure 4.** Calculated and experimental yields of the individual neutral arylated products.<sup>a</sup>



<sup>a</sup>The black filled circles and the regression line and equation are those obtained from the calculated gas-phase activation energies assuming kinetic control with the addition as the rate-determining step. The red open circles indicate the predicted yields obtained by assuming thermodynamic control of the product radicals. The line is that for perfect 1:1 agreement.

The question of whether the product distributions are best described by kinetic or thermodynamic control of the addition reactions can be answered by comparing calculated product distributions with those observed experimentally. Figure 4 shows a plot of the calculated (from transition-state theory) % yield for each product of the addition of *p*-chlorophenyl radical to the monosubstituted benzenes vs. the experimental results expressed as the mean of the product distributions for all reaction conditions with error bars that cover

the range found experimentally. The agreement (RMSE = 15%) is good, so that we can conclude that the product distributions are kinetically controlled. The corresponding correlation obtained by assuming thermodynamic control gives a significantly worse RMSE (21%). The product yields calculated assuming thermodynamic control are shown as red open circles in Figure 4. Note that a statistical factor of 2:2:1 has been used for the 2-, 3-, and 4-isomers, respectively, in order to take the fact that there is only one 4-position available for each substituted benzene into account. The assumption of kinetic control for the arylation of nitrobenzene leads to a predicted 2/4- ratio of 79:16, representing the strongest deviation in this series, compared to 30:69 for thermodynamic control. As experiment (Table 3) gives approximately equal yields of the two isomers with a slight preference for the 2-isomer, the arylation of nitrobenzene can be considered to proceed with a stronger thermodynamic influence.

The combination of our experimental and theoretical results therefore suggests that the addition of aryl radicals to substituted benzenes largely takes place under kinetic control with the radical addition as the rate-determining step and that substituent effects in the product radicals are approximately 50% effective in the transition states. These conclusions are consistent with those of an earlier theoretical study by Zhang<sup>34</sup> on the arylation of aniline. Zhang concluded that aryl radical addition to neutral anilines is the rate-determining step in a kinetically controlled reaction, whereas arylation to protonated anilines may be thermodynamically controlled.

Note that, even though the calculations can predict the product distributions well, the calculated activation energies do not agree completely with the accepted reactivity series  $\text{Ph-NH}_2 > \text{Ph-O}^- > \text{PhCN} > \text{PhNO}_2 \approx \text{PhOMe}$ ,<sup>15a,26,36</sup> in order of increasing calculated activation energy for the most reactive isomer, the calculations predict  $\text{Ph-NH}_2 > \text{Ph-OCH}_3 \approx \text{Ph-NO}_2 > \text{Ph-CN} > \text{benzene}$ .  $\text{Ph-O}^-$  cannot be treated on the same footing as the neutral systems

because ion-pairing and solvation effects are likely to be dominant in determining reactions rates for the phenolate anion.

## Conclusions

In summary, it has been shown that DFT calculations can be used to predict experimental product distributions in radical arylations of benzenes. Although not as regioselective as arylations of alkenes, the aryl radical addition to substituted benzenes still proceeds with a high degree of kinetically controlled selectivity. Substituent effects in the resulting cyclohexadienyl radical adduct are reflected to about 50% in the transition state. All reactions pass through a pre-reaction complex, which was found to be particularly strongly stabilized in the case of an aryl radical attack on phenolate ions. These results give more detailed insight into the long-standing question of how substituents affect the regioselectivity in radical arylations of substituted benzenes.

## Experimental section

**General Experimental:** Solvents and reagents were used as received.  $^1\text{H}$  NMR spectra were recorded on 360 and 600 MHz spectrometers using  $\text{CDCl}_3$  and  $\text{CD}_3\text{OD}$  as solvents referenced to TMS (0 ppm),  $\text{CHCl}_3$  (7.26 ppm) and  $\text{CHD}_2\text{OD}$  (3.31 ppm). Chemical shifts are reported in parts per million (ppm). Coupling constants are in Hertz ( $J$  Hz). The following abbreviations are used for the description of signals: s (singlet), d (doublet), dd (double doublet), t (triplet), q (quadruplet), m (multiplet).  $^{13}\text{C}$  NMR spectra were recorded at 90.6 and 150.9 MHz in  $\text{CDCl}_3$  and  $\text{CD}_3\text{OD}$  using  $\text{CDCl}_3$  (77.0 ppm) and  $\text{CD}_3\text{OD}$  (49.5 ppm) as standard. Chemical shifts are given in parts per million (ppm).  $^{19}\text{F}$  NMR spectra were recorded at 338.8 MHz using  $\text{CFCl}_3$  (0 ppm) or  $\text{C}_6\text{F}_6$  (-164.9 ppm) as standard. Mass spectra were recorded using electron impact (EI). Analytical TLC was carried out on

Merck silica gel plates using short wave (254 nm) UV light and Ninhydrin to visualise components. Silica gel (Kieselgel 60, 40-63  $\mu\text{m}$ , Merck) was used for flash column chromatography. Yields obtained after purification are summarized in Tables 1-3 in the article.

#### **General procedure for the synthesis of biphenyls under Gomberg-Bachmann conditions**

##### **(conditions A): Preparation of the aryl diazotate by diazotization and addition of base: A**

degassed solution of sodium nitrite (20.0 mmol, 1.38 g) in water (10 mL) was added dropwise to an ice-cooled degassed solution of the aniline (20.0 mmol) in hydrochloric acid (3 N, 20 mL) and water (20 mL) over a period of 15 min. The clear solution was stirred for 20 more minutes at 0 °C. An aliquot of this 0.4 M aryldiazonium chloride solution (2.00 mmol, 5.00 mL) was treated with a pre-cooled aqueous solution of sodium hydroxide (4 N, 3 mL).

The resulting solution/suspension of the aryl diazotate was then used for the aryl–aryl coupling. **Radical arylation of substituted benzenes with a previously prepared aryl**

**diazotate:** The previously prepared solution/suspension of the aryl diazotate (2.00 mmol, 5.00 mL) was added dropwise to the substituted benzene (20.0 mmol) at 75–95 °C under vigorous stirring over a period of 10–15 min. After the addition was complete, the mixture was left to stir for 10 more minutes. The resulting reaction mixture was then extracted with organic solvents (e.g. diethyl ether or ethyl acetate, 3  $\times$  75 mL). The combined organic phases were washed with saturated aqueous sodium chloride and dried over sodium sulfate. The solvent was removed under reduced pressure and the resulting product was dried in vacuo. Depending on the product, further purification was carried out by distillation under reduced pressure or column chromatography on silica gel.

**General procedure for the synthesis of biphenyls with MnO<sub>2</sub> under single phase conditions (conditions B):** To a stirred suspension of the substituted benzene (20.0 mmol) and MnO<sub>2</sub> (435 mg, 5.00 mmol) in acetonitrile (5 mL) at r.t., a solution of the arylhydrazine (1.00 mmol) in acetonitrile (2 mL) was added dropwise over a period of 1 h. After completion of the reaction, as monitored by TLC, the reaction mixture was filtered over Celite. The filter cake was further washed with ethyl acetate and the solvents were removed under reduced pressure. The remaining benzene derivative was removed by Kugelrohr distillation, and the products were purified by silica gel chromatography.

**General procedure for the synthesis of biphenyls from arylhydrazine hydrochlorides under air (conditions C):** A mixture of the substituted benzene (20.0 mmol) and aqueous sodium hydroxide (1 N, 1.0 ml) was heated to 60–90 °C and the arylhydrazine hydrochloride was added portion wise in ten batches over a period of 9 hours. The reaction was completed after 24 hours at the given temperature, as monitored by TLC. After removal of water under reduced pressure, the remaining aniline was recovered by Kugelrohr distillation, and the crude biphenyls were purified by column chromatography on silica gel.

4'-Chlorobiphenyl-2-amine (**8a**) and 4'-chlorobiphenyl-4-amine (**8a''**) were prepared under the conditions A, B and C. Separation and purification by column chromatography (hexane / ethyl acetate 10:1) gave **8a** and **8a''** as dark solids. 4'-Chlorobiphenyl-2-amine (**8a**):  $R_f$  = 0.6 (hexane/EtOAc = 4:1) [UV]; mp = 65-67 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 4.09 (bs, 2 H), 6.80 (dd,  $J$  = 1.1 Hz,  $J$  = 8.0 Hz, 1 H), 6.85 (dt,  $J$  = 1.1 Hz,  $J$  = 7.5 Hz, 1 H), 7.10 (dd,  $J$  = 1.6 Hz,  $J$  = 7.6 Hz, 1 H), 7.15-7.21 (m, 1 H), 7.40 (s, 4 H); <sup>13</sup>C NMR (91 MHz, CDCl<sub>3</sub>) δ 115.7 (CH), 118.8 (CH), 126.3 (C<sub>q</sub>), 128.8 (CH), 129.0 (2 × CH), 130.3 (CH), 130.4

(2 × CH), 133.1 (C<sub>q</sub>), 137.9 (C<sub>q</sub>), 143.4 (C<sub>q</sub>); MS (EI) *m/z* (%): 205 (35) [<sup>37</sup>Cl-M<sup>+</sup>], 203 (100) [<sup>35</sup>Cl-M<sup>+</sup>], 168 (45), 167 (75); HRMS (ESI) calcd for C<sub>12</sub>H<sub>11</sub>ClN [M<sup>+</sup>+H]: 204.0575, found: 204.0569. 4'-Chlorobiphenyl-4-amine (**8a''**): R<sub>f</sub> = 0.3 (hexane/EtOAc = 4:1) [UV]; mp = 118-120 °C; <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>) δ 4.09 (bs, 2 H), 6.81 (d, *J* = 8.6 Hz, 2 H), 7.35 (d, *J* = 8.5 Hz, 2 H), 7.39 (d, *J* = 8.6 Hz, 2 H), 7.45 (d, *J* = 8.7 Hz, 2 H); <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 115.4 (2 × CH), 127.5 (2 × CH), 127.8 (2 × CH), 128.7 (2 × CH), 130.2 (C<sub>q</sub>), 132.1 (C<sub>q</sub>), 139.6 (C<sub>q</sub>), 146.1 (C<sub>q</sub>); MS (EI) *m/z* (%): 205 (30) [<sup>37</sup>Cl-M<sup>+</sup>], 203 (100) [<sup>35</sup>Cl-M<sup>+</sup>], 167 (18); HRMS (ESI) calcd for C<sub>12</sub>H<sub>11</sub>ClN [M<sup>+</sup>+H]: 204.0575, found: 204.0582. The spectral data obtained are in agreement with those reported in literature.<sup>15b</sup>

4'-Chloro-5-fluorobiphenyl-2-amine (**8b**) was prepared under the conditions A, B and C. Purification by column chromatography (hexane / ethyl acetate 10:1) gave **8b** as a dark brown oil. 4'-Chloro-5-fluorobiphenyl-2-amine (**8b**): R<sub>f</sub> = 0.6 (hexane/EtOAc = 4:1) [UV]; <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>) δ 3.54 (bs, 2 H), 6.69 (dd, *J*<sub>HF</sub> = 4.7 Hz, *J* = 8.5 Hz, 1 H), 6.83 (dd, *J* = 2.8 Hz, *J*<sub>HF</sub> = 9.0 Hz, 1 H), 6.88 (ddd, *J* = 3.0 Hz, *J*<sub>HF</sub> = 8.1 Hz, *J* = 8.7 Hz, 1 H), 7.37 (d, *J* = 8.7 Hz, 2 H), 7.42 (d, *J* = 8.7 Hz, 2 H); <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 115.3 (d, *J*<sub>CF</sub> = 22.2 Hz, CH), 116.5 (d, *J*<sub>CF</sub> = 22.6 Hz, CH), 116.7 (d, *J*<sub>CF</sub> = 7.7 Hz, CH), 127.4 (d, *J*<sub>CF</sub> = 7.2 Hz, C<sub>q</sub>), 129.1 (2 × CH), 130.3 (2 × CH), 133.6 (C<sub>q</sub>), 136.9 (d, *J*<sub>CF</sub> = 1.7 Hz, C<sub>q</sub>), 139.4 (d, *J*<sub>CF</sub> = 2.2 Hz, C<sub>q</sub>), 156.4 (d, *J*<sub>CF</sub> = 236.9 Hz, C<sub>q</sub>); MS(EI) *m/z* (%): 224 (6), 223 (29) [<sup>37</sup>Cl-M<sup>+</sup>], 222 (18), 221 (100) [<sup>35</sup>Cl-M<sup>+</sup>], 220 (10), 219 (20), 187 (8), 186 (45), 185 (60), 184 (13), 159 (5), 157 (7), 126 (6), 110 (10), 93 (37); HRMS (EI) calcd for C<sub>12</sub>H<sub>9</sub>ClFN [M<sup>+</sup>]: 221.0407, found: 221.0409. The spectral data obtained are in agreement with those reported in literature.<sup>15b</sup>

4'-Chloro-2-hydroxybiphenyl (**9a**) and 4'-chloro-4-hydroxybiphenyl (**9a''**) were prepared under the conditions B and C. Purification by column chromatography (hexane/EtOAc = 10:1



→ 4:1) gave a mixture of **9a** and **9a''**. 4'-Chloro-2-hydroxybiphenyl (**9a**):  $R_f = 0.5$  (hexane/EtOAc = 4:1) [UV];  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  5.06 (bs, 1 H), 6.95 (dd,  $J = 0.9$  Hz,  $J = 8.1$  Hz, 1 H), 6.99 (dt,  $J = 1.2$  Hz,  $J = 7.5$  Hz, 1 H), 7.21 (dd,  $J = 1.7$  Hz,  $J = 7.6$  Hz, 1 H), 7.23-7.29 (m, 1 H), 7.41 (d,  $J = 8.7$  Hz, 2 H), 7.44 (d,  $J = 8.6$  Hz, 2 H). 4'-Chloro-4-hydroxybiphenyl (**9a''**):  $R_f = 0.5$  (hexane/EtOAc = 4:1) [UV];  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  4.82 (bs, 1 H), 6.90 (d,  $J = 8.7$  Hz, 2 H), 7.37 (d,  $J = 8.7$  Hz, 2 H), 7.43 (d,  $J = 8.7$  Hz, 2 H), 7.45 (d,  $J = 8.7$  Hz, 2 H). The spectral data obtained are in agreement with those reported in literature.<sup>37,38</sup>

4'-Chloro-5-fluorobiphenyl-2-ol (**9b**) was prepared under the conditions B and C. Purification by column chromatography (dichloromethane 100%) gave **9b** as a light yellow oil. 4'-Chloro-5-fluorobiphenyl-2-ol (**9b**):  $R_f = 0.3$  (DCM) [UV];  $^1\text{H}$  NMR (360 MHz,  $\text{CDCl}_3$ )  $\delta$  4.70 (bs, 1 H), 6.74-6.78 (m, 1 H), 6.85 (dd,  $J = 3.1$  Hz,  $J_{\text{HF}} = 6.2$  Hz, 1 H), 7.01 (dd,  $J = 8.8$  Hz,  $J_{\text{HF}} = 10.0$  Hz, 1 H), 7.40 (d,  $J = 8.7$  Hz, 2 H), 7.46 (d,  $J = 8.6$  Hz, 2 H);  $^{13}\text{C}$  NMR (91 MHz,  $\text{CDCl}_3$ )  $\delta$  115.5 (d,  $J_{\text{CF}} = 8.2$  Hz, CH), 116.5 (d,  $J_{\text{CF}} = 3.1$  Hz, CH), 116.9 (d,  $J_{\text{CF}} = 25.0$  Hz, CH), 127.9 (d,  $J_{\text{CF}} = 7.7$  Hz,  $\text{C}_q$ ), 129.3 ( $2 \times$  CH), 130.3 ( $2 \times$  CH), 134.3 ( $\text{C}_q$ ), 134.6 ( $\text{C}_q$ ), 148.3 ( $\text{C}_q$ ), 156.0 (d,  $J_{\text{CF}} = 239.2$  Hz,  $\text{C}_q$ ); HRMS (EI) calcd for  $\text{C}_{12}\text{H}_9\text{ClFN}$  [ $\text{M}^+$ ]: 221.01779, found: 221.01749. The spectral data obtained are in agreement with those reported in literature.<sup>39</sup>

4'-Chloro-2-methoxybiphenyl (**10a**), 4'-chloro-3-methoxybiphenyl (**10a'**) and 4'-chloro-4-methoxybiphenyl (**10a''**) were prepared under the conditions A, B and C. Purification by column chromatography (hexane/EtOAc = 12:1 → 4:1) gave a mixture of **10a**, **10a'** and **10a''**. 4'-Chloro-2-methoxybiphenyl (**10a**):  $R_f = 0.2$  (hexane) [UV];  $^1\text{H}$  NMR (360 MHz,  $\text{CDCl}_3$ )  $\delta$  3.81 (s, 3 H), 6.98 (dd,  $J = 0.9$  Hz,  $J = 8.3$  Hz, 1 H), 7.02 (dt,  $J = 1.1$  Hz,  $J = 7.5$  Hz, 1 H), 7.28 (dd,  $J = 1.8$  Hz,  $J = 7.6$  Hz, 1 H), 7.30-7.34 (m, 1 H), 7.36 (d,  $J = 8.8$

Hz, 2 H), 7.46 (d,  $J = 8.8$  Hz, 2 H). *4'-Chloro-3-methoxybiphenyl* (**10a'**):  $R_f = 0.2$  (hexane) [UV];  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  3.86 (s, 3 H), 6.91 (ddd,  $J = 0.9$  Hz,  $J = 2.4$  Hz,  $J = 8.2$  Hz, 1 H), 7.08 (dd,  $J = 1.9$  Hz,  $J = 2.4$  Hz, 1 H), 7.14 (ddd,  $J = 0.9$  Hz,  $J = 1.9$  Hz,  $J = 7.6$  Hz, 1 H), 7.35 (d,  $J = 7.9$  Hz, 1 H), 7.40 (d,  $J = 8.7$  Hz, 2 H), 7.51 (d,  $J = 8.7$  Hz, 2 H). *4'-Chloro-4-hydroxybiphenyl* (**10a''**):  $R_f = 0.2$  (hexane) [UV];  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  3.85 (s, 3 H), 6.97 (d,  $J = 8.9$  Hz, 2 H), 7.37 (d,  $J = 8.7$  Hz, 2 H), 7.45-7.50 (m, 4 H). The spectral data obtained are in agreement with those reported in literature.<sup>40-42</sup>

*4'-Chloro-5-fluoro-2-methoxybiphenyl* (**10b**) and *4'-chloro-6-fluoro-3-methoxybiphenyl* (**10b'**) were prepared under the conditions A, B and C. Separation and purification by column chromatography (hexane / ethyl acetate 10:1) gave **10b** and **10b'** as a brown solids. *4'-Chloro-5-fluoro-2-methoxybiphenyl* (**10b**):  $R_f = 0.6$  (hexane/EtOAc = 4:1) [ $\text{KMnO}_4$ ];  $^1\text{H}$  NMR (360 MHz,  $\text{CDCl}_3$ )  $\delta$  3.78 (s, 3 H), 6.88-6.94 (m, 1 H), 6.97-7.04 (m, 2 H), 7.38 (d,  $J = 8.9$  Hz, 2 H), 7.46 (d,  $J = 8.8$  Hz, 2 H);  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  56.2 ( $\text{CH}_3$ ), 112.4 (d,  $J_{\text{CF}} = 8.3$  Hz, CH), 114.6 (d,  $J_{\text{CF}} = 22.6$  Hz, CH), 117.2 (d,  $J_{\text{CF}} = 23.5$  Hz, CH), 128.3 ( $2 \times \text{CH}$ ), 128.6 ( $\text{C}_q$ ), 130.7 ( $2 \times \text{CH}$ ), 133.4 ( $\text{C}_q$ ), 135.8 (d,  $J_{\text{CF}} = 1.6$  Hz,  $\text{C}_q$ ), 152.5 (d,  $J_{\text{CF}} = 8.1$  Hz,  $\text{C}_q$ ), 157.3 (d,  $J_{\text{CF}} = 240.4$  Hz  $\text{C}_q$ ); MS(EI)  $m/z$  (%): 237 (14), 236 (100), 221 (19), 186 (98), 157 (24); HRMS (EI) calcd for  $\text{C}_{13}\text{H}_{10}\text{ClFO}$  [ $\text{M}^+$ ]: 236.0404, found: 236.0404. *4'-Chloro-5-fluoro-2-methoxybiphenyl* (**10b'**):  $R_f = 0.5$  (hexane/EtOAc = 4:1) [ $\text{KMnO}_4$ ];  $^1\text{H}$  NMR (360 MHz,  $\text{CDCl}_3$ )  $\delta$  3.83 (s, 3 H), 6.83-6.87 (m, 1 H), 6.88-6.94 (m, 1 H), 7.08 (dd,  $J = 8.9$  Hz,  $J_{\text{HF}} = 10.0$  Hz, 1 H), 7.42 (d,  $J = 8.9$  Hz, 2 H), 7.47 (d,  $J = 8.8$  Hz, 2 H);  $^{13}\text{C}$  NMR (91 MHz,  $\text{CDCl}_3$ )  $\delta$  55.8 ( $\text{CH}_3$ ), 114.8 ( $\text{C}_q$ ), 115.3 (d,  $J_{\text{CF}} = 8.0$  Hz, CH), 116.8 (d,  $J_{\text{CF}} = 23.5$  Hz, CH), 120.4 (d,  $J_{\text{CF}} = 7.9$  Hz, CH), 129.2 ( $2 \times \text{CH}$ ), 130.4 ( $2 \times \text{CH}$ ), 133.9 ( $\text{C}_q$ ), 135.7 (d,  $J_{\text{CF}} = 1.6$  Hz,  $\text{C}_q$ ), 146.3 (d,  $J_{\text{CF}} = 20.2$  Hz,  $\text{C}_q$ ), 153.8 (d,  $J_{\text{CF}} = 240.4$  Hz,  $\text{C}_q$ ); MS(EI)  $m/z$  (%): 237 (8), 236 (14), 235 (4), 234 (9), 199 (5), 193 (7), 163 (9), 137 (6), 127 (5), 117 (5), 76 (6), 57 (4), 53 (6), 43 (4), 27 (6), 18 (100); HRMS (EI) calcd for  $\text{C}_{13}\text{H}_{10}\text{ClFO}$

[M<sup>+</sup>]: 236.0404, found: 236.0405. The spectral data obtained are in agreement with those reported in literature.<sup>15b</sup>

4'-Chlorobiphenyl-2-carbonitrile (**11a**), 4'-chlorobiphenyl-3-carbonitrile (**11a'**) and 4'-chlorobiphenyl-4-carbonitrile (**11a''**) were prepared under the conditions A, B and C. Purification by column chromatography (hexane/EtOAc = 10:1 → 4:1) gave a mixture of **11a**, **11a'** and **11a''**. 4'-Chlorobiphenyl-2-carbonitrile (**11a**): R<sub>f</sub> = 0.5 (hexane/EtOAc = 4:1) [UV]; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.44–7.51 (m, 6 H), 7.65 (dt, *J* = 1.4 Hz, *J* = 8.5 Hz, 1 H), 7.77 (ddd, *J* = 0.5 Hz, *J* = 1.4 Hz, *J* = 7.8 Hz 1 H); <sup>13</sup>C NMR (91 MHz, CDCl<sub>3</sub>) δ 111.2 (C<sub>q</sub>), 118.4 (CH), 123.4 (CH), 127.9 (CH), 129.9 (CH), 130.1 (2 × CH), 132.9 (2 × CH), 133.9 (C<sub>q</sub>), 135.0 (C<sub>q</sub>), 136.5 (C<sub>q</sub>), 144.2 (C<sub>q</sub>). 4'-Chlorobiphenyl-3-carbonitrile (**11a'**): R<sub>f</sub> = 0.5 (hexane/EtOAc = 4:1) [UV]; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.55 (t, *J* = 7.8 Hz, 1 H), 7.63–7.67 (m, 3 H), 7.73 (d, *J* = 0.9 Hz, *J* = 8.7 Hz, 2 H), 7.77 (d, *J* = 7.9 Hz, 1 H), 7.83 (t, *J* = 1.5 Hz, 1 H). 4'-Chlorobiphenyl-4-carbonitrile (**11a''**): R<sub>f</sub> = 0.5 (hexane/EtOAc = 4:1) [UV]; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.45 (d, *J* = 8.7 Hz, 2 H), 7.52 (d, *J* = 8.8 Hz, 2 H), 7.65 (d, *J* = 8.7 Hz, 2 H), 7.73 (d, *J* = 8.7 Hz, 2 H); <sup>13</sup>C NMR (91 MHz, CDCl<sub>3</sub>) δ 111.3 (C<sub>q</sub>), 118.7 (C<sub>q</sub>), 127.6 (2 × CH), 128.5 (2 × CH), 129.3 (2 × CH), 132.7 (2 × CH), 134.9 (C<sub>q</sub>), 137.6 (C<sub>q</sub>), 146.1 (C<sub>q</sub>). The spectral data obtained are in agreement with those reported in literature.<sup>43,44</sup>

4'-Chloro-5-fluorobiphenyl-2-carbonitrile (**11b**) and 4'-chloro-6-fluorobiphenyl-3-carbonitrile (**11b'**) were prepared under the conditions A, B and C. Purification by column chromatography (hexane/EtOAc = 10:1) gave a mixture of **11b** and **11b'**. 4'-Chloro-5-fluorobiphenyl-2-carbonitrile (**11b**): R<sub>f</sub> = 0.6 (Hexan/EtOAc = 4:1) [UV]; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.16 (ddd, *J* = 2.6 Hz, *J*<sub>HF</sub> = 7.8 Hz, *J* = 8.6 Hz, 1 H), 7.19 (dd, *J* = 2.6 Hz, *J*<sub>HF</sub> = 9.5 Hz, 1 H), 7.46–7.50 (m, 4 H), 7.77 (dd, *J*<sub>HF</sub> = 5.5 Hz, *J* = 8.6 Hz, 1 H); DEPTQ (151 MHz, CDCl<sub>3</sub>) δ 107.3 (C<sub>q</sub>), 115.5 (d, *J*<sub>CF</sub> = 22.5 Hz, CH), 117.2 (d, *J*<sub>CF</sub> = 23.2 Hz, CH), 117.

(C<sub>q</sub>), 129.2 (2 × CH), 129.9 (2 × CH), 135.5 (d,  $J_{CF}$  = 1.6 Hz, C<sub>q</sub>), 135.6 (C<sub>q</sub>), 136.2 (d,  $J_{CF}$  = 9.7 Hz, C<sub>q</sub>), 147.2 (d,  $J_{CF}$  = 9.1 Hz, C<sub>q</sub>), 164.8 (d,  $J_{CF}$  = 257.4 Hz, C<sub>q</sub>); GC-MS (EI)  $m/z$ : 233 ( $^{37}\text{Cl-M}^+$ ), 231 ( $^{35}\text{Cl-M}^+$ ), 196, 176, 169, 149, 97; HRMS (EI) calcd for C<sub>13</sub>H<sub>7</sub>ClFNNa: 254.0143, found: 254.0148. *4'-Chloro-6-fluorobiphenyl-3-carbonitrile (11b')*:  $R_f$  = 0.6 (Hexan/EtOAc = 4:1) [UV];  $^1\text{H}$  NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.26 (m, 1 H), 7.43-7.45 (m, 4 H), 7.64 (ddd,  $J$  = 2.2 Hz,  $J_{\text{HF}}$  = 4.5 Hz,  $J$  = 8.5 Hz, 1 H), 7.73 (dd,  $J_{\text{HF}}$  = 2.1 Hz,  $J$  = 7.1 Hz, 2 H).

*4'-Chloro-2-nitrobiphenyl (12a)*, *4'-chloro-3-nitrobiphenyl (12a')* and *4'-chloro-4-nitrobiphenyl (12a'')* were prepared under the conditions A, B and C. Purification by column chromatography (hexane/EtOAc = 15:1) gave a mixture of **12a**, **12a'** and **12a''**. *4'-Chloro-2-nitrobiphenyl (12a)*:  $R_f$  = 0.6 (hexane/EtOAc = 4:1) [UV];  $^1\text{H}$  NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.16–7.19 (m, 2 H), 7.31–7.34 (m, 3 H), 7.43 (dt,  $J$  = 1.2 Hz,  $J$  = 8.0 Hz, 1 H), 7.54 (td,  $J$  = 1.2 Hz,  $J$  = 7.6 Hz 1 H), 7.80 (dd,  $J$  = 1.2 Hz,  $J$  = 8.0 Hz 1 H). *4'-Chloro-3-nitrobiphenyl (12a')*:  $R_f$  = 0.6 (hexane/EtOAc = 4:1) [UV];  $^1\text{H}$  NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.45 (d,  $J$  = 7.6 Hz, 2 H), 7.55 (d,  $J$  = 7.6 Hz, 2 H), 7.61 (t,  $J$  = 8.6 Hz, 1 H), 7.87 (d,  $J$  = 7.6 Hz, 1 H), 8.20 (m, 1 H), 8.39 (t, 1 H). *4'-Chloro-4-nitrobiphenyl (12a'')*:  $R_f$  = 0.6 (hexane/EtOAc = 4:1) [UV];  $^1\text{H}$  NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.47 (d,  $J$  = 8.4 Hz, 2 H), 7.55 (d,  $J$  = 8.4 Hz, 2 H), 7.70 (d,  $J$  = 8.4 Hz, 2 H), 8.29 (d,  $J$  = 8.4 Hz, 2 H);  $^{13}\text{C}$  NMR (91 MHz, CDCl<sub>3</sub>)  $\delta$  124.2 (2 × CH), 127.6 (2 × CH), 128.6 (2 × CH), 129.4 (2 × CH), 135.3 (C<sub>q</sub>), 137.2 (C<sub>q</sub>), 146.3 (C<sub>q</sub>), 147.3 (C<sub>q</sub>). The spectral data obtained are in agreement with those reported in literature.<sup>44-46</sup>

**Acknowledgment.** The authors would like to thank the Deutsche Forschungsgemeinschaft for financial support within the projects HE5413/2-2 and GRK 1910/A6 and B3.

**Supporting Information Available.** Computational results of the density-functional calculations, tables of atom coordinates and absolute energies, NMR spectra of biaryl compounds and GC chromatograms of biaryl compounds **10b** and **10b'**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Gomberg, M.; Bachmann, W. E. *J. Am. Chem. Soc.* **1924**, *46*, 2339.
- (2) For reviews and mechanistic studies on the Gomberg-Bachmann reaction and related radical aryl-aryl coupling reactions, see: (a) Dermer, O. C.; Edmison, M. T. *Chem. Rev.* **1957**, *57*, 77. (b) Rüchardt, C.; Merz, E. *Tetrahedron Lett.* **1964**, *36*, 2431. (c) Bolton, R.; Williams, G. H. *Chem. Soc. Rev.* **1986**, *15*, 261. (d) Studer, A.; Curran, D. P. *Nature Chem.* **2014**, *6*, 765. (e) Studer, A.; Curran, D. P. *Angew. Chem. Int. Ed.* **2011**, *50*, 5018.
- (3) For reviews on radical aryl-aryl coupling reactions, see: (a) Studer, A.; Bossart, M. in *Radicals in Organic Synthesis*, eds. Renaud, P.; Sibi, M. 1st ed., Wiley-VCH **2001**, vol. 2, 62. (b) Bowman, W. R.; Storey, J. M. D. *Chem. Soc. Rev.* **2007**, *36*, 1803. (c) Studer, A.; Curran, D. *Angew. Chem. Int. Ed.* **2011**, *50*, 5018. (d) Pratsch, G.; Heinrich, M. R. in *Radicals in Synthesis III, Topics in Current Chemistry*, eds. Gansäuer, A.; Heinrich, M. R., Springer **2012**, vol. 320. (e) Bonin, H.; Sauthier, M.; Felpin, F.-X. *Adv. Synth. Catal.* **2014**, *356*, 645.
- (4) Galli, C. *Chem. Rev.* **1988**, *88*, 765.
- (5) For the most advanced and versatile Gomberg-Bachmann reactions under classical conditions, see: Baedle, J. R.; Korzeniowski, S. H.; Rosenberg, D. E.; Garcia-Slanga, B. J.; Gokel, G. W. *J. Org. Chem.* **1984**, *49*, 1594.
- (6) For reviews on aromatic C-H activation reactions, see: (a) Dyker, G. *Handbook of C-H Transformations*, Wiley-VCH, Weinheim, **2005**. (b) Ackermann, L. *Top. Organomet.*

- Chem.* **2008**, *24*, 35. (c) Li, B.-J.; Yang, S.-D.; Shi, Z.-J. *Synlett* **2008**, 949. (d) Lersch, M.; Tilset, M. *Chem. Rev.* **2005**, *105*, 2471. (d) Gorelsky, S. I. *Coord. Chem. Rev.* **2013**, *257*, 153. (e) Cacchi, S.; Fabrizi, G.; Goggiamani, A. in *Innovative Catalysis in Organic Synthesis* **2012**, 211. (f) Chen, D. Y.-K.; Youn, S. W. *Chem. Eur. J.* **2012**, *18*, 9452.
- (7) (a) Hari, D. P.; Schroll, P.; König, B. *J. Am. Chem. Soc.* **2012**, *134*, 2958. (b) Zoller, J.; Fabry, D. C.; Rueping, M. *ACS Catal.* **2015**, *3*, 3900. (c) Xue, D.; Jia, Z.-H.; Zhao, C.-J.; Zhang, Y.-Y.; Wang, C.; Xiao, J. *Chem. Eur. J.* **2014**, *20*, 2960.
- (8) Base-mediated arylation: (a) Yanagisawa, S.; Ueda, K.; Taniguchi, T.; Itami, K. *Org. Lett.* **2008**, *10*, 4673. (b) Sun, C.-L.; Li, H.; Yu, D.-G.; Yu, M.; Zhou, X.; Lu, X.-Y.; Huang, K.; Zheng, S.-F.; Li, B.-J.; Shi, Z.-J. *Nat. Chem.* **2010**, *2*, 1044. (c) Shirakawa, E.; Itoh, K.-I.; Higashino, T.; Hayashi, T. *J. Am. Chem. Soc.* **2010**, *132*, 15537. (d) Liu, W.; Cao, H.; Zhang, H.; Chung, K. H.; He, C.; Wang, H.; Kwong, F. Y.; Lei, A. *J. Am. Chem. Soc.* **2010**, *132*, 16737. (e) Dewanji, A.; Murarka, S.; Curran, D.; Studer, A. *Org. Lett.* **2013**, *15*, 6102. (f) Budén, M. E.; Guastavino, J. F.; Rossi, R. A. *Org. Lett.* **2013**, *15*, 1174.
- (9) (a) Demir, A. S.; Findik, H. *Tetrahedron* **2008**, *64*, 6196. (b) Chen, Z.-X.; Wang, G.-W. *J. Org. Chem.* **2005**, *70*, 2380. (c) Ravi, M.; Chauhan, P.; Kant, R.; Shukla, S. K.; Yadav, P. P. *J. Org. Chem.* **2015**, *80*, 5369.
- (10) Crisóstomo, F. P.; Martín, T.; Carrillo, R. *Angew. Chem. Int. Ed.* **2014**, *53*, 2181.
- (11) For the rate of addition of aryl radicals to benzene, see: Scaiano, J. C.; Stewart, L. C. *J. Am. Chem. Soc.* **1983**, *105*, 3609.
- (12) For the so far most detailed overview, see: M. J. Perkins in J. K. Kochi: Free Radicals. Vol. 2, Wiley, New York **1973**, p. 231.
- (13) (a) Wetzel, A.; Pratsch, G.; Kolb, R.; Heinrich, M. R. *Chem. Eur. J.* **2010**, *16*, 2547. (b) Maity, P.; Kundu, D.; Ranu, B. C. *Eur. J. Org. Chem.* **2015**, 1727.

- (14) (a) Pratsch, G.; Wallaschkowski, T.; Heinrich, M. R. *Chem. Eur. J.* **2012**, *18*, 11555. (b) Höfling, S. B.; Bartuschat, A. L.; Heinrich, M. R. *Angew. Chem. Int. Ed.* **2010**, *49*, 9769.
- (15) (a) Jasch, H.; Scheumann, J.; Heinrich, M. R. *J. Org. Chem.* **2012**, *77*, 10699. (b) Hofmann, J.; Jasch, H.; Heinrich, M. R. *J. Org. Chem.* **2014**, *79*, 2314. (c) Jiang, T.; Chen, S.-Y.; Zhang, G.-Y.; Zeng, R.-S.; Zou, J.-P. *Org. Biomol. Chem.* **2014**, *12*, 6922. (d) Jiang, T.; Chen, S.-Y.; Zhuang, H.; Zeng, R.-S.; Zou, J.-P. *Tetrahedron Lett.* **2014**, *55*, 4549.
- (16) Hey, D. H. in *Advances in Free Radical Chemistry*, Williams, G. H., Ed.; Logos-Academic, London, **1967**, Vol. II, p. 47.
- (17) (a) Dannley, R. L.; Sternfeld, M. *J. Am. Chem. Soc.* **1954**, *76*, 4543. (b) Cadogan, J. I. G.; Hey, D. H.; Williams, G. H. *J. Chem. Soc.* **1955**, 1425.
- (18) Ito, R.; Migita, T.; Morikawa, N.; Simamura, O. *Tetrahedron* **1965**, *21*, 955.
- (19) Chlorophenyl radicals have shown a comparably low regioselectivities in arylations of such diverse arenes as PhNO<sub>2</sub>, PhCl, PhOMe, PhMe. See ref. 12.
- (20) Giese, B. *Angew Chem. Int. Ed.* **1983**, *22*, 753.
- (21) Curran, D. P.; Keller, A. I. *J. Am. Chem. Soc.* **2006**, *128*, 13706.
- (22) Petrillo, G.; Novi, M.; Dell'Erba, C.; Tavani, C.; Berta, G. *Tetrahedron* **1990**, *46*, 7977.
- (23) DeTar, D. F.; Scheifele, H. J. *J. Am. Chem. Soc.* **1951**, *73*, 1442.
- (24) Shih, C.; Hey, D. H.; Williams, G. H. *J. Chem. Soc.* **1958**, 1885.
- (25) Reactivity series derived from average yields from Tables 1-3: aniline (64%), phenol/phenolate (59%), benzonitrile (50%), anisole (41%), nitrobenzene (39%).
- (26) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, M.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.;

Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A. J.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Keith, T. A.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, O.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian 09, Revision C.01, Revision C.01; Gaussian, Inc.: Wallingford CT, **2010**.

(27) Becke, A. D., *J. Chem. Phys.* **1993**, *98*, 5648.

(28) Lee, C. T.; Yang, W. T.; Parr, R. G., *Phys. Rev. B* **1988**, *37*, 785.

(29) Dunning, T. H. Jr., *J. Chem. Phys.*, **1989**, *90*, 1007.

(30) Kendall, R. A., Dunning, T. H. Jr., Harrison, R. J., *J. Chem. Phys.*, **1992**, *96*, 6796.

(31) G. C. Tiu, F.-M. Tao, *Chem. Phys. Lett.* **2006**, *428*, 42.

(32) (a) James, D. G. L.; Suart, R. D. *Chem. Commun.* **1966**, 484. (b) James, D. G. L.; Suart, R. D. *Trans. Faraday Soc.* **1968**, *64*, 2735 and 2752.

(33) (a) De Tar, D. F. *J. Am. Chem. Soc.* **1967**, *89*, 4058. (b) See also ref. 12.

(34) Zhang, X. *J. Quantum Chem.* **2015**, *115*, 1658.

(35) Anglada, J. M.; Besalú, E.; Bofill, J. M.; Crehuet, R. *J. Comput. Chem.* **1999**, *20*, 1112 and 1130.

(36) Atkinson, D. J.; Perkins M. J.; Ward, P. *J. Chem. Soc. C* **1971**, 3240.

(37) Hoshiya, N.; Shimoda, M.; Yoshiyuki, H.; Yamashita, Y.; Shuto, S.; Arisawa, M. *J. Am. Chem. Soc.* **2010**, *132* (21), 7270-7272.



- (38) Gallon, B. J.; Kojima, R. W.; Kaner, R. B.; Diaconescu, P. L. *Angew. Chem., Int. Ed.* **2007**, *46*, 7251-7254.
- (39) Kralj, A.; Kurt, E.; Tschammer, N.; Heinrich, M R. *ChemMedChem*, **2014**, *9*, 151-168.
- (40) Denmark, S. E.; Smith, R. C.; Chang, W.-T. T.; Muhuhi, J. M. *J. Am. Chem. Soc.* **2009**, *131*, 3104.
- (41) Lu, J.-M.; Ma, H.; Ma, D.; Shao, L.-X.; Li, S.-S. *Tetrahedron* **2010**, *66*, 5185.
- (42) Zhou, W.-J.; Wang, K.-H.; Wang, J.-X.; Gao, Z.-R. *Tetrahedron* **2010**, *66*, 7633.
- (43) Zhang, L.; Ang, G. Y.; Chiba, S. *Org. Lett.* **2010**, *12*, 3682-3685.
- (44) Xu, C.; Yin, L.; Huang, B.; Liu, H.; Cai, M. *Tetrahedron* **2016**, *72*, 2065-2071.
- (45) Elumalai, V.; Sandtorv, A. H.; Bjørsvik, H.-R. *Eur. J. Org. Chem.* **2016**, *7*, 1344-1354.
- (46) Al-Amin, M.; Akimoto, M.; Tameno, T.; Ohki, Y.; Takahashi, N.; Hoshiya, N.; Shuto, S.; Arisawa, M. *Green Chem.* **2013**, *15*, 1142-1145.