



Cite this: DOI: 10.1039/c6tb02456d

Fluorescent flavonoids for endoplasmic reticulum cell imaging†

Lucas McDonald,^a Bin Liu,^{‡a} Alexandra Taraboletti,^a Kyle Whiddon,^a Leah P. Shriver,^{ab} Michael Konopka,^a Qin Liu^b and Yi Pang^{*a}

Visualization of subcellular organelles *in vivo* is critical for basic biomedical research and clinical applications. Two new flavonoids with an amide substituent were synthesized and characterized. The flavonoids were nearly non-fluorescent in aqueous environment, but exhibited two emission peaks (one λ_{em} at 495–536 nm and the other at 570–587 nm) in organic solvents, which were assigned to the excited normal (N*) and tautomer (T*) emission. When the dyes were examined on oligodendrocyte cells, they were found to selectively accumulate in the endoplasmic reticulum (ER), a eukaryotic organelle involved in lipid and protein synthesis, giving fluorescence turn-on. The ER-selective flavonoids could be a valuable tool due to its low molecular mass (<500), large Stokes' shift, low toxicity, and biocompatibility.

Received 20th September 2016,
Accepted 16th November 2016

DOI: 10.1039/c6tb02456d

www.rsc.org/MaterialsB

Introduction

Optical visualization of subcellular organelles and biochemical processes *in vivo* is critical for basic biomedical research and the development of novel clinical diagnostics.^{1–3} The endoplasmic reticulum (ER) plays a central role in the biosynthesis of lipids and proteins. In the lumen of the endoplasmic reticulum, proteins fold and mature before being delivered to other compartments of cells or released extracellularly. An accumulation of unfolded or misfolded proteins causes disturbances in the normal functions of the ER, which induces a cellular stress response (or ER stress) that activates the unfolded protein response (UPR).^{1,2} UPR activation increases ER abundance to match needs by mediating expansion of the ER membrane.³ Since ER stress is involved in the feedback loops that establish and maintain homeostasis, it plays a significant role in various diseases⁴ such as autoimmunity⁵ and metabolic disease.^{6,7} Because of the importance of the ER in disease, there is significant interest in developing new probes to track ER dynamics.⁸ An effective ER imaging probe has potential to identify perturbations in ER function that are associated with early disease pathology.

Oligodendrocytes are the myelinating glial cells of the central nervous system. The myelin sheath produced by these cells is a lipid-rich structure that insulates axons in the central

nervous system (CNS) and promotes rapid conduction of nerve impulses.⁹ Myelin is a poorly hydrated structure containing 40% water, and the dry mass of myelin consists of ~70% lipids and 30% proteins. During the active phase of myelination (the process of forming myelin sheath), each oligodendrocyte in the CNS must produce as much as ~5000 μm^2 of myelin surface area per day, and ~ 10^5 myelin-associated protein molecules per minute.¹⁰ ER stress can disturb lipid and protein synthesis and this has been linked to oligodendrocyte perikaryon degeneration and myelin sheath disintegration.¹¹ In addition, ER stress in oligodendrocytes has been implicated in dysmyelinating diseases.¹²

Considerable interests exist in searching for novel fluorescent probes that can target specific biological tissues/molecules to meet the need for advanced bioimaging.¹³ Among the known probes, few are capable of targeting ER, partly due to the challenge in controlling ER selectivity. A recent ER-probe uses a nitro-furan group to target ER-associated proteins.¹⁴ The commercial ER-Tracker Green and Red contains a “gliberclamide” segment (Scheme 1), which binds to the sulphonylurea receptors of ATP-sensitive K^+ channels^{15,16} that are prominent on ER.¹⁷ In the commercial ER probes, the necessary “gliberclamide” segment for ER selectivity has the mass of 494 Da, whose relative large moiety could cause significant perturbations to normal cell activity upon introduction. In order to avoid this potential side effect, an ideal ER probe should have the lowest possible molecular weight (e.g. Mw below 500 Da). In addition, the probe should be non-toxic for cells.


As a broad class of natural substances normally occurring in diet, flavonoids exhibit a variety of beneficial health effects, including antioxidant properties and anticancer activities.^{18,19} With simple/minor structural modification, the flavonoids can

^a Department of Chemistry, University of Akron, Akron, Ohio 44325, USA.
E-mail: yp5@uakron.edu


^b Department of Biology, University of Akron, Akron, Ohio 44325, USA

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c6tb02456d

‡ Current address: Shenzhen Key Laboratory of Special Functional Materials, College of Materials Science and Engineering, Shenzhen University, Shenzhen 518060, China.



Scheme 1 Structures of commercial ER-Tracker, where the fluorophore is shown in blue color.



Scheme 2 Chemical structures of flavonoids **1–2** and synthesis of **2**.

also exhibit “excited-state intramolecular photon-transfer (ESIPT),” which provides intense fluorescence with large Stokes Shifts.²⁰ The combination of the unique optical properties and biocompatibility makes flavonoid compounds a valuable template for sensor design. Our recent studies show that the flavonoid **1** (Scheme 2) with a suitable amino substituent (R_2N) is an environmentally sensitive dye, which can be tailored to target various biomolecular structures.^{21,22} Thus, *via* attaching a functional group at the C-3 position of “chromen-4-one” segment, flavonoid **1a** can label mitochondria.²¹ Flavonoid **1b** shows a large fluorescence turn-on upon selective binding to albumin proteins.²² The observed protein selectivity also raises the possibility for other protein binding-induced fluorescence turn-on. Herein we disclose that the flavonoid **2** with a small amide group on the C-6 position shows excellent selectivity for ER localization in mammalian cells.

Results and discussion

Synthesis of flavonoids

The new flavonoids **2a–2b** include a donor–acceptor (D–A) interaction, which retains the desirable fluorescent solvatochromic property. Flavonoids **2** were synthesized by reaction of *N*-(3-acetyl-4-hydroxyphenyl)butyramide **3** with 4-(dialkylamino)benzaldehyde **4** in two steps (Scheme 2), *i.e.* by Claisen–Schmidt condensation followed by Algar–Flynn–Oyamada reaction.^{23,24} The crude

product was purified by recrystallization from hexane/ethanol mixture in good yield.

Optical properties

Both flavonoid compounds showed a similar absorption, revealing λ_{\max} at 406 nm and 411 nm for **2a** and **2b**, respectively (Fig. 1). The emission of **2a** revealed two peaks with λ_{em} at 495 and 570 nm, which could be attributed to the excited normal form (N^*) and the tautomeric form (T^*), respectively. The tautomer emission (from T^*) was associated with the excited state intramolecular proton transfer (ESIPT) (see Scheme 3).²⁵ In contrast, the emission λ_{em} from the N^* and T^* of **2b** was at 536 and 587 nm, with about equal intensity. The observed large difference in the ratio of N- and T-emission indicated that the amino substituent in the electron-donor segment had a profound impact on the emissive property of flavonoids. By using rhodamine 6G in ethanol (QY = 95%)²⁶ as a reference, the fluorescence quantum yield (QY) of **2a** and **2b** were determined to be 26% and 47%, respectively. Both **2a** and **2b** exhibited large Stokes' shifts (~ 170 nm), which is highly desirable for fluorescence imaging. The observed large Stokes' shift could be attributed to a combination of the intramolecular charge transfer (ICT) and ESIPT processes occurring in the flavonoids.

Compound **2** appeared to be non-fluorescent in aqueous solution, possibly due to excessive hydrogen bonding to chromen-4-one unit (see **1** in Scheme 2). In a non-aqueous solution, however, the flavonoid gave a large fluorescence turn-on (Fig. 2). In a less polar solvent such as THF and toluene, the flavonoid gave a





Fig. 1 UV-vis (broken line) and fluorescence spectra (solid line) of **2a** and **2b** in CH_2Cl_2 (10 μM) at room temperature. The fluorescence excitation at 410 nm. The N^* and T^* indicate the emission from normal or tautomeric form.



Scheme 3 The ESIPT process of a representative flavonoid, showing the structural change from N-tautomer to T-tautomer.




Fig. 2 Fluorescence of **2b** in different solvents at room temperature.

predominant emission from the excited tautomer (T^*). It was also interesting to note that the T^* emission wavelength was basically not influenced by the solvent polarity, although the tautomer had a more polar structure. This was in sharp contrast to the N^* emission, whose emission wavelength was notably shifted to a longer wavelength as solvent polarity increased (e.g. from 487 nm in toluene to 540 nm in DMSO, see Fig. 2). One possible reason was that the solvent reorganization was not significant during the short lifetime of the excited tautomer T^* .

The environmental response of **2** was further examined on proteins. The weak fluorescence of **2b** in aqueous solution was increased moderately upon addition of 1 equiv. of bovine serum albumin (BSA) or human serum albumin (HSA), giving only one emission peak at $\lambda_{em} \approx 532$ nm (ESI,† Fig. S2). On the basis of the emission wavelength (532 nm), the observed emission from the albumin binding was likely from the N^* form (Fig. 2), although it was not clear why the emission from the T^* form was inhibited upon binding proteins.

Cytotoxicity of flavonoids was examined by using an MTT assay, showing that the half maximal inhibitory concentration (IC_{50}) for **2a–2b** was 25 μ M. Low cytotoxicity of **2** encouraged us to further examine the staining of the flavonoids in MO3.13 cells. This human cell line has the properties of myelinating oligodendrocytes, and has abundant myelin basic proteins (MBP) which are almost exclusively found in oligodendrocyte.²⁷ When the cells were incubated with **2** and imaged by confocal microscopy, the dye was found to quickly penetrate into cells, giving strong fluorescence (Fig. 3). Weak fluorescence could be observed immediately after adding dyes into cells and mounting the sample onto microscope (approximately in less than 1 minute). Since the dye was relatively non-fluorescent in an aqueous environment, the observed fluorescence was assumed to arise from its interaction with cellular lipids or hydrophobic pockets of proteins.

Colocalization of **2** and ER-Tracker in oligodendrocytes

To the MO3.13 cell media were added ER-Tracker™ Red (concentration 1 μ M) along with flavonoid **2** to be tested (concentration 500 nM). Fluorescent probe DRAQ5 (10 μ M) was also added to the tissue culture media for staining cell nuclei. The fluorescence




Fig. 3 Confocal imaging of oligodendrocyte cells incubated with **2b** (10 μ M) at <1, 15, 30, and 60 minutes after probe addition. Magnification is 100 \times .

confocal imaging (Fig. 4B) revealed that flavonoid **2** stained the cell components surrounding the nuclei. The sharp circles in fluorescence image of **2** (Fig. 3) were outer contour of nuclei, since they are not stained and non-fluorescent.

The non-uniform stain of **2** on the cells suggests that the dyes might be selectively binding to intracellular organelles. In order to confirm the intracellular location and distribution of **2**, we examined the colocalization of our dye with the commercial ER probe, ER-Tracker™ Red. Incubation of MO3.13 cells with ER-Tracker™ Red showed robust fluorescence in structures surrounding the nucleus, a pattern consistent with ER staining (Fig. 4A and D). Staining with both compounds **2a** and **2b** showed similar localization (Fig. 4E and B) and colocalization was observed when cells were incubated with both the commercial ER dye and our probes (Fig. 4C and F). The stained cells were scanned repeated at 5 minute interval for 3 hours (ESI,† Fig. S12), which revealed good stability in comparison with commercial ER Tracker (ESI,† Fig. S13).

It is well known that ER forms an interconnected network of membrane-enclosed sacs which are embedded with transmembrane proteins that contain hydrophobic structural motifs. It is attractive to speculate that the flavonoid dyes were binding to these types of hydrophobic structures. Such ER binding-induced fluorescence turn-on would be valuable for dynamic tracking, since **2** was nearly non-fluorescent in the aqueous environment. The assumption was confirmed by observation of clear cell contours without post-staining washing (Fig. 3), as the dyes outside the cells did not give observable fluorescence. Direct comparison also revealed that flavonoid **2b** gave stronger green fluorescence than **2a** under the same experimental conditions (Fig. 4B and E), illustrating that the ER binding-induced fluorescence turn on was also sensitive to substituents on the flavonoid.




Fig. 4 Confocal imaging of MO3.13 cells incubated with ER-Tracker (1 μM) (panels A and D), flavonoid compounds **2b** (panel B) or **2a** (panel E) (500 nM, for each dye respectively). Panels (C) and (F) are the merged images of ER Tracker Red and flavonoid probes. The nuclei of cells were stained with DRAQ5 (shown in purple).

In order to verify the ER selectivity, flavonoid **2b** was further examined by staining in GL261 cells (astrocyte derived glioblastoma), which is a mouse glioblastoma multiforme (GBM) cell line and its growth is known to respond to ER stress.²⁸

Colocalization of **2b** with the commercial ER-Tracker™ Red revealed the identical patterns (Fig. 5A and B), showing that the flavonoid was indeed selective to ER. The ER selectivity was also examined on SIM-A9 cells (semi adherent microglial cells).²⁹




Fig. 5 Confocal imaging of GL261 cells (panels A–C) and SIM A9 cells (panels D–F), which were incubated with ER-Tracker (1 μM, panels A and D), flavonoid **2b** (500 nM, panels B and E) for 30 minutes. Panels (C) and (F) are the merged images of ER Tracker Red and flavonoid probes. The nuclei of cells were stained with DRAQ5 (shown in purple).




Fig. 6 (A) Confocal image of zebrafish (48 hpf) at 10 \times magnification, after staining with flavonoid dye **2b**. The images were taken from the head and tail sections. (B) and (C) Fluorescence confocal images of zebrafish from the tail section (approximate position is shown by a red square), at 20 \times and 100 \times magnification, respectively.

As a type of glial cell located throughout the brain, microglial cells account for 10–15% of all cells found within the brain,³⁰ and ER stress plays a significant role in the microglial cell death.³¹ Although the ER distribution in GL261 cells is quite different from that in SIM-A9 cells (Fig. 5A and D), colocalization of **2b** with the commercial ER-Tracker™ Red revealed the same patterns.

Zebrafish staining

The ability of **2b** to penetrate cells (Fig. 3), in addition to its low toxicity, encouraged us to further examine its potential *in vivo* applications. When zebrafish of 48 hours post fertilization (hpf) was exposed to **2b**, bright green fluorescence could be observed from the entire zebrafish (Fig. 6), indicating that flavonoid dye could easily penetrate into the fish for *in vivo* applications. Interestingly, the detailed cell structure could be clearly visible from the stained zebrafish (Fig. 6C), as the penetrated flavonoid dye did not exhibit fluorescence in aqueous solution.

Experiment section

Synthesis of **2b**

N-(3-Acetyl-4-hydroxyphenyl)butyramide (10 mmol) was added to a solution of 4-(diphenylamino)benzaldehyde (10 mmol) in ethanol (50 mL), then 50 mL of aqueous NaOH (4 g, 100 mmol) solution was added slowly to the solution at 0 °C over 1 hour period. The mixture was stirred at reflux (~80 °C) for 6 hours, and then cooled to room temperature for another 48 hours. H₂O₂ solution (10 mL of 30%) was slowly added into the reaction solution over 1 hour while cooled in an ice-water bath. After stirring at room temperature for 96 hours, the mixture was poured into ice water and then placed into the refrigerator overnight. The precipitate was collected *via* filtration, and

washed with cold ethanol. The product was purified by recrystallization from ethanol three times. Yield = 17%. ¹H NMR (*d*₆-DMSO, 500 MHz): δ = 10.15 (s, 1H, -OH), 9.35 (s, 1H, -NH), 8.43 (d, 1H, *J* = 3.0 Hz), 8.09 (d, 2H, *J* = 9.0 Hz), 7.90 (dd, 1H, *J*₁ = 9.0 Hz, *J*₂ = 2.5 Hz), 7.66 (d, 1H, *J* = 9.0 Hz), 7.38 (t, 4H), 7.16 (m, 6H), 7.04 (d, 2H, *J* = 9.0 Hz), 2.37 (t, 2H), 1.65 (m, 2H), 0.95 (t, 3H). ¹³C NMR (*d*₆-DMSO, 125 MHz): *d* = 172.76, 171.80, 150.88, 149.06, 146.85, 145.92, 138.44, 136.32, 130.23, 129.39, 125.80, 124.65, 124.50, 121.90, 121.19, 119.13, 113.23, 38.77, 18.96, 14.07. MS, calcd for [C₃₁H₂₆N₂O₄ + Na]⁺ *m/z*: 513.2; found: 513.2.

Synthesis of **2a**

Compound **2a** was prepared from *N*-(3-acetyl-4-hydroxyphenyl)-butyramide (50 mmol) and 4-(dimethylamino)benzaldehyde (50 mmol) in ethanol (100 mL) under the same conditions used for **2b**. The product was purified by recrystallization from hexane/ethanol (v/v = 1/1). Yield = 21%. ¹H NMR (*d*₆-DMSO, 500 MHz): δ = 10.13 (s, 1H, -OH), 9.07 (s, 1H, -NH), 8.40 (d, 1H, *J* = 2.5 Hz), 8.11 (d, 2H, *J* = 9.0 Hz), 7.89 (dd, 1H, *J*₁ = 9.0 Hz, *J*₂ = 2.5 Hz), 7.67 (d, 1H, *J* = 9.0 Hz), 6.85 (d, 2H, *J* = 9.0 Hz), 3.00 (s, 6H), 2.34 (t, 2H), 1.66 (m, 2H), 0.95 (t, 3H). ¹³C NMR (*d*₆-DMSO, 125 MHz): *d* = 157.43, 157.02, 139.20, 138.55, 135.40, 126.91, 125.79, 119.78, 116.29, 113.27, 110.68, 110.24, 105.65, 104.43, 41.42, 40.27, 22.88, 18.58. ESI-MASS *m/z* calcd for [C₂₁H₂₂N₂O₄ + H]⁺: 367.2; found: 367.2.

Fluorescence quantum yield

The fluorescence quantum yields (QY) were measured using rhodamine 6G (sigma) as the standard ($\Phi_{\text{R}} = 0.95$, ethanol).²⁶ The fluorescence quantum yields were calculated by using the following equation:

$$\Phi_{\text{s}} = \Phi_{\text{r}} \times (A_{\text{r}} \times n_{\text{s}}^2 \times F_{\text{s}}) / (A_{\text{s}} \times n_{\text{r}}^2 \times F_{\text{r}})$$

where the subscripts s and r refer to the sample and the standard (rhodamine 6G), respectively. Φ is the quantum yield, *F* is the integrated emission intensity, *A* is the absorbance, and *n* is the refractive index of the solution.

Live imaging

Imaging was performed on a Nikon A1 confocal system with a 100 \times Plan Apo λ , NA = 1.45 oil objective with both GaAsP detectors and high sensitivity low noise PMTs for detection. The excitation used for our dyes was 405 nm with standard DAPI, FITC, and Texas Red filters. The ER-Tracker™ was excited using 560 nm and with a 600/50 nm bandpass filter used for emission. The nuclear dye, DRAQ5, was excited at 620 nm and 680/75 nm bandpass filter was used for emission. All imaging was done in an Okolab Bold Cage Incubator at 37 °C, and images were processed using NIS Elements or ImageJ Pro imaging software.

Cell culturing and staining

MO3.13 cells (oligodendrocytes), GL261 cells (astrocyte derived glioblastoma) or SIM A9 cells (semi adherent microglial) were plated on microscopy dishes at a density of 5 \times 10⁵ cells per mL

in DMEM media supplemented with 10% FBS and 1% penicillin/streptomycin. Cells were allowed to attach overnight at 37 °C and in 5% CO₂. After cells attached, ER-Tracker™ Red (Sigma E34250) was added to the cell media at a final concentration of 1 μM along with a final concentration at 500 nM of the flavonoid dye to be tested. Cells were left to incubate for 2 hours, and then washed 5 times with PBS. Live Cell Imaging Solution (Thermo-Fisher A14291DJ) was then added, along with 10 μM of DRAQ5 (Sigma 62251) fluorescent probe, to stain cell nuclei. Cells were then left to incubate at room temperature for 30 minutes before imaging.

Zebrafish breeding and staining

Zebrafish were maintained as described in the Zebrafish Book by the University of Oregon. They were kept in 28.5 °C water for mating, male and female zebrafish were maintained in one tank on a 12 h light/12 h dark cycle. Spawning was triggered by light stimulation in the morning and eggs were immediately fertilized. All embryos were maintained in E3 medium (15 mM NaCl, 0.5 mM KCl, 1 mM MgSO₄, 1 mM CaCl₂, 0.15 mM KH₂PO₄, 0.05 mM Na₂HPO₄, 0.7 mM NaHCO₃, and 10⁻⁵% methylene blue at 7.5 pH). All animal related procedures were approved by the Care and Use of Animals in Research Committee at The University of Akron.

A stock solution of the dye was prepared in DMSO at 1 mM. All stains were done at 10 μM with a 12 h incubation time and the fish were washed 3 times with E3 medium to limit potential background noise from aggregates. All fish were anesthetized with 0.5 mL of MS-2,2,2 20–30 minutes prior to imaging to limit movement. Once imaging was completed all the fish were euthanized using excess MS-2,2,2 before disposal.

Conclusion

In summary, we report the first example of flavonoid-based probes for ER sensing in living eukaryotic cells. The D-A type flavonoid **2** exhibits many attractive features, which include fluorescence turn-on, protein interaction, low toxicity, good biocompatibility, efficient cellular uptake, and ER-targeting. The peculiar ER-targeting has been independently observed in all the cell lines examined. The low molecular mass of **2** (<500 Dalton), in addition to its low toxicity, is anticipated to cause less perturbation to cellular function during imaging. It appears that the amide group plays an important role in the observed ER selectivity, as the control compounds without the amide side chain do not exhibit good selectivity to ER (ESI,† Fig. S3). The observed ER selectivity from **2** could be attributed to flavonoid's binding to specific proteins. Further study will be carried out to fine tune the ER selectivity.

Acknowledgements

This work was supported by NIH (Grant No. 1R15EB014546-01A1). We also thank the Coleman endowment from the University of Akron for partial support.

References

- 1 C. Xu, B. Bailly-Maitre and J. C. Reed, Endoplasmic reticulum stress: cell life and death decisions, *J. Clin. Invest.*, 2005, **115**, 2656–2664.
- 2 P. Walter and D. Ron, The unfolded protein response: from stress pathway to homeostatic regulation, *Science*, 2011, **334**, 1081–1086.
- 3 I. Kim, W. Xu and J. C. Reed, Cell death and endoplasmic reticulum stress: disease relevance and therapeutic opportunities, *Nat. Rev. Drug Discovery*, 2008, **7**, 1013–1030.
- 4 H. Yoshida, ER stress and diseases, *FEBS J.*, 2007, **274**, 630–658.
- 5 D. J. Todd, A. H. Lee and L. H. Glimcher, The endoplasmic reticulum stress response in immunity and autoimmunity, *Nat. Rev. Immunol.*, 2008, **8**, 663–674.
- 6 S. Fu, S. M. Watkins and G. Hotamisligil, The Role of Endoplasmic Reticulum in Hepatic Lipid Homeostasis and Stress Signaling, *Cell Metab.*, 2012, **15**, 623–634.
- 7 L. Ozcan and I. Tabas, Role of Endoplasmic Reticulum Stress in Metabolic Disease and Other Disorders, *Annu. Rev. Med.*, 2012, **63**, 317–328.
- 8 P. van Bergeijk, C. C. Hoogenraad and L. C. Kapitein, Right Time, Right Place: Probing the Functions of Organelle Positioning, *Trends Cell Biol.*, 2016, **26**, 121–134.
- 9 N. Baumann and D. Pham-Dinh, Biology of Oligodendrocyte and Myelin in the Mammalian Central Nervous System, *Physiol. Rev.*, 2001, **81**, 871–927.
- 10 S. E. Pfeiffer, A. E. Warrington and R. Bansal, The oligodendrocyte and its many cellular processes, *Trends Cell Biol.*, 1993, **3**, 191–197.
- 11 J. Praet, C. Guglielmetti, Z. Berneman, A. Van der Linden and P. Ponsaerts, Cellular and molecular neuropathology of the cuprizone mouse model: clinical relevance for multiple sclerosis, *Neurosci. Biobehav. Rev.*, 2014, **47**, 485–505.
- 12 A.-S. Dhaunchak and K.-A. Nave, A common mechanism of PLP/DM20 misfolding causes cysteine-mediated endoplasmic reticulum retention in oligodendrocytes and Pelizaeus-Merzbacher disease, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 17813–17818.
- 13 J. Wu, W. Liu, J. Ge, H. Zhang and P. Wang, New sensing mechanisms for design of fluorescent chemosensors emerging in recent years, *Chem. Soc. Rev.*, 2011, **40**, 3483–3495.
- 14 J. M. Meinig, L. Fu and B. R. Peterson, Synthesis of Fluorophores that Target Small Molecules to the Endoplasmic Reticulum of Living Mammalian Cells, *Angew. Chem., Int. Ed.*, 2015, **54**, 9696–9699.
- 15 F. M. Ashcroft and F. M. Gribble, Correlating structure and function in ATP-sensitive K⁺ channels, *Trends Neurosci.*, 1998, **21**, 288–294.
- 16 A. Hambrock, C. Löffler-Walz and U. Quast, Glibenclamide binding to sulphonylurea receptor subtypes: dependence on adenine nucleotides, *Br. J. Pharmacol.*, 2002, **136**, 995–1004.
- 17 A. J. Smith, T. K. Taneja, J. Mankouri and A. Sivaprasadarao, Molecular cell biology of K_{ATP} channels: implications for neonatal diabetes, *Expert Rev. Mol. Med.*, 2007, **9**, 1–17.

- 18 R. J. Nijveldt, E. van Nood, D. E. van Hoom, P. G. Boelens, K. van Norren and P. A. van Leeuwen, Flavonoids: a review of probable mechanisms of action and potential applications, *Am. J. Clin. Nutr.*, 2001, **74**, 418–425.
- 19 A. Garcia-Lafuente, E. Guillamon, A. Villares, M. A. Rostagno and J. A. Martinez, Flavonoids as anti-inflammatory agents: implications in cancer and cardiovascular disease, *Inflammation Res.*, 2009, **58**, 537–552.
- 20 A. S. Klymchenko, G. Duportail, Y. Mely and A. P. Demchenko, Ultrasensitive two-color fluorescence probes for dipole potential in phospholipid membranes, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 11219–11224.
- 21 B. Liu, M. Shah, Z. Ge, Q. Liu and Y. Pang, Biocompatible Flavone-Based Fluorogenic Probes for Quick Wash-Free Mitochondrial Imaging in Living Cells, *ACS Appl. Mater. Interfaces*, 2014, **6**, 21638–21644.
- 22 B. Liu, Y. Pang, R. Bouhenni, E. Duah, S. Paruchuri and L. McDonald, A step toward simplified detection of serum albumin on SDS-PAGE using an environment-sensitive flavone sensor, *Chem. Commun.*, 2015, **51**, 11060–11063.
- 23 T. R. Gormley and W. I. O'Sullivan, Flavanoid epoxides XIII: acid and base catalysed reactions of 2'-tosyloxychalcone epoxides. Mechanism of the algar-flynn-oyamada reaction, *Tetrahedron*, 1973, **29**, 369–373.
- 24 B. Liu, J. Wang, G. Zhang, R. Bai and Y. Pang, Flavone-Based ESIPT Ratiometric Chemodosimeter for Detection of Cysteine in Living Cells, *ACS Appl. Mater. Interfaces*, 2014, **6**, 4402–4407.
- 25 V. V. Shynkar, A. S. Klymchenko, E. Piemont, A. P. Demchenko and Y. Mely, Dynamics of Intermolecular Hydrogen Bonds in the Excited States of 4'-Dialkylamino-3-hydroxyflavones. On the Pathway to an Ideal Fluorescent Hydrogen Bonding Sensor, *J. Phys. Chem. A*, 2004, **108**, 8151–8159.
- 26 D. Magde, R. Wong and P. G. Seybold, Fluorescence Quantum Yields and Their Relation to Lifetimes of Rhodamine 6G and Fluorescein in Nine Solvents: Improved Absolute Standards for Quantum Yields, *Photochem. Photobiol.*, 2002, **75**, 327–334.
- 27 K. Iwata, C. C. Café-Mendes, A. Schmitt, J. Steiner, T. Manabe, H. Matsuzaki, P. Falkai, C. W. Turck and D. Martins-de-Souza, The human oligodendrocyte proteome, *Proteomics*, 2013, **13**, 3548–3553.
- 28 K. Suzuki, A. Gerelchuluun, Z. Hong, L. Sun, J. Zenkoh, T. Moritake and K. Tsuboi, Celecoxib enhances radiosensitivity of hypoxic glioblastoma cells through endoplasmic reticulum stress, *Neuro-Oncology*, 2013, **15**, 1186–1199.
- 29 K. Nagamoto-Combs, J. Kulas and C. K. Combs, A novel cell line from spontaneously immortalized murine microglia, *J. Neurosci. Methods*, 2014, **233**, 187–198.
- 30 L. J. Lawson, V. H. Perry and S. Gordon, Turnover of resident microglia in the normal adult mouse brain, *Neuroscience*, 1992, **48**, 405–415.
- 31 B. M. Costa, H. Yao, L. Yang and S. Buch, Role of Endoplasmic Reticulum (ER) Stress in Cocaine-Induced Microglial Cell Death, *J. Neuroimmune Pharmacol.*, 2013, **8**, 705–714.