# ACS Medicinal Chemistry Letters

Letter

Subscriber access provided by BIU Pharmacie | Faculté de Pharmacie, Université Paris V

## Missing Selectivity of Targeted 4#-Phorbol Prodrugs Expected to be Potential Chemotherapeutics

Ilari Tarvainen, Tomas Zimmermann, Pia Heinonen, Maria Helena Jäntti, Jari Yli-Kauhaluoma, Virpi Talman, Henrik Franzyk, Raimo K Tuominen, and Søren Brøgger Christensen

> ACS Med. Chem. Lett., Just Accepted Manuscript • DOI: 10.1021/ acsmedchemlett.9b00554 • Publication Date (Web): 23 Dec 2019

Downloaded from pubs.acs.org on December 25, 2019

### Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

#### 

## Missing Selectivity of Targeted 4β-Phorbol Prodrugs Expected to be Potential Chemotherapeutics

Ilari Tarvainen,<sup>†</sup> Tomáš Zimmermann,<sup>§,‡</sup> Pia Heinonen,<sup>†</sup> Maria Helena Jäntti,<sup>†</sup> Jari Yli-Kauhaluoma,<sup>∥</sup> Virpi Talman,<sup>†,⊥</sup> Henrik Franzyk,<sup>§</sup> Raimo K Tuominen,<sup>†</sup> Søren Brøgger Christensen<sup>§\*</sup>

<sup>†</sup> Drug Research Program, Division of Pharmacology and Pharmacotherapy, Faculty of Pharmacy, University of Helsinki, Helsinki, Finland

<sup>§</sup> Department of Drug Design and Pharmacology, Faculty of Health and Medical Sciences, University of Copenhagen, Jagtvej 162, DK-2100, Copenhagen, Denmark

<sup>‡</sup> Department of Chemistry of Natural Compounds, Faculty of Food and Biochemical Technology, University of Chemistry and Technology, Prague, Czech Republic

<sup>I</sup> Drug Research Program, Division of Pharmaceutical Chemistry and Technology, Faculty of Pharmacy, University of Helsinki, Helsinki, Finland

<sup>⊥</sup> National Heart and Lung Institute, Imperial College London, London, United Kingdom



#### Abstract

Targeting cytotoxic  $4\beta$ -phorbol esters toward cancer tissue was attempted by conjugating a  $4\beta$ -pborbol derivative with substrates for the proteases prostate-specific antigen (PSA) and prostate-specific membrane antigen (PSMA) expressed in cancer tissue. The hydrophilic peptide moiety was hypothesized to prevent penetration of the prodrugs into cells and prevent interaction with PKC. Cleavage of the peptide in cancer tumors was envisioned to release lipophilic cytotoxins, which subsequently penetrate into cancer cells. The  $4\beta$ -phorbol esters were prepared from  $4\beta$ -phorbol isolated from *Croton tiglium* seeds, while the peptides were prepared by solid-phase synthesis. Cellular assays revealed activation of PKC by the prodrugs and efficient killing of both peptidase positive as well as peptidase negative cells. Consequently no selectivity for enzyme expressing cells was found.

**Keywords:** 4β-Phorbol ester, Protease-assisted targeting, Targeted chemotherapy, Prodrug, Prostate-specific antigen. Prostate-specific membrane antigen,

**Memoriam**: This article is written in memory of the valuable contributions professor Maurizio Botta has offered to medicinal chemistry

#### **ACS Medicinal Chemistry Letters**

Prostate cancer (PCa) is a major cause of death by cancer in men in high-income countries.<sup>1</sup> In the initial stage, PCa mainly consists of cells that are androgen-dependent, and the growth can be retarded by hormone therapy.<sup>2</sup> Unfortunately, in later stages hormone refractory cells dominate (castration-resistant prostate cancer, CRPC).<sup>2, 3</sup> At this stage the use of common chemotherapeutics is complicated by the slow proliferation of the cancer tissue, since chemotherapeutics like taxanes, doxorubicine or vincristine target the proliferative stages of cancer. Thus, selectivity is obtained by the faster division rate for cancer cells.<sup>3-6</sup> Therefore an urgent need for drugs against late-stage PCa exists.

Pre-clinical evidence supports the idea that drugs targeting protein kinase C (PKC) may be useful in treatment of CRPC.<sup>7</sup> The PKC family comprises ten serine/threonine kinases, which can be divided into three groups: i) conventional PKC (cPKCs: - $\alpha$ , - $\beta$ I, - $\beta$ II and - $\gamma$ , ii) novel PKCs (nPKCs: - $\delta$ , - $\epsilon$ , - $\theta$  and - $\eta$ ) and iii) atypical PKCs (aPKCs: - $\zeta$ , -I and  $\lambda$ ) Expression and function of different PKC isoforms are context- and cell type-specific<sup>8-11</sup>. High expression of PKC $\delta$  has been reported in prostate cancer, and activation of PKC $\delta$  induces apoptosis in LNCaP PCa cells.<sup>9, 12-15</sup> PKC $\epsilon$  is generally overexpressed in PCa and downregulation of PKC $\epsilon$  induces apoptosis.<sup>16-20</sup>

PKC-activating diterpenoids related to PMA (1, Fig. 1) have been in clinical trials.<sup>21-23</sup> Tiglianol tiglate (ECB-46) awaits approval by FDA and EMA for treatment of mast cell tumours in dogs<sup>24</sup> and is entering phase IIA clinical trials for treatment of head and neck squamous cell carcinoma (HNSCC) in humans. Ingenol 3-angelate has under the trade name Picato<sup>R</sup> been approved by FDA in 2012 as a topical gel for the treatment of actinic keratosis (preliminary stage of skin cancer).<sup>25, 26</sup> Since PKC is present in virtually all cells, administration of phorbol esters may affect normal physiology in a broad sense. Selectivity of cancer therapies may be obtained by taking advantage of proteases present in tumors.<sup>27</sup> Prostate specific antigen (PSA), a peptidase expressed by the prostate and PCa is a diagnostic marker for prostate cancer and it has been suggested to be involved in cancer invasion and metastasis.<sup>28-32</sup> The missing activity of PSA in the blood caused by complexation with proteins like blood albumin,<sup>3, 27</sup> makes the enzyme a potential facilitator for selective drug delivery. Since both PSA and prostate specific membrane antigen (PSMA) are expressed by PCa even when they become more undifferentiated and anaplastic, they appear to be promising tools in the targeting of toxins for tumors even in CRPC <sup>33, 34</sup>. In the case of thapsigargin (Tg1, Supporting Information Fig. S27) selectivity toward cancer tissue was obtained by conjugation of 8-O-12-aminododecanoyl-8-O-debutanoylthapsigargin with peptides that are substrates for human glandular kallikrein 2, hK2,<sup>35, 36</sup> PSA or PSMA (mipsagargin, Tg6, Fig. S27).<sup>37</sup> Mipsagargin has successfully passed clinical trial 2 (For details see Supporting Information paragraph S3).<sup>38</sup> Based on the above findings 4 $\beta$ -phorbol esters **4-6** (Fig. 1) were designed and expected to display a similar behaviour in the organism as the thapsigargin analogs. By

conjugating the toxin with a substrate for the proteases, penetration into cells is ideally only possible after enzymatic cleavage by either PSA or PSMA.<sup>30, 37</sup> Encouraged by the above-mentioned findings and hypotheses we have attempted to develop prodrugs of 4 $\beta$ -phorbol esters for selective targeting of PSA- and PSMA-expressing cancer cells.

ACS Paragon Plus Environment



**Fig. 1.** Target compounds and starting material:  $4\beta$ -Phorbol 12-O-myristate 13-O-acetate (1), toxin 2 obtained after cleavage of prodrugs 4 and 5 with hK2 or PSA, respectively, while toxin 3 is obtained after cleavage of prodrug 6 with PSMA.  $4\beta$ -Phorbol (7). Compound 8 is the starting material for synthesis of compounds 2 – 3 and compound 9 for 4 – 6.

The starting material 4 $\beta$ -phorbol (7) was obtained from seeds of *Croton tiglium* L. (Euphorbiaceae) (for details see Supporting Information paragraph S2.5.1). By a few synthetic steps 4 $\beta$ -pborbol was converted into the cytotoxins **2** and **3** via **8** (Supportin Information S2.5.2). The peptides needed for preparing the prodrugs 4 – 6 were prepared by solid phase syntheses For syntheses and characterization of the 4 $\beta$ -phorbol toxins and prodrugs see Supporting Information S2.5.3 – S2.5.10.

Binding to PKC as Measured by [<sup>3</sup>H]PDBu Displacement assay. Compounds 2 and 3 as well as prodrugs 4 - 6) were tested for binding to the C1 domains of PKC $\alpha$  in a 96-well plate filtration assay as described earlier<sup>39</sup> at a concentration range of 0.01-10 µM. All new compounds (i.e., 2 -6) displaced [<sup>3</sup>H]4 $\beta$ -phorbol 12,13-dibutyrate ([<sup>3</sup>H]PDBu) as efficiently as PMA (1) (Fig. 2) except for prodrug 5, for which an approximately ten times higher concentration was required to achieve a displacement comparable to that of the other compounds. Thus, the presence of a peptide moiety in the prodrugs did not nullify their affinities to the C1 domain of PKC $\alpha$ .



**Fig. 2**. Displacement binding curves of prodrugs 4 - 6. Toxins 2 and 3 and PMA. Binding of [<sup>3</sup>H]PDBu (10 nM) to PKC $\alpha$  was measured in the presence of increasing concentrations of the tested compounds. The PKC $\alpha$  was obtained from a lysate if cells overexpressing the enzyme. The data is presented as mean of residual [3H]PDBu binding (% of control) from three parallel samples in a single representative experiment.

*Cell Death as Measured by Cell viability assays.* The effect of the compounds on viability of PCa cells was determined by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay (Fig.4). The previously established PSMA prodrug mipsagargin (**Tg6**) and the PSMA cleavage product Asp-12-AD-thapsigargin (**Tg3**) were used as reference compounds <sup>37</sup>. The PCa cell lines, used in the present study, represent different types of PCa: androgen-unresponsive DU145 and PC3 cells, which do not express PSA or PSMA, and androgen-responsive 22Rv1 and LNCaP cells, which both are PSA- and PSMA-positive.<sup>40, 41</sup> The maximal effect of the PSA cleavage product **2** with the highest concentration gave rise to a reduction in viability of PSA/PSMA-positive LNCaP and 22Rv1 cells to ~60% and ~10%, respectively, and in PSA/PSMA-negative DU145 and PC3 cells to ~40% and ~30%, respectively (Fig. 3). PSA prodrug **5** reduced the viability to below 50% only in LNCaP cells (~40%) at 20  $\mu$ M and 40  $\mu$ M concentrations, whereas PSA prodrug **4** reduced the viability to below 50% at the highest concentration not only in PSA-positive 22Rv1 (to ~25%) but also in PSA-negative PC3 cells (~15%) at the highest concentration. To our surprise, the PSMA cleavage product **3** had almost no effect on cell viability in any of the PCa cell lines. The PSMA prodrug **6**, however, decreased the viability to ~10% at the

highest concentration only in the PSMA-negative PC3 cells (Fig. 3). The reference compound **Tg3** decreased the viability concentration-dependently in all PC cell lines, and its maximal effect (achieved with the highest concentration) was a reduction in viability to ~4% for LNCaP, 27% for 22Rv1, 20% for DU145, and ~15% for PC3 cells. Surprisingly, the other reference compound (i.e., **Tg6**) demonstrated a similar reduction in the viability in PSMA-negative PC3 cells (to ~32%) as seen for the PSMA-positive LNCaP and 22Rv1 cells (to ~35% and ~40%, respectively; Fig. 3). PMA is known to promote PKC-induced apoptosis in the LNCaP cell line <sup>42</sup>. In accordance with this, the pan-PKC inhibitor Gö6983 was able to dampen the effect of 20 µM of compound **5** in

LNCaP cells (Fig. 3), indicating that the cytotoxic effect indeed is PKC-mediated. The compounds did not induce distinct damage to the cell membranes during the 72-h incubation with any of the concentrations as determined by the LDH test (Fig. S2).



**Fig. 3.** Effects of phorbol prodrugs **4** - **6**, and cleavage products **2** and **3** as well as **Tg3** and **Tg6** on viability of PCa cell lines. (A) PC3; (B) LNCaP; (C) DU145; (D) 22Rv1 and the effect of PKC inhibitor Gö6983 (1  $\mu$ M) on the effect of 20  $\mu$ M of **2** - **6**, **Tg3**, **Tg6** and 100 nM PMA on viability of LNCaP, 22Rv1, DU145 and PC3 cells (E). Cell viability was measured

after 72 h incubation with the compounds by utilizing the MTT assay. The data is presented as mean of cell viability (% of control) (n=3).

#### Effects on ERK1/2 phosphorylation and protein expression of PKC and PSMA

Extracellular signal-regulated kinases 1 and 2 (ERK1/2) are members of the mitogen-activated protein kinase (MAPK) signalling cascade that controls several cellular processes <sup>43</sup>. PKC-mediated ERK1/2 phosphorylation is one of the initial rapid events in PMA-treated LNCaP cells <sup>42</sup> Since the novel 4β-phorbol-derived compounds compete with PDBu *in vitro*, their ability to modulate ERK activity was investigated in living cells.<sup>44</sup> PSA/PSMA-positive 22Rv1 and PSA/PSMA-negative DU145 PCa cell lines were exposed to 20 µM of compounds **2** - **6**, **Tg3** and **Tg6** and to 10 nM of PMA for 30 min. Phorbol-derived compounds **2** - **5** induced substantial ERK1/2 phosphorylation in 22Rv1 cells (Fig. 4). The ERK1/2 phosphorylation was even more distinct than after PMA exposure, except in cells treated with compound **6**. The phorbol derivatives also induced ERK1/2 phosphorylation in DU145 cells, but the magnitude of the effect was distinctively smaller than in 22Rv1 cells (Fig. 4).

Downregulation of PKC has been suggested to explain the tumor promotion caused by phorbol esters.<sup>10, 45</sup> In addition PMA is able to induce androgen receptor downregulation in PCa cells, which is associated with PSMA downregulation. <sup>46, 47</sup> To understand the effects of our compounds on the expression of PKC and PSMA we decided to investigate the effects of the phorbol derivatives on the expression levels of PKC $\alpha$  and PKC $\delta$  and PSMA in 22Rv1 cells. A 24 h exposure to 20  $\mu$ M **3** and **6** increased the expression of PKC $\alpha$  while the incubation with **2** and **4** had no effect. However, PKC $\alpha$ , PKC $\delta$  and PSMA expression was reduced upon 24 h exposure to all compounds, including **3** and **6** (Fig. S4 and Table S3). Indeed, our results support the hypothesis that phorbol ester induced down regulation of PKC is associated with down regulation of PSMA. Probably the PKC activating effects of our toxins caused a down regulation of PKC and PSMA.



**Fig. 4**. Effects of phorbol derivatives **2** - **6** on ERK1/2 phosphorylation in PC cells. Quantifications from DU145 and 22Rv1 cells. Data is presented as mean + SEM (N = 3; \*P < 0.05 vs ctrl, Welch's t-test). The cells were treated with 20  $\mu$ M of different phorbol derivatives and PMA for 30 min. The cells were harvested, and then ERK1/2 phosphorylation was analysed by using Western blotting with detection as described in the Experimental Section.

#### Conclusion

In the present study, we synthesised, characterised and evaluated ability of the PSA/PSMAactivable 4β-phorbol ester prodrugs 4 - 6 and the corresponding cytotoxins 2 and 3 to displace <sup>3</sup>H PDBu from PKC and to decrease the viability of PSMA/PSA-positive as well as PSMA/PSAnegative PCa cell lines. In addition, we performed studies on their abilities to increase ERK1/2 phosphorylation. The synthesised PSA/PSMA-activable 4β-phorbol ester prodrugs were designed to contain peptide sequences that are specifically cleaved by either hK2 (i.e., 4), PSA (i.e., 5), or PSMA (i.e., 6). All phorbol-derived compounds showed low nanomolar binding affinity to the C1 domain of recombinant human PKCa, as shown by their displacement of PBDu (Fig. 3). Compound 3 induced cytotoxicity only to a limited extent, which may be explained by the lack of ability to penetrate the cell membrane due to its zwitterionic nature at physiological pH. Analogously, compound 3 and prodrug 6 provoked phosphorylation of ERK1/2 only to a limited extent (Fig. 3 and 4). Disappointingly prodrugs 4 - 6 showed no selectivity for PSA/PSMA-positive cell lines (i.e., LNCaP/22Rv1) over PSA/PSMA-negative cell lines (i.e., DU145 and PC3). This observation strongly infers that despite conjugation to a hydrophilic peptide the prodrugs (compounds 4 - 6) retain an ability to penetrate cell membranes. The poor activity of 3 and 6 in the viability assay and in the phosphorylation assay indicates that these highly charged molecules can be taken up by the cells only to a limited extent. The results of the present study do not support the hypothesis that the designed PSA/PSMA-targeted prodrugs are capable of providing selective toxicity to PSA/PSMA-expressing PCa cells. Importantly the present results obtained for the known

prodrug mipsagargin (**Tg6**, Supplementary Information Fig. S26) does not support previous observations of selectivity for peptidase expressing cell lines<sup>30</sup> since **Tg6** exhibited clear toxicity both on the PSMA-negative cell line PC3 and on the PSMA-expressing cell lines LNCaP and 22Rv1.<sup>37</sup> A similar poor selectivity has recently been reported by Akinboy et al.<sup>3</sup> fo PSA-targeted *O*-8-(morpholine-4-carbonyl-His-Ser-Ser-Lys-Leu-Phe-GIn-Leu-*N*-12-aminododecanoyl)-*O*-8-debutanoyl-thapsigargin (**Tg5**). The missing selectivity of the 4β-phorbol alanlogs are even more

surprising than the missing selectivity of the thapsigargin analogs since the first mentioned are calculated to have lower logP values. In conclusion, the proposed targeted therapy involving conjugation to peptides that are selectively cleaved by proteases present in cancer tissue appears to lack the desired selectivity with  $4\beta$ -phorbol and thapsigargin analogs.

### ASSSOCIATED CONTENT

Supporting information

The Supporting Information is available free of charge on the ACS Publications website at DOI Experimental procedures, spectral data for all compounds, spectra for al target compounds and supporting biological data are available.

### AUTHOR INFORMATION

CORRESPONDING AUTHOR

E-mail: soren.christensen@sund.ku.dk

## ORCID

Ilari Tarvainen 0000-0003-3386-465X

Tomáš Zimmermann 0000-0003-0529-723X

Pia Heinonen 0000-0002-0988-8013

Maria Helena Jäntti 0000-0003-1468-3861

Jari Yli-Kauhaluoma 0000-0003-0370-7653

Virpi Talman 0000-0002-2702-6505

Henrik Franzyk 0000-0002-2822-1927

Raimo Kalevi Tuominen 0000-0001-7603-0749

Søren Brøgger Christensen 0000-0002-7067-5586

### **Author Contributions**

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript. Ilari Tarvainen and Tomáš Zimmermann has contributed equally to the manuscript.

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGEMENTS

This project was funded by the Jane and Aatos Erkko Foundation, Doctoral Programme in Drug Research (University of Helsinki), University of Copenhagen and University of Chemistry and Technology in Prague. We thank Ms. Johanna Mosorin for technical assistance.

## ABBREVIATIONS

Boc: tert-butyloxycarbonyl; CRPC: castration-resistant prostate cancer, 2-CTC: 2-chlorotrityl chloride; 2-CI-Trt: 2-chlorotrityl; DCM: dichloromethane; DIPEA: *N,N*-diisopropylethylamine; DMAP: 4-(*N*,*N*-dimethylamino)pyridine; DMF: *N*,*N*-dimethylformamide; EDC: 1-ethyl-3-(3dimethylaminopropyl)carbodiimide; Fmoc: fluorenylmethyloxycarbonyl; GPCR: G-protein-coupled receptor; HBTU: 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate; HFIP: 1,1,1,3,3,3-hexafluoro-2-propanol; InsP<sub>3</sub>: inositol-1,4,5-trisphosphate; MTT: methylthiazolyldiphenyltetrazolium bromide; MW: microwave; NMP: N-methyl-2-pyrrolidone; PCa: prostate cancer; PDBu: phorbol-12,13-dibutyrate; PKC: protein kinase C; PMA: 48-phorbol 12myristate 13-acetate; PSA: prostate-specific antigen; PSMA: prostate-specific membrane antigen; PyBOP: benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate; SPPS: solid-phase peptide synthesis; TBDMS: tert-butyldimethylsilyl; THF: tetrahydrofuran

### References

- (1) Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R. L.; Torre, L. A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* **2018**, 68, 394-424.
- (2) Hwang, C. Overcoming docetaxel resistance in prostate cancer: a perspective review. *Ther. Adv. Med. Oncol.* **2012**, 4, 329-340, 12.
- (3) Akinboye, E. S.; Brennen, W. N.; Denmeade, S. R.; Isaacs, J. T. Albumin-linked prostatespecific antigen-activated thapsigargin- and niclosamide-based molecular grenades targeting the microenvironment in metastatic castration-resistant prostate cancer. *Asian J Urol* **2019**, 6, 99-108.
- (4) Chan, K. S.; Koh, C. G.; Li, H. Y. Mitosis-targeted anti-cancer therapies: where they stand. *Cell Death Dis.* **2012**, 3, e411, 11 pp.
- (5) Berges, R. R.; Vukanovic, J.; Epstein, J. I.; CarMichel, M.; Cisek, L.; Johnson, D. E.; Veltri, R. W.; Walsh, P. C.; Isaacs, J. T. Implication of cell kinetic changes during the progression of human prostatic cancer. *Clin.Cancer Res.* **1995**, 1, 473-480.
- (6) Pinski, J.; Parikh, A.; Bova, G. S.; Isaacs, J. T. Therapeutic implications of enhanced G0/G1 checkpoint control induced by coculture of prostate cancer cells with osteoblasts. *Cancer Res.* **2001**, 61, 6372-6376.
- (7) Inoue, T.; Ogawa, O. Role of signaling transduction pathways in development of castrationresistant prostate cancer. *Prostate Cancer* **2011**, 647987, 7 pp.

1	
2	
3 1	
4 5	(8) Cornford, P.; Evans, J.; Dodson, A.; Parsons, K.; Woolfenden, A.; Neoptolemos, J.; Foster, C. S.
6	<b>1000</b> 154 137 144
7	<b>1999</b> , 154, 137-144.
8	(9) Villar, J.; Arenas, M. I.; MacCartny, C. M.; Bianquez, M. J.; Tirado, O. M.; Notario, V.
9	PCPH/ENTPD5 expression enhances the invasiveness of human prostate cancer cells by a Protein Kinase Co-
10	Dependent Mechanism. <i>Cancer Res.</i> <b>2007,</b> 67, 10859-10868.
11	(10) Antal, C. E.; Hudson, A. M.; Kang, E.; Zanca, C.; Wirth, C.; Stephenson, N. L.; Irotter, E. W.;
12	Gallegos, L. L.; Miller, C. J.; Furnari, F. B.; Hunter, T.; Brognard, J.; Newton, A. C. Cancer-Associated Protein
14	Kinase C Mutations Reveal Kinase's Role as Tumor Suppressor. Cell (Cambridge, MA, United States) 2015,
15	160, 489-502.
16	(11) Pandian, S. S.; Sneddon, A. A.; Bestwick, C. S.; McClinton, S.; Grant, I.; Wahle, K. W. J.; Heys,
17	S. D. Fatty Acid regulation of protein kinase C isoforms in prostate cancer cells. <i>Biochem. Biophys. Res.</i>
18	<i>Commun.</i> <b>2001,</b> 283, 806-812.
19	(12) Rusnak, J. M.; Lazo, J. S. Downregulation of protein kinase C suppresses induction of
20	apoptosis in human prostatic carcinoma cells. <i>Exp. Cell Res.</i> <b>1996,</b> 224, 189-99.
21	(13) Lamm, M. L. G.; Long, D. D.; Goodwin, S. M.; Lee, C. Transforming growth factor-β1 inhibits
22	membrane association of protein kinase C $\alpha$ in a human prostate cancer cell line, PC3. Endocrinology <b>1997</b> ,
25 24	138, 4657-4664.
25	(14) Fujii, T.; Garcia-Bermejo, M. L.; Bernabo, J. L.; Caamano, J.; Ohba, M.; Kuroki, T.; Li, L.; Yuspa,
26	S. H.; Kazanietz, M. G. Involvement of protein kinase C delta (PKCdelta) in phorbol ester-induced apoptosis
27	in LNCaP prostate cancer cells. Lack of proteolytic cleavage of PKCdelta. J. Biol. Chem. 2000, 275, 7574-82.
28	(15) Kharait, S.; Dhir, R.; Lauffenburger, D.; Wells, A. Protein kinase Cdelta signaling downstream
29	of the EGF receptor mediates migration and invasiveness of prostate cancer cells. Biochem. Biophys. Res.
30	<i>Commun.</i> <b>2006,</b> 343, 848-56.
31	(16) Aziz, M. H.; Manoharan, H. T.; Church, D. R.; Dreckschmidt, N. E.; Zhong, W.; Oberley, T. D.;
32	Wilding, G.; Verma, A. K. Protein Kinase Cɛ interacts with signal transducers and activators of transcription
33 24	3 (Stat3), phosphorylates Stat3 Ser727, and regulates its constitutive activation in Prostate cancer. Cancer
35	Res. <b>2007,</b> 67, 8828-8838.
36	(17) Gundimeda, U.; Schiffman, J. E.; Chhabra, D.; Wong, J.; Wu, A.; Gopalakrishna, R. Locally
37	Generated Methylseleninic Acid Induces Specific Inactivation of Protein Kinase C Isoenzymes: relevance to
38	selenium-induced apoptosis in prostate cancer cells. J. Biol. Chem. 2008, 283, 34519-34531.
39	(18) Sarveswaran, S.; Gautam, S. C.; Ghosh, J. Wedelolactone, a medicinal plant-derived
40	coumestan, induces caspase-dependent apoptosis in prostate cancer cells via downregulation of PKCe
41	without inhibiting Akt. Int. J. Oncol. 2012, 41, 2191-2199.
42	(19) Sarveswaran, S.; Thamilselvan, V.; Brodie, C.; Ghosh, J. Inhibition of 5-lipoxygenase triggers
43	apoptosis in prostate cancer cells via down-regulation of protein kinase C-epsilon. <i>Biochim. Biophys. Acta,</i>
44 15	Mol. Cell Res. 2011, 1813, 2108-2117.
46	(20) BinHafeez, B.; Zhong, W.; Fischer, J. W.; Mustafa, A.; Shi, X.; Meske, L.; Hong, H.; Cai, W.;
47	Havighurst, T.; Kim, K. M.; Verma, A. K. Plumbagin, a medicinal plant (Plumbago zeylanica)-derived 1,4-
48	naphthoquinone, inhibits growth and metastasis of human prostate cancer PC-3M-luciferase cells in an
49	orthotopic xenograft mouse model. <i>Mol. Oncol.</i> <b>2013,</b> 7, 428-439.
50	(21) Gobbi, G.; Mirandola, P.; Carubbi, C.; Micheloni, C.; Malinverno, C.; Lunghi, P.; Bonati, A.;
51	Vitale, M. Phorbol ester-induced PKCe down-modulation sensitizes AML cells to TRAIL-induced apoptosis
52	and cell differentiation. <i>Blood</i> <b>2009,</b> 113, 3080-3087.
53 54	(22) Schaar, D.; Goodell, L.; Aisner, J.; Cui, X. X.; Han, Z. T.; Chang, R.; Martin, J.; Grospe, S.: Dudek.
54 55	L.; Riley, J.; Manago, J.; Lin, Y.; Rubin, E. H.; Conney, A.: Strair, R. K. A phase I clinical trial of 12- O-
56	tetradecanoylphorbol-13-acetate for patients with relapsed/refractory malignancies. <i>Cancer Chemother</i> .
57	Pharmacol. <b>2006,</b> 57, 789-795.
58	(23) Han, Z. T.; Tong, Y. K.; He. L. M.: Zhang, Y.: Sun. J. Z.: Wang, T. Y.: Zhang, H.: Cui, Y. L.:
59	Newmark, H. L.; Conney, A. H.; Chang, R. L. 12-O-tetradecanovlphorbol-13-acetate (TPA)-induced increase
60	

1 2 3 4 in depressed white blood cell counts in patients treated with cytotoxic cancer chemotherapeutic drugs. 5 Proc. Natl. Acad. Sci. U. S. A. 1998, 95, 5362-5365. 6 (24) Miller, J.; Campbell, J.; Blum, A.; Reddell, P.; Gordon, V.; Schmidt, P.; Lowden, S. Dose 7 Characterization of the Investigational Anticancer Drug Tigilanol Tiglate (EBC-46) in the Local Treatment of 8 9 Canine Mast Cell Tumors. Front Vet Sci 2019, 6, 106. 10 Tzogani, K.; Pignatti, F.; Nagercoil, N.; Hemmings Robert, J.; Samir, B.; Gardette, J.; Demolis, (25) 11 P.; Salmonson, T. The European Medicines Agency approval of ingenol mebutate (Picato) for the cutaneous 12 treatment of non-hyperkeratotic, non-hypertrophic actinic keratosis in adults: Summary of the scientific 13 assessment of the Committee for Medicinal Products for Human Use (CHMP). Eur. J. Dermatol. 2014, 24, 14 457-463. 15 Ersvaer, E.; Kittang, A. O.; Hampson, P.; Sand, K.; Gjertsen, B. T.; Lord, J. M.; Bruserud, O. The (26) 16 protein kinase C agonist PEP005 (ingenol 3-angelate) in the treatment of human cancer: a balance between 17 efficacy and toxicity. Toxins (Basel) 2010, 2, 174-194. 18 19 (27) Aloysius, H.; Hu, L. Targeted Prodrug Approaches for Hormone Refractory Prostate Cancer. 20 Med. Res. Rev. 2015, 35, 554-585. 21 Ishii, K.; Otsuka, T.; Iguchi, K.; Usui, S.; Yamamoto, H.; Sugimura, Y.; Yoshikawa, K.; Hayward, (28) 22 S. W.; Hirano, K. Evidence that the prostate-specific antigen (PSA)/Zn2+ axis may play a role in human 23 prostate cancer cell invasion. Cancer Letters (Amsterdam, Netherlands) 2004, 207, 79-87. 24 (29) Denmeade, S. R.; Jakobsen, C. M.; Janssen, S.; Khan, S. R.; Garrett, E. S.; Lilja, H.; Christensen, 25 S. B.; Isaacs, J. T. Prostate-Specific Antigen-Activated Thapsigargin Prodrug as Targeted Therapy for Prostate 26 Cancer. J. Natl. Cancer Inst. 2003, 95, 990-1000. 27 (30) Denmeade, S. R.; Isaacs, J. T. Engineering enzymatically activated "molecular grenades" for 28 29 cancer. Oncotarget. 2012, 3, 666-667. 30 (31) Denmeade, S. R.; Nagy, A.; Gao, J.; Lilja, H.; Schally, A. V.; Isaacs, J. T. Enzymatic activation of 31 a doxorubicin-peptide prodrug by prostate-specific antigen. Cancer Res. 1998, 58, 2537-2540. 32 Denmeade, S. R.; Isaacs, J. T.; Buckley, J. T. Proaerolysin containing protease activation (32) 33 sequences and methods use for treatment of prostate cancer. Eur. Pat. 2 518 142 B1, July 15, 2015. 34 (33) Rajasekaran, A. K.; Anilkumar, G.; Christiansen, J. J. Is prostate-specific membrane antigen a 35 multifunctional protein? Am. J. Physiol. 2005, 288, C975-C981. 36 Williams, S. A.; Singh, P.; Isaacs, J. T.; Denmeade, S. R. Does PSA play a role as a promoting (34) 37 agent during the initiation and/or progression of prostate cancer? Prostate 2007, 67, 312-329. 38 39 Lovgren, J.; Airas, K.; Lilja, H. Enzymatic action of human glandular kallikrein 2 (hK2). (35) 40 Substrate specificity and regulation by Zn2+ and extracellular protease inhibitors. European Journal of 41 Biochemistry 1999, 262, 781-789. 42 (36) Janssen, S.; Rosen, D. M.; Ricklis, R. M.; Dionne, C. A.; Lilja, H.; Christensen, S. B.; Isaacs, J. T.; 43 Denmeade, S. R. Pharmacokinetics, biodistribution, and antitumor efficacy of a human glandular kallikrein 2 44 (hK2)-activated thapsigargin prodrug. Prostate 2006, 66, 358-368. 45 Denmeade, S. R.; Mhaka, A. M.; Rosen, D. M.; Brennen, W. N.; Dalrymple, S.; Dach, I.; Olesen, (37) 46 C.; Gurel, B.; DeMarzo, A. M.; Wilding, G.; Carducci, M. A.; Dionne, C. A.; Moeller, J. V.; Nissen, P.; 47 48 Christensen, S. B.; Isaacs, J. T. Engineering a prostate-specific membrane antigen-activated tumor 49 endothelial cell prodrug for cancer therapy. Sci. Transl. Med. 2012, 4, 140ra86, 13 pp. 50 Mahalingam, D.; Mahalingam, D.; Arora, S. P.; Sarantopoulos, J.; Peguero, J.; Campos, L.; Cen, (38) 51 P.; Rowe, J.; Allgood, V.; Tubb, B. A Phase II, Multicenter, Single-Arm Study of Mipsagargin (G-202) as a 52 Second-Line Therapy Following Sorafenib for Adult Patients with Progressive Advanced Hepatocellular 53 Carcinoma. Cancers (Basel) 2019, 11. 54 Boije af Gennas, G.; Talman, V.; Aitio, O.; Ekokoski, E.; Finel, M.; Tuominen, R. K.; Yli-(39) 55 Kauhaluoma, J. Design, Synthesis, and Biological Activity of Isophthalic Acid Derivatives Targeted to the C1 56 Domain of Protein Kinase C. J. Med. Chem. 2009, 52, 3969-3981. 57 58 59

1
2
2
ر ۸
4
5
6
7
8
9
10
11
12
13
14
15
16
10
1/
18
19
20
21
22
23
24
25
26
27
28
20
29
50 21
31
32
33
34
35
36
37
38
39
40
41
42
ד∠ ⊿2
40 44
44
45
46
47
48
49
50
51
52
53
54
14

(40) Bennett, N. C.; Hooper, J. D.; Johnson, D. W.; Gobe, G. C. Expression profiles and functional associations of endogenous androgen receptor and caveolin-1 in prostate cancer cell lines. *Prostate* (*Hoboken, NJ, U. S.*) **2014,** 74, 478-487.

(41) Ghosh, A.; Wang, X.; Klein, E.; Heston, W. D. W. Novel role of prostate-specific membrane antigen in suppressing prostate cancer invasiveness. *Cancer Res.* **2005**, 65, 727-731.

(42) Tanaka, Y.; Gavrielides, M. V.; Mitsuuchi, Y.; Fujii, T.; Kazanietz, M. G. Protein Kinase C Promotes Apoptosis in LNCaP Prostate Cancer Cells through Activation of p38 MAPK and Inhibition of the Akt Survival Pathway. *J. Biol. Chem.* **2003**, 278, 33753-33762.

(43) Chang, L.; Karin, M. Mammalian MAP kinase signalling cascades. *Nature (London, United Kingdom)* **2001,** 410, 37-40.

(44) Schonwasser, D. C.; Marajs, R. M.; Marshall, C. J.; Parker, P. J. Activation of the mitogenactivated protein kinase/extracellular signal-regulated kinase pathway by conventional, novel, and atypical protein kinase C isotypes. *Mol. Cell. Biol.* **1998**, 18, 790-798.

(45) Fuerstenberger, G.; Berry, D. L.; Sorg, B.; Marks, F. Skin tumor promotion by phorbol esters is a two-stage process. *Proc. Natl. Acad. Sci. U. S. A.* **1981**, 78, 7722-6.

(46) Itsumi, M.; Shiota, M.; Yokomizo, A.; Takeuchi, A.; Kashiwagi, E.; Dejima, T.; Inokuchi, J.; Tatsugami, K.; Uchiumi, T.; Naito, S. PMA induces androgen receptor downregulation and cellular apoptosis in prostate cancer cells. *J. Mol. Endocrinol.* **2014,** 53, 31-41.

(47) Liu, T.; Wu, L. Y.; Fulton, M. D.; Johnson, J. M.; Berkman, C. E. Prolonged androgen deprivation leads to downregulation of androgen receptor and prostate-specific membrane antigen in prostate cancer cells. *Int. J. Oncol.* **2012**, 41, 2087-2092.

#### **Graphical Abstract**

Missing Selectivity of Targeted 4β-Phorbol Prodrugs Expected to be Potential Chemotherapeutics

Ilari Tarvainen, Tomáš Zimmermann, Pia Heinonen, Maria Helena Jäntti, Jari Yli-Kauhaluoma, Virpi Talman, Henrik Franzyk, Raimo K Tuominen, Søren Brøgger Christensen\*

