

Preparation of 5-Alkyl-3-carboxymethylrhodanines and Their Aldose Reductase Inhibitory Activity

Yoshitaka OHISHI,* Teruo MUKAI, Michiko NAGAHARA, Motoyuki YAJIMA, and Norio KAJIKAWA

Central Research Institute, Kaken Pharmaceutical Co., Ltd., 14, Minamikawara-machi, Shinomiya, Yamashina-ku, Kyoto 607, Japan.

Received September 4, 1991

5-Alkyl-3-carboxymethylrhodanines (2) were prepared from **5-alkylmethylidene-3-carboxymethylrhodanines (1)**. The *exo* double bond of **1** was successfully reduced with NaBH₄. The 1,4-addition reaction path was confirmed on the basis of proton nuclear magnetic resonance spectrum of the product (**4b**) obtained from the reduction of **3** using NaBD₄. Optical resolution of the *tert*-butyl compound (**2i**) was achieved upon epimerization–crystallization method using L-3-amino-ε-caprolactam. The alkyl compounds (**2**) and the optical active compounds ((+)-**2i**, (–)-**2i**) were evaluated for aldose reductase inhibitory potency.


Keywords 5-alkyl-3-carboxymethylrhodanine; 5-alkylmethylidene-3-carboxymethylrhodanine; NaBH₄ reduction; *exo* carbon–carbon double bond; imide group; 1,4-addition; optical resolution; epimerization–crystallization method; aldose reductase inhibitory potency

Since Dvornik's proposal in 1973,¹⁾ many kinds of compounds have been prepared and their aldose reductase (AR) inhibitory activity have been examined in the search to create drugs for treatment of diabetic complications. AR inhibitors are divided into structural types, many compounds with carboxyl group have been particularly well studied²⁾ and several representative compounds are now being tested in clinical stage.³⁾ Recently we reported synthesis and AR inhibitory activity of two kinds of novel AR inhibitors having carboxyl group: benzo[*b*]furans derivatives possessing a carboxymethylsulfamoyl group⁴⁾ and 5-alkylmethylidene-3-carboxymethylrhodanines (**1**).⁵⁾ In the course of the studies on 3-carboxymethylrhodanine derivatives, the AR inhibitory activity of 5-alkyl-3-carboxymethylrhodanines (**2**) was compared with that of 5-alkylmethylidene-3-carboxymethylrhodanines (**1**). In this paper, we describe reduction of **1** to **2** and aldose reductase inhibitory potency of **2**.

There is appreciable interaction between *exo* carbon–carbon double bond at 5-position and functions (C=O, S) of the rhodanine ring. We expect that reduction of the *exo* double bond of **1** may alter the inhibitory potency. Selective reduction of the *exo* double bond of **1a–o** was carried out with NaBH₄ in *N,N*-dimethylformamide (DMF) without reduction of carbonyl and thiocarbonyl group of the rhodanine moiety under mild conditions to give the corresponding saturated compounds (**2a–o**) in reasonable yields (Chart 1, Table I).

The use of NaBH₄ for the reduction of aldehydes and ketones is now well established,⁶⁾ and a few applications to reduction of carbon–carbon double bond conjugated with nitro,⁷⁾ nitrile,⁸⁾ carbonyl⁹⁾ and ester functions¹⁰⁾ have been reported. To the best of our knowledge, the present successful reduction of the carbon–carbon double bond conjugated with the imide group may be the first finding in NaBH₄ reduction.

The reduction mechanism of the double bond in 5-*tert*-butylmethylidene-3-methoxycarbonylmethylrhodanine (**3**), more suitable than the carboxyl compound (**1i**) for identification of the mechanism, was investigated using NaBD₄. The methylidene compound (**3**) was treated with 0.28 eq of NaBH₄ in DMF to afford the 5-(2,2-dimethylpropyl) compound (**4a**) in 65% yield. In the proton nuclear magnetic resonance (¹H-NMR) of **4a** in CDCl₃, methylene proton signals of 2,2-dimethylpropyl group were detected at 1.81 (1H, double doublet, *J*=14.5, 11.1 Hz) and 2.38 ppm (1H, double doublet, *J*=14.5, 2.0 Hz), and 5-H signal was observed at 4.18 ppm (1H, double doublet, *J*=11.1, 2.0 Hz). The same procedure was carried out except for replacement of NaBH₄ with NaBD₄ and the reduced deuterium compound (**4b**) was obtained. Two kinds of methylene proton signals in the 2,2-dimethylpropyl group were detected at 1.79 (0.5H, doublet, *J*=11.0 Hz) and 2.36 ppm (0.5H, doublet, *J*=1.5 Hz) and two kinds of 5-H signals (0.5H, doublet, *J*=11.0 Hz and 0.5H, doublet, *J*=1.5 Hz) were observed at 4.18 ppm in the ¹H-NMR spectrum of **4b**. In addition, **4b** contained one deuterium atom based on the mass spectra (MS) (*M*⁺, 276). Consequently, it was confirmed that the reduced product (**4b**) was 5-(1-monodeuterio-2,2-dimethylpropyl)-3-methoxycarbonylmethylrhodanine. Since the reduction resulted in the formation of two asymmetric carbons, **4b**



	R ¹	R ²	yield of 2 (%)
a	CH(CH ₃) ₂	H	59.5
b	CH ₂ CH(CH ₃) ₂	H	31.5
c	CH(CH ₂ CH ₃) ₂	H	50.6
d	CH(CH ₃)CH ₂ CH ₃	H	23.2
e	C ₆ H ₁₁	H	63.6
f	(CH ₂) ₃ CH ₃	H	65.0
g	(CH ₂) ₅ CH ₃	H	41.2
h	(CH ₂) ₂ CH ₃	H	42.8
i	C(CH ₃) ₃	H	79.0
j	CH ₃	CH ₃	42.6
k	CH ₂ CH ₃	CH ₂ CH ₃	61.2
l ^{a)}	CH(CH ₃) ₂	CH ₃	49.7
m ^{a)}	CH ₂ CH ₃	CH ₃	34.3
n ^{a)}	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	81.4
o ^{a)}	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃	45.4

a) A mixture of the *E*- and *Z*-isomer.

Chart 1

TABLE I. Physical Data for the 5-Alkylmethyl-3-carboxymethylrhodanines (2)

Compd.	mp (°C)	¹ H-NMR (ppm, in CDCl ₃)	Formula and analysis		MS (<i>m/z</i>)	IR (cm ⁻¹)
			Calcd	Found		
			C	H		
2a	135—138	0.98 (6H, d, <i>J</i> = 5.2 Hz, CH ₃ × 2), 1.14—2.25 (3H, m, (CH ₃) ₂ CHCH ₂), 4.28 (1H, dd, <i>J</i> = 4.2, 10.0 Hz, 5-H), 4.75 (2H, s, CH ₂ COOH), 10.11 (1H, s, COOH)	C ₉ H ₁₃ NO ₃ S ₂ 43.71 (43.87	5.30 5.33)	247 (M ⁺), 229, 191	2860, 1707
2b	115—116.5	0.93 (6H, d, <i>J</i> = 6.0 Hz, CH ₃ × 2), 1.10—2.55 (5H, m, (CH ₃) ₂ CHCH ₂ CH ₂), 4.29 (1H, dd, <i>J</i> = 5.6, 8.0 Hz, 5-H), 4.80 (2H, s, CH ₂ COOH), 10.11 (1H, s, COOH)	C ₁₀ H ₁₅ NO ₃ S ₂ 45.96 (46.06	5.78 5.75)	261 (M ⁺), 243, 191	2880, 1710
2c	124—126	0.88 (6H, t, <i>J</i> = 6.0 Hz, CH ₃ × 2), 1.10—1.65 (5H, m, CH ₃ CH ₂ × 2 and (CH ₃ CH ₂) ₂ CH), 1.72—2.30 (2H, m, (CH ₃ CH ₂) ₂ CHCH ₂), 4.33 (1H, dd, <i>J</i> = 4.4, 10.2 Hz, 5-H), 4.72 (2H, s, CH ₂ COOH), 9.73 (1H, s, COOH)	C ₁₁ H ₁₇ NO ₃ S ₂ 47.98 (48.18	6.22 6.48)	275 (M ⁺), 257, 191	2860, 1730
2d	120—127	0.86—1.03 (6H, m, CH ₃ × 2), 1.23—2.56 (5H, m, CH ₃ CH ₂ CH(CH ₃)CH ₂), 4.21—4.48 (1H, m, 5-H), 4.82 (2H, s, CH ₂ COOH), 10.29 (1H, s, COOH)	C ₁₀ H ₁₅ NO ₃ S ₂ 45.96 (45.85	5.78 6.03)	261 (M ⁺), 243, 191	2870, 1740
2e	149—152	0.98—2.25 (13H, m, CH ₂ CHCH ₂ CH ₂ CH ₂ CH ₂ CH ₂), 4.29 (1H, dd, <i>J</i> = 4.4, 10.2 Hz, 5-H), 4.73 (2H, s, CH ₂ COOH), 10.39 (1H, s, COOH)	C ₁₂ H ₁₇ NO ₃ S ₂ 50.15 (49.86	5.96 6.05)	287 (M ⁺), 191	2840, 1708
2f	107—109	0.89 (3H, t, <i>J</i> = 5.0 Hz, CH ₃), 1.38 (6H, m, CH ₃ (CH ₂) ₃), 2.07 (2H, m, CH ₃ (CH ₂) ₃ CH ₂), 4.31 (1H, dd, <i>J</i> = 5.2, 8.0 Hz, 5-H), 4.80 (2H, s, CH ₂ COOH), 10.06 (1H, s, COOH)	C ₁₀ H ₁₅ NO ₃ S ₂ 45.96 (46.22	5.78 5.76)	261 (M ⁺), 243, 191	3050, 1732, 1713
2g	78—79	0.87 (3H, t, <i>J</i> = 4.2 Hz, CH ₃), 1.29 (10H, m, CH ₃ (CH ₂) ₅), 2.05 (2H, m, CH ₃ (CH ₂) ₅ CH ₂), 4.26 (1H, dd, <i>J</i> = 5.4, 8.2 Hz, 5-H), 4.73 (2H, s, CH ₂ COOH), 10.04 (1H, s, COOH)	C ₁₂ H ₁₉ NO ₃ S ₂ 49.80 (49.64	6.62 6.48)	289 (M ⁺), 171	3040, 1741, 1715
2h	143—145	0.92 (3H, t, <i>J</i> = 5.8 Hz, CH ₃), 1.42 (4H, m, CH ₃ (CH ₂) ₂), 2.11 (2H, m, CH ₃ (CH ₂) ₂ CH ₂), 4.38 (1H, dd, <i>J</i> = 5.2, 8.0 Hz, 5-H), 4.72 (2H, s, CH ₂ COOH), 8.93 (1H, s, COOH)	C ₉ H ₁₃ NO ₃ S ₂ 43.71 (43.56	5.30 5.13)	247 (M ⁺), 229, 191	3030, 1732, 1713
2i	165.5—167	1.00 (9H, s, CH ₃ × 3), 1.77 (1H, dd, <i>J</i> = 11.2, 15.0 Hz, (CH ₃) ₃ CCH ₂), 2.40 (1H, dd, <i>J</i> = 2.0, 15.0 Hz, (CH ₃) ₃ CCH ₂), 4.18 (1H, dd, <i>J</i> = 2.0, 11.2 Hz, 5-H), 4.79 (2H, s, CH ₂ COOH), 7.32 (1H, s, COOH)	C ₁₀ H ₁₅ NO ₃ S ₂ 45.96 (45.71	5.78 5.97)	261 (M ⁺), 243, 191	3050, 1743, 1726
2j	115—117	0.98 (3H, d, <i>J</i> = 6.0 Hz, CH ₃), 1.08 (3H, d, <i>J</i> = 6.0 Hz, CH ₃), 2.63 (1H, m, (CH ₃) ₂ CH), 4.32 (1H, d, <i>J</i> = 4.0 Hz, 5-H), 4.78 (2H, s, CH ₂ COOH), 10.14 (1H, s, COOH)	C ₈ H ₁₁ NO ₃ S ₂ 41.19 (41.03	4.75 4.61)	233 (M ⁺), 215, 191	2870, 1720
2k	105—107	0.90 (3H, t, <i>J</i> = 6.0 Hz, CH ₃), 1.00 (3H, t, CH ₃), 1.10—1.80 (4H, m, CH ₃ CH ₂ × 2), 2.19 (1H, m, (CH ₃ CH ₂) ₂ CH), 4.50 (1H, d, <i>J</i> = 3.6 Hz, 5-H), 4.75 (2H, s, CH ₂ COOH), 10.39 (1H, s, COOH)	C ₁₀ H ₁₅ NO ₃ S ₂ 45.96 (45.84	5.78 6.03)	261 (M ⁺), 191	3050, 1735
2l	143—147	0.91 (3H, d, <i>J</i> = 6.8 Hz, CH ₃), 0.98 (6H, d, <i>J</i> = 5.6 Hz, (CH ₃) ₂ CH), 1.25—2.50 (2H, m, CH × 2), 4.35—4.51 (1H, m, 5-H), 4.77 (2H, s, CH ₂ COOH), 10.34 (1H, s, COOH)	C ₁₀ H ₁₅ NO ₃ S ₂ 45.96 (46.03	5.78 5.83)	261 (M ⁺), 191	3050, 1740
2m	112—115	0.94 (3H, d, <i>J</i> = 7.0 Hz, CH ₃ CH <), 0.97 (3H, t, <i>J</i> = 6.4 Hz, CH ₃ CH ₂), 1.35 (2H, m, CH ₃ CH ₂), 2.43 (1H, m, CH ₃ CH <), 4.36—4.47 (1H, m, 5-H), 4.78 (2H, s, CH ₂ COOH), 10.33 (1H, s, COOH)	C ₉ H ₁₃ NO ₃ S ₂ 43.71 (43.60	5.30 5.02)	247 (M ⁺), 191, 173	3050, 1730
2n	122—124	0.92 (6H, t, CH ₃ × 2), 1.10—1.80 (8H, m, (CH ₃ CH ₂ CH ₂) ₂ CH), 2.35 (1H, m, (CH ₃ CH ₂ CH ₂) ₂ CH), 4.49 (1H, d, <i>J</i> = 3.6 Hz, 5-H), 4.80 (2H, s, CH ₂ COOH), 10.58 (1H, s, COOH)	C ₁₂ H ₁₉ NO ₃ S ₂ 49.80 (49.89	6.62 6.71)	289 (M ⁺), 191, 173	3050, 1734
2o	68—101	0.92 (6H, t, CH ₃ × 2), 1.10—1.70 (6H, m, CH ₃ CH ₂ CH ₂ CH(CH ₃ CH ₂)), 2.28 (1H, m, CH ₃ CH ₂ CH ₂ CH(CH ₃ CH ₂)), 4.50 (1H, d, <i>J</i> = 3.6 Hz, 5-H), 4.78 (2H, s, CH ₂ COOH), 10.16 (1H, s, COOH)	C ₁₁ H ₁₇ NO ₃ S ₂ 47.98 (47.75	6.22 5.92)	275 (M ⁺), 191	3050, 1710




Chart 2





TABLE II. Aldose Reductase-Inhibitory Activity

Compd.	IC ₅₀ (× 10 ⁻⁷ M)	Compd.	IC ₅₀ (× 10 ⁻⁷ M)
2a	1.1	2n	1.2
2b	2.0	2o	0.84
2c	0.79	(+)-2i	0.85
2d	1.1	(-)-2i	0.85
2e	0.90	4a	> 100
2f	0.63	6	> 100
2g	0.81	7	> 100
2h	1.8	8	> 100
2i	1.1	9	> 100
2j	1.9	10a	> 100
2k	3.1	10b	> 100
2l	0.76	10c	> 100
2m	2.0	Sorbinil	2.0

was obtained as a mixture of the two diastereomers in the ratio of 1:1. These results suggest that the reduction normally proceeds in the 1,4-addition process *via* an intermediate (A) (Chart 2).¹¹⁾

These 5-alkyl compounds (2) in the present report contain at least one asymmetric carbon atom. Resolution of 2 was carried out using (±)-5-(2,2-dimethylpropyl) compound (2i) as a typical compound. Treatment of 2i with one equivalent of D-3-amino-ε-caprolactam resulted in the formation of a salt (5a) in 60% yield. On the other hand, 2i was treated with 1 eq of L-3-amino-ε-caprolactam to give a salt (5b) in 64% yield. The enantiomeric acids, (+)-2i ([α]_D +69.4) and (-)-2i ([α]_D -68.7) were obtained from 5a and 5b respectively, in the usual manner. As 2i has acidic methine proton at 5-position, it is presumed that (±)-2i can be resolved by epimerization crystallization.¹²⁾ Attempts of the resolution of the acid (2i) using optical active α-methylbenzylamine, 1-(1-naphthyl)ethylamine and menthol were unsuccessful (Chart 3).

In addition, alkyl group could be introduced into 5-position of 2i to give 5-dialkylated compounds (6, 7) and their AR inhibition potency was checked. Alkylation of 2i with methyl iodide and isopropyl iodide gave corresponding 5-methyl-5-(1,1-dimethylpropyl) compound (6) and 5-isopropyl-5-(1,1-dimethylpropyl) compound (7) in poor yields (5–10%). Both methyl ester (4a) and isopropyl ester (8) were obtained as main products in these alkylations. Steric hindrance among the neopentyl group, sulfur atom at 1-position and carbonyl group at 4-position of 2i may make the replacement of 5-H by the alkyl groups difficult.¹³⁾

Ethyl ester (9) and some amides (10a–c) of 2i also were prepared to check their utility in *in vivo* screening (Chart 4).¹⁴⁾

AR Inhibitory Activity All of the rhodanine derivatives prepared were tested for their ability to inhibit AR

obtained from rat lens. The 50% inhibition of enzyme activity values are shown in Table II. All of the compounds (2a–o) displayed the potent inhibitory activity, with an IC₅₀ of 10⁻⁷–10⁻⁸ M. The *n*-pentyl derivative (2f) showed the highest activity among the compounds having five carbon chain at 5-position (2b, d, f, i, k, l). The branched isomers were less active than 2f, and 2k displayed the lowest activity (one fifth of the inhibitory activity of 2f) in this series. The compound (2b) was also 3 times less active than 2f.

Table II suggests that the activity of 2 is related to the magnitude of lipophilia of the substituted group at 5-position. It is noteworthy that the derivatives having five to seven carbon alkyl group at 5-position show the effective potency among the compounds in the present report: 2f, l having five carbon alkyl group, 2c, o having six carbon alkyl group and 2e, g having seven carbon alkyl group display the potent inhibitory activity, with an IC₅₀ of 10⁻⁸ M, but all of the compounds substituted with three or four carbon alkyl group exhibit less activity, with IC₅₀ of 10⁻⁷ M. The five to seven carbon alkyl groups at 5-position may interact appropriately with a secondary hydrophobic site presents on the enzyme to enhance the activity.¹⁵⁾ The 5-dialkylated compounds (6, 7) show over 10⁻⁵ M of IC₅₀. Steric hindrance caused by the geminal dialkyl group presumably makes the interaction with the hydrophobic site difficult.

The optically active (+)-2i and (-)-2i were as active as (±)-2i. It is speculated that the AR is not sensitive to optical activity of the compound (2i) or that the AR racemizes optically active 2i.

The rhodanine compounds (2) prepared by the reduction of the 5-*exo* double bond of 1 in this series had similar activity to the parent compounds (1).⁵⁾ Thus, it is assumed that bonding type of the 5-substitution group is independent of the affinity with the hydrophobic site.

Experimental

Melting points were determined in open capillaries with a Thomas-Hoover melting point apparatus and are uncorrected. Infrared (IR) spectra were recorded on a Shimadzu IR-17G spectrometer. ¹H-NMR spectra were measured on a JEOL PS-100 and a JNM-EX270 spectrometer with chemical shifts given in δ values with tetramethylsilane (TMS) as an internal standard. Low-resolution MS were obtained with a Hitachi M-52 instrument, and optical rotation was obtained with a Perkin-Elmer 241 polarimeter.

3-Carboxymethyl-5-(2,2-dimethylpropyl)rhodanine (2i) General Procedure for **2a–h** and **2j–o**: A solution of NaBH₄ (0.18 g, 4.62 mmol) in dry DMF (7.5 ml) was added dropwise to a solution of **1i** (1.5 g, 6.17 mmol) in dry DMF (7.5 ml) at 0°C, and the whole was stirred for 1 h at 0°C. The mixture was poured into ice-cold 2N HCl and extracted with ether. The extract was washed with saturated brine, dried over Na₂SO₄ and evaporated under reduced pressure. The yellow residue was crystallized from C₂H₄Cl₂–C₆H₁₂ (1:5) to give **2i** (1.2 g, 79%). Physical and spectral data are given in Table I.

3-Methoxycarbonylmethyl-5-(2,2-dimethylpropyl)rhodanine (4a) A mixture of **1i** (15 g, 57.9 mmol), *p*-toluenesulfonic acid monohydrate (1.5 g), and CH₃OH (20 ml) in dry C₆H₆ (150 ml) was stirred for 5 h at 80°C. After the usual workup, the product was recrystallized from ethyl acetate to give 5-*tert*-butylmethylidene-3-methoxycarbonylmethylrhodanine (**3**) (13.2 g, 83.5%) as colorless prisms, mp 66.0–68.0°C. ¹H-NMR (CDCl₃) δ : 1.24 (9H, s, C(CH₃)₃), 3.77 (3H, s, COOCH₃), 4.82 (2H, s, CH₂), 7.05 (1H, s, CH). MS m/z : 273 (M⁺), 213.

NaBH₄ (78 mg, 2.0 mmol) was added to a dry DMF solution (7 ml) of **3** (2.0 g, 7.3 mmol) at 0°C. The reaction mixture was stirred for 3 h at 0°C and then CH₃OH (3 ml) was added. After the mixture was adjusted to pH 7.0 with 2N HCl, the whole was concentrated under reduced pressure to give a pale yellow oil. The oil was purified by column chromatography (SiO₂, 5% ethyl acetate in C₆H₆) to give colorless prisms (**4a**, 1.3 g, 65%), mp 50.0–51.5°C. ¹H-NMR (CDCl₃) δ : 1.00 (9H, s, CH₃ × 3), 1.81 (1H, dd, J = 14.5, 11.1 Hz, one H of (CH₃)₃CCH₂), 2.37 (1H, dd, J = 14.5, 2.0 Hz, one H of (CH₃)₃CCH₂), 3.77 (3H, s, COOCH₃), 4.18 (1H, dd, J = 11.1, 2.0 Hz, 5-H), 4.75 (2H, s, NCH₂). MS m/z : 275 (M⁺), 243. Anal. Calcd for C₁₁H₁₇NO₃S₂: C, 47.98; H, 6.22. Found: C, 47.71; H, 6.35.

A Diastereomeric Mixture of 5-(1-Monodeuterio-2,2-dimethylpropyl)-3-methoxycarbonylmethylrhodanine (4b) NaBD₄ (84 mg, 2.0 mmol) was added to a dry DMF solution (7 ml) of **3** (2.0 g, 7.3 mmol) at 0°C. The reaction mixture was treated in a similar manner as preparation of **4a**. The deuterium compound (**4b**) was obtained as colorless prisms (1.2 g, 60%), mp 49–54°C. ¹H-NMR (CDCl₃) δ : 1.00 (9H, s, CH₃ × 3), 1.79 (0.5H, d, J = 11.0 Hz, (CH₃)₃CCHD), 2.36 (0.5H, d, J = 1.5 Hz, (CH₃)₃CCDH), 3.76 (3H, s, COOCH₃), 4.18 (0.5H, d, J = 11.0 Hz, 5-H), 4.18 (0.5H, d, J = 1.5 Hz, 5-H), 4.75 (2H, s, NCH₂). MS (m/z): 276 (M⁺), 244.

Optical Resolution of 2i 1) An aqueous solution (25 ml) of D-3-amino- ϵ -caprolactam (2.5 g, 191 mmol) was added slowly to a solution of **2i** (5.0 g, 191 mmol) in EtOH (5 ml) at room temperature. The mixture was allowed to stand for 5 h. The crystals were filtered and recrystallized from C₂H₄Cl₂–C₆H₁₂ (1:1) to give **5a** (4.5 g, 60%); mp 148.0–152.0°C, $[\alpha]_D^{25}$ –8.1° (c = 1.00, CHCl₃). ¹H-NMR (CDCl₃) δ : 1.00 (9H, s, CH₃ × 3), 1.40–2.45 (6H, m, NH₂CHCH₂CH₂CH₂), 1.77 (1H, dd, J = 14, 10 Hz, one H of (CH₃)₃CCH₂), 2.42 (1H, dd, J = 14, 2 Hz, one H of (CH₃)₃CCH₂), 3.07–3.45 (2H, m, CONHCH₂), 3.85–4.20 (1H, m, NH₂CH), 4.25 (1H, dd, J = 10, 2 Hz, SCHCO), 4.49 (2H, brs, NCH₂COOH), 7.56 (4H, brs, NH, NH₂, COOH).

2) **5b** was prepared with L-3-amino- ϵ -caprolactam by the same procedure as above, yield 63%; mp 152.0–156.0°C, $[\alpha]_D^{25}$ +7.3° (c = 1.00, CHCl₃). ¹H-NMR (CDCl₃) δ : 1.00 (9H, s, CH₃ × 3), 1.35–2.40 (6H, m, NH₂CHCH₂CH₂CH₂), 1.70 (1H, dd, J = 14, 10 Hz, one H of (CH₃)₃CCH₂), 2.34 (1H, dd, J = 14, 2 Hz, one H of (CH₃)₃CCH₂), 3.05–3.40 (2H, m, CONHCH₂), 3.88–4.15 (1H, m, NH₂CH), 4.22 (1H, dd, J = 10, 2 Hz, SCHCO), 4.50 (2H, brs, NCH₂COOH), 7.72 (4H, brs, NH, NH₂, COOH).

3) 0.1N HCl (25.7 ml) was added dropwise to an aqueous solution (200 ml) of **5a** (1.0 g, 2.57 mmol) at 0°C. The solution was extracted with ether, and the extract washed with saturated brine, dried over Na₂SO₄ and evaporated under reduced pressure. The residue was recrystallized from C₂H₄Cl₂–C₆H₁₂ (1:1) to give (+)-**2i** (0.52 g, 77%); mp 159.0–162.0°C, $[\alpha]_D^{25}$ +69.4° (c = 1.00, CHCl₃). ¹H-NMR (CDCl₃) δ : 0.99 (9H, s, CH₃ × 3), 1.78 (1H, dd, J = 10.2, 14.6 Hz, one H of (CH₃)₃CCH₂), 2.41 (1H, dd, J = 2.0, 14.6 Hz, one H of (CH₃)₃CCH₂), 4.19 (1H, dd, J = 2.0,

10.2 Hz, 5-H), 4.78 (2H, s, CH₂COOH), 9.56 (1H, s, COOH). MS m/z : 261 (M⁺), 243, 191. Anal. Calcd for C₁₀H₁₅NO₃S₂: C, 45.95; H, 5.78. Found: C, 46.20; H, 5.83.

4) (–)-**2i** was prepared from **5b** using the same procedure as above: yield 84%, mp 157.0–159.0°C, $[\alpha]_D^{25}$ –68.7° (c = 1.00, CHCl₃). ¹H-NMR (CDCl₃) δ : 0.99 (9H, s, CH₃ × 3), 1.78 (1H, dd, J = 10.2, 14.6 Hz, one H of (CH₃)₃CCH₂), 2.41 (1H, dd, J = 2.0, 14.6 Hz, one H of (CH₃)₃CCH₂), 4.19 (1H, dd, J = 2.0, 10.2 Hz, 5-H), 4.78 (2H, s, CH₂COOH), 9.01 (1H, s, COOH). MS m/z : 261 (M⁺), 243, 191. Anal. Calcd for C₁₀H₁₅NO₃S₂: C, 45.95; H, 5.78. Found: C, 45.94; H, 5.73.

3-Carboxymethyl-5-methyl-5-(2,2-dimethylpropyl)rhodanine (6) Methyl iodide (7.0 g, 49.3 mmol) was added to a mixture of **2i** (7.5 g, 29 mmol), acetic acid (2.6 g, 43.5 mmol) and sodium acetate (3.6 g, 43.5 mmol) in DMF (150 ml) at 30°C. The mixture was stirred at 50°C for 10 h, then the solvent was evaporated off under reduced pressure, and the residue was extracted with ether. The ether layer was washed with 0.5N HCl followed by brine, and dried over Na₂SO₄. Removal of the ether gave a pale yellow oil. The oil was purified by column chromatography on silica gel (8% ethyl acetate in hexane) to give **6** as pale yellow prisms (0.8 g, 10%) and **4a** (3.9 g, 49%). **6**: mp 85.5–88.0°C. ¹H-NMR (CDCl₃) δ : 0.88 (9H, s, CH₃ × 3), 2.28 (1H, d, J = 15 Hz, one H of (CH₃)₃CCH₂), 2.49 (1H, d, J = 15 Hz, one H of (CH₃)₃CCH₂), 3.38 (3H, s, 5-CH₃), 4.23 (2H, s, NCH₂), 7.81 (1H, brs, COOH). MS m/z : 275 (M⁺), 243. Anal. Calcd for C₁₁H₁₇NO₃S₂: C, 47.98; H, 6.22. Found: C, 47.70; H, 6.13.

3-Carboxymethyl-5-(2,2-dimethylpropyl)-5-isopropylrhodanine (7) Isopropyl iodide (7.0 g, 41.5 mmol) was added to a mixture of **2i** (5.0 g, 19.2 mmol), acetic acid (1.7 g, 28.8 mmol) and sodium acetate (2.4 g, 28.8 mmol) in DMF (120 ml) at 30°C. The mixture was stirred at 80°C for 15 h. After workup in a similar manner as for **6**, the residue was purified by column chromatography on silica gel (8% ethyl acetate in hexane) to give **7** (0.3 g, 5%) and 5-(2,2-dimethylpropyl)-3-isopropoxy-carbonylmethylrhodanine (**8**) (2.5 g, 43%) as yellow prisms. **7**: mp 96.0–98.0°C. ¹H-NMR (CDCl₃) δ : 0.98 (9H, s, (CH₃)₃C), 1.13 (6H, d, J = 6.0 Hz, (CH₃)₂CH), 1.72 (1H, d, J = 15.0 Hz, one H of CCH₂), 2.16 (1H, d, J = 15.0 Hz, one H of CCH₂), 4.20 (2H, s, NCH₂), 4.53 (1H, m, CH), 8.01 (1H, brs, COOH). MS m/z : 303 (M⁺), 286, 260. Anal. Calcd for C₁₃H₂₁NO₃S₂: C, 51.46; H, 6.98. Found: C, 51.60; H, 6.93. **8**: mp 72.5–75.0°C. ¹H-NMR (CDCl₃) δ : 0.90 (9H, s, (CH₃)₃C), 1.14 (6H, d, J = 7.0 Hz, CH(CH₃)₂), 1.50 (1H, dd, J = 14.5, 11.1 Hz, one H of CCH₂), 2.17 (1H, dd, J = 14.5, 2.0 Hz, one H of CCH₂), 3.76 (1H, dd, J = 11.1, 2.0 Hz, 5-H), 4.24 (2H, s, CH₂N), 4.58 (1H, m, CH(CH₃)₂), 8.00 (1H, brs, COOH). MS m/z : 303 (M⁺), 261. Anal. Calcd for C₁₃H₂₁NO₃S₂: C, 51.46; H, 6.98. Found: C, 51.37; H, 6.99.

3-Ethoxycarbonylmethyl-5-(2,2-dimethylpropyl)rhodanine (9) A mixture of **2i** (20.0 g, 76.5 mmol), absolute EtOH (20 ml) and *p*-toluenesulfonic acid (1.0 g) was refluxed for 8 h with azeotropic removal of water. The reaction mixture was washed successively with 2N HCl, saturated aqueous NaHCO₃ and saturated brine. The solution was dried over Na₂SO₄ and evaporated off under reduced pressure. The residue was recrystallized from EtOH–H₂O (5:1) to give **9** (18.0 g, 62%), mp 51.0–52.0°C. ¹H-NMR (CDCl₃) δ : 1.00 (9H, s, (CH₃)₃C), 1.27 (3H, t, J = 7.0 Hz, CH₃CH₂), 1.77 (1H, dd, J = 10.4, 14.4 Hz, one H of (CH₃)₃CCH₂), 2.40 (1H, dd, J = 2.0, 14.4 Hz, one H of (CH₃)₃CCH₂), 4.17 (1H, dd, J = 2.0, 10.4 Hz, 5-H), 4.21 (2H, q, J = 7.0 Hz, CH₃CH₂), 4.71 (2H, s, CH₂COO). MS m/z : 289 (M⁺). Anal. Calcd for C₁₂H₁₉NO₃S₂: C, 49.80; H, 6.62. Found: C, 49.66; H, 6.65.

3-Diethylaminocarbonylmethyl-5-(2,2-dimethylpropyl)rhodanine (10a) Isobutyl chloroformate (1.0 g, 7.61 mmol) was added slowly to a mixture of **2i** (2.0 g, 7.56 mmol), triethylamine (0.8 g, 7.71 mmol) and dry THF (14 ml) at –13°C, and the mixture was stirred for 15 min at –13°C. A solution of diethylamine (0.6 g, 7.66 mmol) in CHCl₃ (14 ml) was added dropwise and the resulting mixture was stirred for 1 h at 0°C and then 1 h at room temperature. The reaction mixture was poured into cold water and extracted with ethyl acetate. The extract was washed successively with 2N HCl, saturated aqueous NaHCO₃ and saturated brine. The solution was dried over Na₂SO₄ and evaporated off under reduced pressure. The residual oil was distilled under reduced pressure to give **10a** (1.4 g, 58%), bp 260°C/0.1 mmHg. ¹H-NMR (CDCl₃) δ : 1.00 (9H, s, (CH₃)₃C), 1.12 (3H, t, J = 7.6 Hz, CH₃CH₂), 1.31 (3H, t, J = 7.6 Hz, CH₃CH₂), 1.82 (1H, dd, J = 10.0, 14.8 Hz, one H of (CH₃)₃CCH₂), 2.41 (1H, dd, J = 2.0, 14.8 Hz, one H of (CH₃)₃CCH₂), 3.40 (4H, q, J = 7.6 Hz, CH₃CH₂ × 2), 4.22 (1H, dd, J = 2.0, 10.0 Hz, 5-H), 4.81 (2H, s, CH₂CO). MS m/z : 316 (M⁺), 301, 283. Anal. Calcd for C₁₄H₂₄N₂O₂S₂: C, 53.13; H, 7.64. Found: C, 52.86; H, 7.55.

3-Dimethylaminocarbonylmethyl-5-(2,2-dimethylpropyl)rhodanine (10b)

The amide (**10b**) was prepared with dimethylamine using a procedure similar to that above, mp 104.0–105.5°C from (C₂H₄Cl₂–C₆H₁₂ (1:1)), yield 82%. ¹H-NMR (CDCl₃) δ: 1.00 (9H, s, (CH₃)₃C), 1.81 (1H, dd, *J* = 10.0, 14.8 Hz, one H of (CH₃)₃CCH₂), 2.41 (1H, dd, *J* = 2.0, 14.8 Hz, one H of (CH₃)₃CCH₂), 2.98 (3H, s, NCH₃), 3.10 (3H, s, NCH₃), 4.20 (1H, dd, *J* = 2.0, 10.0 Hz, 5-H), 4.81 (2H, s, CH₂CO). MS *m/z*: 288 (M⁺). Anal. Calcd for C₁₂H₂₀N₂O₂S₂: C, 49.97; H, 6.99. Found: C, 50.23; H, 6.73.

5-(2,2-Dimethylpropyl)-3-piperidylcarbonylmethylrhodanine (10c)

The amide (**10c**) was prepared with piperidine by use of a similar procedure as above, mp 142.0–144.0°C from (C₂H₄Cl₂–C₆H₁₂ (1:1)), yield 72%. ¹H-NMR (CDCl₃) δ: 1.00 (9H, s, (CH₃)₃C), 1.65–2.02 (7H, m, NCH₂CH₂CH₂CH₂CH₂ and one H of (CH₃)₃CCH₂), 2.39 (1H, dd, *J* = 2.0, 14.4 Hz, one H of (CH₃)₃CCH₂), 3.48 (4H, brs, NCH₂CH₂CH₂CH₂CH₂), 4.17 (1H, dd, *J* = 2.0, 10.0 Hz, 5-H), 4.79 (2H, s, CH₂CO). MS *m/z*: 328 (M⁺), 313, 295. Anal. Calcd for C₁₅H₂₄N₂O₂S₂: C, 54.85; H, 7.36. Found: C, 55.13; H, 7.18.

Enzyme Inhibitory Activity Aldose reductase activity was measured by the method of Hoyman and Kinoshita.¹⁶ Assays were performed at 30°C in 0.1 M sodium phosphate buffer (pH 6.2) containing 1.5 mM DL-glyceraldehyde, 0.25 mM reduced nicotinamide adenine diphosphate (NADPH) and an appropriate amount of enzyme (supernatant of homogenates of rat lens) in a total volume of 1.5 ml. The effect of an inhibitor on the enzyme activity was determined by adding 15 μl of dimethylsulfoxide solution of a test compound to the reaction mixture. The concentration of the inhibitor giving IC₅₀ was estimated from the least-squares regression line in the plot of the logarithm of inhibition concentration versus remaining activity.

Acknowledgement The authors wish to thank Professor Shunsaku Ohta of Kyoto Pharmaceutical University for his valuable advice.

References and Notes

- 1) D. Dvornik, N. Simard-Duquesne, M. Krami, K. Sestanj, K. H. Gabbay, J. H. Kinoshita, S. D. Varma, and L. O. Merola, *Science*, **182**, 1146 (1973).
- 2) C. A. Lipinski and N. J. Hutson, *Ann. Rep. Med. Chem.*, **19**, 169 (1984); P. F. Kador, J. H. Kinoshita, and N. E. Sharpless, *J. Med. Chem.*, **28**, 841 (1985); J. DeRuiter, R. F. Borne, and C. A. Mayfield, *J. Med. Chem.*, **32**, 145 (1989); J. DeRuiter, B. E. Swearingen, V. Wandrekar, and C. A. Mayfield, *ibid.*, **32**, 1033 (1989); J. Wrobel, J. Millen, J. Sredy, A. Dietrich, J. M. Kelly, B. J. Gorham, and K. Sestanj, *ibid.*, **32**, 2493 (1989); B. L. Mylari, E. R. Larson, T. A. Beyer, W. J. Zembrowski, C. E. Aldinger, M. F. Dee, T. W. Siegal, and D. H. Singleton, *J. Med. Chem.*, **34**, 108 (1991).
- 3) a) R. J. C. Guy, S. G. Gilbey, M. Sheehy, P. Asselman, and P. J. Watkins, *Diabetologia*, **31**, 214 (1988); A. A. F. Sima, V. Bril, V. Nathaniel, J. A. J. McEwen, M. B. Brown, S. A. Lattimer, and D. A. Greene, *New. Engl. J. Med.*, **319**, 548 (1988); b) D. G. Shand, *Diabetic Med.*, **3**, 559A (1986); Y. Harati, E. Niakan, J. P. Comstock, and C. Logon, *Ann. Neurol.*, **22**, 129 (1987); c) N. Sakamoto, J. H. Kinoshita, P. F. Kador, and N. Hotta, *Excerpta Medica Congress Series*, **760**, 422 (1988); *idem, ibid.*, **760**, 451 (1988).
- 4) Y. Ohishi, T. Mukai, M. Nagahara, M. Yazima, and N. Kajikawa, *Chem. Pharm. Bull.*, **37**, 2398 (1989).
- 5) Y. Ohishi, T. Mukai, M. Nagahara, M. Yazima, and N. Kajikawa, *Chem. Pharm. Bull.*, **38**, 1911 (1990).
- 6) H. C. Brown, "Boranes in Organic Chemistry," Cornell University Press, Ithaca, N.Y., 1972, p. 209.
- 7) H. Shechter, D. E. Ley, and E. B. Robertson, Jr., *J. Am. Chem. Soc.*, **78**, 4984 (1956); H. H. Baer and W. Rank, *Can. J. Chem.*, **50**, 1292 (1972).
- 8) H. Le Moal, R. Carrie, and M. Bargain, *Comp. Rend.*, **251**, 2541 (1960); S. S. Kulp and C. B. Caldwell, *J. Org. Chem.*, **45**, 171 (1980); S. S. Kulp and R. Szorko, *ibid.*, **53**, 5573 (1988).
- 9) K. Adank, H. A. Pfenniger, W. G. Stoll, and M. Viscontini, *Helv. Chim. Acta*, **46**, 1030 (1963); J. Bottin, O. Eisenstein, C. Minot, and N. T. Anh, *Tetrahedron Lett.*, **1972**, 3015.
- 10) a) S. B. Kadin, *J. Org. Chem.*, **31**, 620 (1966); b) J. H. Schauble, G. J. Walter, and J. G. Morin, *J. Org. Chem.*, **39**, 755 (1974).
- 11) Schauble *et al.* reported a mechanistic aspect of the nucleophilic 1,4-addition of NaBH₄ to methyl α-(*p*-nitrophenyl)-*trans*-cinnamate (ref. 10b).
- 12) H. Leuches and J. Wutke, *Ber.*, **46**, 2425 (1913); H. Leuche, *ibid.*, **54**, 830 (1921); W. C. Ashley and R. L. Shriner, *J. Am. Chem. Soc.*, **54**, 4410 (1932); M. K. Hargreaves and M. A. Khan, *J. Chem. Soc., Perkin Trans. 2*, **1973**, 1204; M. A. Kahn and A. M. Kahn, *J. Chem. Soc., Perkin Trans. 1*, **1979**, 75; K. Arai, Y. Ohata, K. Takakuwa, and T. Iizumi, U. S. Patent 4417070 (1983) [*Chem. Abstr.*, **100**, 138788c (1984)].
- 13) There may be steric repulsion among the *tert*-butyl group in the neopentyl moiety, hydrogen at 5-position, sulfur at 1-position and carbonyl group at 4-position of **2i** considering the space-filling model (CPK precision molecular models). The repulsion of the 5-dialkyl compounds (**6**, **7**) may be much larger than that of **2i**.
- 14) These data will be published and discussed elsewhere.
- 15) J. DeRuiter and C. A. Mayfield, *J. Pharm. Sci.*, **76**, 149 (1987).
- 16) S. Hoyman and J. H. Kinoshita, *J. Biol. Chem.*, **240**, 877 (1965).