## Direct, efficient, and inexpensive formation of $\alpha$ -hydroxyketones from olefins by hydrogen peroxide oxidation catalyzed by the 12-tungstophosphoric acid/cetylpyridinium chloride system

Yanfei Zhang,<sup>a</sup> Zongxuan Shen,<sup>a</sup> Jingting Tang,<sup>a</sup> Yan Zhang,<sup>a</sup> Lichun Kong<sup>b</sup> and Yawen Zhang<sup>\*a</sup>

Received 6th January 2006, Accepted 7th February 2006 First published as an Advance Article on the web 13th March 2006 DOI: 10.1039/b518200j

The direct ketohydroxylation of a variety of 1-aryl-1alkenes with  $H_2O_2$ , catalyzed by the inexpensive 12tungstophosphoric acid/cetylpyridinium chloride system under very mild conditions, was achieved. Various acyloins were obtained in good yields and high regioselectivies.

 $\alpha$ -Hydroxyketones (acyloins) are important building blocks in organic synthesis and form parts of the structures of various biologically active compounds, such as cortisone acetate 1 and daunomycinone  $2^{1,2}$  (Fig. 1).



Fig. 1 Acyloin substructure in cortisone acetate 1 and daunomycinone 2.

It is known that acyloins can be prepared by the benzoin condensation, the acyloin condensation, the  $\alpha$ -hydroxylation of carbonyl compounds and the oxidation of diols, whereas the direct oxidation of C–C double bonds has only rarely been reported.<sup>3</sup> Since, in these cases, toxic cyanide compounds must be employed, or the reaction must be performed under an inert atmosphere with dry solvent, the development of an efficient and convenient method to obtain  $\alpha$ -hydroxyketones would be significant. Recently, a new straightforward oxidation of C–C double bonds to  $\alpha$ -hydroxyketones using catalytic amounts of RuCl<sub>3</sub> and stoichiometric amounts of Oxone under buffered conditions has been reported by Plietker.<sup>4</sup> A variety of substituted olefins were oxidized to  $\alpha$ -hydroxyketones in good-to-excellent regioselectivities and yields by this method.

Heteropoly acids (HPA) such as 12-molybdophopsphoric acid (MPA) or 12-tungstophosphoric acid (WPA) are often used, not only for the oxidation of organic substrates, but also for many acid-catalyzed reactions, because they possess the dual catalytic functions of oxidizing ability and strong acidity.<sup>5</sup> It has been reported that the combination of HPA with cetylpyridinium chloride (CPC) could catalyze the epoxidation of olefins such as 1-octene, the ketonization of alcohols and diols, and the oxidative cleavage of 1,2-diols and olefins.<sup>5</sup>

We have found that direct ketohydroxylation of some olefins with  $H_2O_2$  could be achieved under similar conditions. It turns out that when olefins are treated with hydrogen proxide in chloroform containing a catalytic amount of WPA (10 mol%) combined with CPC (30 mol%) as phase-transfer catalyst at 60 °C,  $\alpha$ -hydroxyketones can obtained in good-to-excellent yields with good regioselectivities (Scheme 1). There are a number of interesting features of this methodology worth pointing out: (i)  $H_3PW_{12}O_{40}$  is less expensive and less toxic than RuO<sub>4</sub> or OsO<sub>4</sub>; (ii)  $H_2O_2$  as the oxidant is of great advantage to the environment and industry because it generates  $H_2O$  as the sole by-product, it has a high content of active oxygen species, and it is less expensive than organic peroxides and peracids; (iii) the experiment is easy to carry out, because no special precautions such as high temperature, inert gas atmosphere or dry solvents are necessary.

$$R^{1} \xrightarrow{R^{2}} H_{3}PW_{12}O_{40} (10 \text{ mol}\%) \xrightarrow{O} R^{2}$$

$$CPC (30 \text{ mol}\%) \xrightarrow{R^{2}} R^{1} OH$$

$$H_{2}O_{2} (30\%, 10 \text{ eq})$$

$$CHCI_{3}, 60 \ ^{\circ}C$$

Scheme 1

We choose styrene as our working model to study the influences of the catalyst, phase-transfer catalyst, solvents and temperatures (Table 1).

Table 1 Effects of the catalyst, solvent, and temperature on the ketohydroxylation of styrene"

-	-			
Entry	Catalyst	Solvent	T∕°C	Yield (%) <sup>b</sup>
1	WPA/CPC	CHCl <sub>3</sub>	60	86.4
2	WPA	CHCl <sub>3</sub>	60	nd
3	$CWP^{c}$	CHCl <sub>3</sub>	60	42.4
4	WPA/CPC	$CH_2Cl_2$	40	62.5
5	WPA/CPC	CH <sub>3</sub> CN	60	36.4
6	WPA/CPC	PhCH <sub>3</sub>	60	73.9
7	WPA/CPC	H <sub>2</sub> O	60	nd
8	WPA/CPC	CHCl <sub>3</sub>	-20	nd
9	WPA/CPC	CHCl,	4	10

 $^a$  All reactions were performed on a 6.5 mmol scale in CHCl<sub>3</sub> (15 mL) using WPA (H<sub>3</sub>PW<sub>12</sub>O<sub>40</sub>, 0.4 mol%), CPC (1.2 mol%) and 10 equiv. of H<sub>2</sub>O<sub>2</sub> (30%), and were stopped after 24 h.  $^b$  Isolated yields (nd = not determined).  $^c$  CWP: prepared according to ref. 5.

<sup>&</sup>lt;sup>a</sup>Key Laboratory of Organic Synthesis of Jiangsu Province, College of Chemistry and Chemical Engineering, Suzhou University, Suzhou, 215123, China. E-mail: zhangyw@suda.edu.cn; Fax: +86 0512 65880305; Tel: +86 0512 65880339

<sup>&</sup>lt;sup>b</sup>College of Chemistry and Life Science, Zhejiang Normal University, Jinhua, 321004, China. E-mail: sky35@zjnu.cn

To evaluate the potential of  $H_3PW_{12}O_{40}$  for the ketohydroxylation of styrene, three different catalyst systems were tested (entries 1–3). Combination of WPA with CPC produced *in situ* was found to be more effective than  $[\pi$ -C<sub>6</sub> $H_5NC_{16}H_{33}]_3[PO_4(WO_3)_4]$  (CWP).

When the WPA/CPC combination (which produces CWP with evolution of hydrogen chloride) was employed as the catalyst, the reaction medium is acidic, while in the CWP-catalyzed reaction, the medium is not acidic. The fact that ketohydroxylation proceeded more rapidly with the WPA/CPC combination suggested that the tungsten-catalyzed reaction is accelarated in acidic media. WPA alone could not catalyze the ketohydroxylation of styrene (entry 2), probably due to its poor solubility in organic solvents. Therefore, the reaction was conducted in various solvents using the WPA/CPC catalyst. It was observed that chloroform was the best solvent. The same reaction in toluene or acetonitrile led to a considerable decrease of the yield (entries 1, 4–7). A lower temperature decreased the conversion significantly (entry 9), while no reaction was found at -20 °C.

It is noteworthy that the insoluble WPA/CPC dissolved slowly under the action of  $H_2O_2$ , to form the real catalyst, which subsequently led to homogeneous catalytic direct oxidation of C–C double bond. During the reaction, the system gradually changed from turbid to clear. The infrared spectra (Fig. 2) showed that the  $H_2O_2$ -treated WPA/CPC (spectrum 2) has a peroxo-bond absorption peak v(O–O) at 840 cm<sup>-1</sup> and 846 cm<sup>-1</sup>, while CWP itself (spectrum 1) does not have a peroxo-bond absorption peak at about 840 cm<sup>-1</sup>. Instead, an absorption peak at 895 cm<sup>-1</sup>, which corresponds to the asymmetric vibration of W–O–W, was found.



Fig. 2 Infrared spectra of (1) CWP, (2)  $H_2O_2$  (30%)-treated CWP, and (3) recovered  $H_2O_2$  (30%)-treated catalyst.

After extraction of the product, the aqueous phase was evaporated, which allowed the recovery of the catalyst. As shown in Table 2, the recovered catalyst can be recycled at least once without significant loss of activity.

 Table 2
 Recycling the catalyst for the ketohydroxylation of styrene

Run "	Yield (%) <sup>b</sup>	
1	86.4	
2	81.8	
3	71.5	
4	64.7	

<sup>*a*</sup> All reactions were performed under the conditions of entry 1, Table 1, except that the catalyst was recovered from the preceding run. <sup>*b*</sup> Isolated yield.



Scheme 2 A proposed mechanism.

A mechanistic explanation is proposed as shown in Scheme 2. According to previous mechanistic research on the WPA/CPCcatalyzed epoxidation of olefins with hydrogen peroxide, when mixed with commercially available 35%  $H_2O_2$ ,  $H_3[PW_{12}O_{40}]$  is oxidized to  $\{PO_4[WO(O_2)_2]_4\}^{3-}$ , which was identified as the real catalyst.7 This peroxo tungsten complex behaves as a 1,3-dipolar reagent, M<sup>+</sup>-O-O<sup>-</sup>, which reacts with the olefin to form a fivemembered dioxametallocycle adduct. Degradation of the adduct produces the corresponding epoxide<sup>8</sup> and a PW<sub>3</sub> species, which quickly turns to the peroxo tungsten catalyst in the presence of hydrogen peroxide. The electrophilic attack of the peroxo tungsten complex on the above epoxide results in the formation of a six-membered cyclic intermediate, which degrades to produce the final oxidation product, a hydroxyketone, together with a  $PW_3$  moiety, which is oxidized back to the peroxo tungsten catalyst and then enters the next cycle. As evidence for the above mechanism, phenyloxirane could be oxidized to the corresponding hydroxyketone in high yield under the reaction conditions (entry 19, Table 3). The fact that 1,1-disubstituted and trisubstituted ethylenes did not give hydroxyketones is another piece of evidence for this, because, according to the above mechanistic explanation, a hydrogen atom  $\alpha$ - to the phenyl group is necessary. In the absence

 Table 3
 WPA/CPC-catalyzed oxidation of various olefins<sup>a</sup>

 Entry	Substrate	Product	Time/h	Yield (%) <sup>b</sup>
1°	Ph	0 	24	86.4
	1	Ph CH <sub>2</sub> OH		
2		<b>2</b> O	48	74.6
	3	Ph		
		ОН <b>4</b>		
3	Ph		24	75.7
	5	Ph j O Ph OH		
4	$\sim$	<b>6</b> O	48	73.0
	Ph SO <sub>2</sub> Ph	Ph SO <sub>2</sub> Ph		
		ОН 8		
5	Ph	$\hat{\square}$	48	78.5
	9	Ph OAc OH		
6	$\sim$ $\sim$	10 O	17	75.6
	Ph <sup>2</sup> N <sub>3</sub> 11	Ph N <sub>3</sub>		
		ОН <b>12</b>		
7			20	79.4
	Ph <sup>-</sup> N	Ph N OH		
	0 13	0 14		
8	PhOPh		48	20, 22
	15	F''                   OPh         F''                   OPh           O         OH         0         0         0         0         0         0         0         0         0         0         0         0         0         16a         16b         16b         16b         16b         0		
9	CH <sub>3</sub>	CH3	48	76.2
	p-F-C <sub>6</sub> H <sub>4</sub> -Cl-o	p-F-C <sub>6</sub> H <sub>4</sub>		
10	17	18 O	12	76.7
	Ph	Ph 20		
11	<b>19</b> Ph	2 <b>U</b>	12	80.7
		Ph		
	21	22		

## Table 3 (Contd.)



<sup>*a*</sup> All reactions were performed on a 2 mmol scale in CHCl<sub>3</sub> (15 mL) using WPA (H<sub>3</sub>PW<sub>12</sub>O<sub>40</sub>, 10 mol%), CPC (30 mol%), and 10 equiv. of H<sub>2</sub>O<sub>2</sub> (30%) at 60 °C. <sup>*b*</sup> Isolated yields (nd = not determined). <sup>*c*</sup> The reaction was carried out under the conditions given in Table 1.

of WPA, styrene was not oxidized at all, even after 20 h of stirring with CPC/hydrogen peroxide at 60 °C. This excludes a radical pathway.

However, when extended to substrates other than styrene, the above conditions did not work. It was found that the amount of WPA and CPC must be increased to 10 mol% and 30 mol%, respectively, for the reaction to proceed at an acceptable rate.

We investigated the direct oxidation of various olefins under the modified conditions.<sup>6</sup> Table 3 shows the results of an intense screening of scope and limitations. It is important that many electron-rich alkenes were successfully transformed into the corresponding acyloins with remarkably high regioselectivities and yields (entries 1–7). We were even not able to detect any regioisomeric products from the crude mixtures. All the substrates that gave hydroxyketones as the products were 1-aryl-1-alkenes. In other words, an aromatic ring attached to the terminal sp<sup>2</sup> carbon is necessary for the hydroxyketonization to occur. Without this kind of substituent, only epoxidation was observed (entries 12– 14). 1,1-Disubstituted and trisubstituted ethylenes cleaved during the course of oxidation (entries 9–11). It is noteworthy that the conversion is strongly dependent on the electronic properties of the C–C double bond. In general, electron-rich substrates are ketohydroxylated much faster than electron-poor starting materials; in fact, no reaction was observed with those alkenes that had moderately strong electron-withdrawing groups on C2 (entries 15–18).

In conclusion, we have demonstrated that various 1-aryl-1alkenes can be oxidized by  $H_2O_2$  to hydroxyketones in the presence of WPA/CPC under mild reaction conditions.

## Acknowledgements

The authors are grateful to the Key Laboratory of Organic Synthesis of Jiangsu Province, China, for financial support.

## **References and notes**

- 1 L. F. Fieser and M. Fieser, in: *Steroids*, Reinhold Publishing, New York, 1959, p. 600.
- 2 T. Li and Y.-L. Wu, J. Am. Chem. Soc., 1981, 103, 7007.
- 3 (a) S.-I. Murahashi, Pure Appl. Chem., 1992, 64, 403; (b) Rutheniumcatalyzed reactions: S.-I. Murahashi, T. Saito, H. Hanaoka, Y. Murakami, T. Naota, H. Kumobayashi and S. Akutagawa, J. Org. Chem., 1993, 58, 2929; (c) S.-I. Murahashi, T. Naota and H. Hanaoka, Chem. Lett., 1993, 1767; (d) T. Takai, T. Yamada and T. Mukaiyama, Chem.

Lett., 1991, 1499; (e) S. Baskaran, J. Das and S. Chandrasekaran, J. Org. Chem., 1989, 54, 5182.

- 4 (a) B. Plietker, J. Org. Chem., 2003, 68, 7123; (b) B. Plietker, M. Niggemann and A. Pollrich, Org. Biomol. Chem., 2004, 2, 1116; (c) B. Plietker, Org. Lett., 2004, 6, 289; (d) B. Plietker, J. Org. Chem., 2004, 69, 8287.
- 5 Y. Ishii, K. Yamawaki, T. Ura, H. Yamada, T. Yoshida and M. Ogawa, *J. Org. Chem.*, 1988, **53**, 3587.
- 6 General procedure for the oxidation of olefins: To a stirred mixture of 12-tungstophosphate acid (0.60 g, 0.21 mmol), CPC (0.21 g, 0.62 mmol) and 30%  $H_2O_2$  (2.0 mL, 19.7 mmol) in chloroform (15 mL) in a 100 mL round-bottomed flask equipped with a magnetic stirring bar was added alkene (2 mmol). The mixture was allowed to react

under reflux, and was monitored by TLC. After the reaction was complete, the reaction mixture was filtered, and the filtrate treated with a solution of 10% sodium hydrogen sulfite (to decompose unreacted H<sub>2</sub>O<sub>2</sub>) and then with 10% sodium hydroxide. The product was extracted with chloroform (15 mL  $\times$  3), and the organic phase was washed with brine (100 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to a colorless oil. Pure product was obtained by silica gel column chromatography (petroleum ether–ethyl acetate, 10 : 1–2, as the eluent).

- 7 D. C. Duncan, R. C. Chambers, E. Hecht and C. L. Hill, J. Am. Chem. Soc., 1995, 117, 681.
- 8 (a) H. Mimoun, Angew. Chem., Int. Ed. Engl., 1982, 21, 734; (b) H. Mimoun, J. Mol. Catal., 1980, 7, 1.