

BIBLIOGRAPHIE

- [1] W. D. Weringa, D. H. Williams, J. Feeny, J. P. Brown & R. W. King, J. chem. Soc., Perkin Trans. 1, 443 (1972); R. M. Smith, R. W. Johnson & R. D. Guthrie, Chem. Commun. 1972, 361; G. Lukacs & R. M. Smith, Bull. Soc. chim. France 10, 3995 (1972).
- [2] S. W. Gunner, R. D. King, W. G. Overend & N. R. Williams, J. chem. Soc. [C], 1954 (1970).
- [3] H. Yanagisawa, M. Kinoshita & S. Umezawa, Bull. Soc. chim. Japan 42, 1719 (1969).
- [4] H. Yanagisawa, M. Kinoshita & S. Urnezawa, Bull. Soc. chim. Japan 43, 246 (1970).
- [5] J. M. J. Tronchet, J. M. Bourgeois, Carbohydrate Res. 29, 373 (1973).
- [6] A. Rosenthal, K. S. Ong, D. Baker, Carbohydrate Res. 13, 113 (1970).

297. Stereospecific Synthesis of a 9, 11, 12, 13, 13a, 14-hexahydro-dibenzo(f, h)pyrrolo(1,2-b)isoquinoline-alkaloid


Preliminary communication

by Ludwig Faber and Wolfgang Wiegrebe

Pharmazeutisches Institut, Berne

(12. XI. 73)

Several syntheses of the title-type alkaloids starting from chiral proline-derivatives, are described in the literature [1-4]. All these routes produce racemates because they include intermediates with a carbonyl-group α to the asymmetrically substituted carbon-atom C(13a). We avoided racemisation by using the proline derivative III [5] and cyclising the phenanthrene-moiety at a late stage of our synthetic route, shown in the following schemes:




*) diastereomers were isolated +) mixture of diastereomers R = OCH₃

We obtained III by LiBH_4 -reduction of 5-oxo-proline methyl ester to 5-oxo-prolinol which was subsequently tosylated. The optical purity of 5-oxo-prolinol was only 51% of that of *Hardegger's* [5] product, who used LiAlH_4 . This disadvantage is compensated by a better yield of 5-oxo-prolinol, as compared with the LiAlH_4 -reduction.

Anisole + homoveratrotylchloride $\rightarrow$ I [6]. – I + 65% HNO_3 in glacial $\text{AcOH} \rightarrow$ II, m.p. 197° ; NMR. (chem. shift in δ): H_1 : 6.80, 1s; H_2 : 7.83, 1s; $\text{H}_{3a,b}$: 8.09, 2d $\times$ d, $J_0 = 7$ Hz, $J_m = 1.7$ Hz; $\text{H}_{4a,b}$: 7.03, 2d $\times$ d, $J_0 = 7$ Hz, $J_m = 1.7$ Hz. – II + III in acetone/ K_2CO_3 , 110° , 4–5 atm $\rightarrow$ IV; yellow oil; M^+ : 428. NMR.: H_5 : 5.67, 1t, $J = 7.5$ Hz; $[\alpha]_D^{20} = +22^\circ$ (MeOH). – IV + NaBH_4 (70% EtOH) $\rightarrow$ V. Va, low Rf, NMR.: H_7 : 4.74, 1d, $J = 7$ Hz; H_8 : 3.00, 1s. – Vb, high Rf, NMR.: H_7 : 4.77, 1d, $J = 7$ Hz; H_8 : 3.00, 1s. – V (a or b) + 1.25% HCl in glacial AcOH (20°) $\rightarrow$ VI (a or b respectively). VIa (high Rf); m.p. 152° ; phenylgroups *cis*; NMR.: H_6 : 5.08, 1d, $J = 8.5$ Hz; $[\alpha]_D^{19} = +89^\circ$ (CHCl_3). – VIb (low Rf); m.p. 194° ; phenylgroups *trans*; NMR.: H_6 : 4.83, 1d, $J = 1.7$ Hz; $[\alpha]_D^{20} = +43.5^\circ$ (CHCl_3). – VI (a or b) + H_2 (Raney-Ni/EtOH) $\rightarrow$ VII (a or b respectively). MS.: M^+ : 382, dominating peaks at 203 and 175. – VII (mixture of diastereomers) + NaNO_2 /2.5% aqueous HCl , followed by 30% $\text{H}_3\text{PO}_2 \rightarrow$ VIII. MS. of both diastereomers: M^+ : 367, dominating peaks 203 and 175. – VIII (mixture of diastereomers) + LiAlH_4 /THF $\rightarrow$ IX; diastereomers (colourless oils) were separated. IXa (low Rf), $[\alpha]_D^{25} = -214^\circ$ (MeOH); IXb (high

Scheme 2



$\text{R} = \text{OCH}_3$

Rf), $[\alpha]_D^{25} = -173^\circ$ (MeOH). MS. (both diastereomers): M^+ 353, base peak: 189. – IX (mixture of a and b) + BrCN in abs. $C_6H_6 \rightarrow$ X. IR. ($CHCl_3$): 2200/cm. MS.: M^+ : 378, dominating peaks 283 and 95. – X (4.6×10^{-5} M in C_6H_{12} , saturated with air) was irradiated (254 nm) $\rightarrow$ XI; m.p. 151–2°; UV.: λ_{max} (log ϵ): 257 (4.73), 285 (4.44), 311 (3.90), 342 (2.89), 359 nm (2.31); M^+ : 376; $[\alpha]_D^{25} = +74.7$ ($CHCl_3$). – XI + $LiAlH_4$ in THF $\rightarrow$ XII (viscous oil). M^+ : 351, small intensity [7], dominating peaks 349, 282 and 70; $[\alpha]_D^{25} = +8.1^\circ$. – XII + 98% $HCOOH/180^\circ \rightarrow$ XIII; cyclisation (without further purification) with $POCl_3$ /toluene $\rightarrow$ XIV; UV.: λ_{max} : 259; 268; 282; 320; 420 nm. XIV + $NaBH_4$ /MeOH $\rightarrow$ XV, m.p. 206–11°. $[\alpha]_D^{25} = +66^\circ$ ($CHCl_3$); antofine: $[\alpha]_D^{25} = -131^\circ$ ($CHCl_3$). XV was identical with natural antofine [8] (TLC., UV., MS., IR.) but shows opposite optical rotation, 50% in magnitude for the reason stated. Because XV has S-configuration, antofine is of the R-configuration. This is in accord with our degradation of antofine to D-proline [9].

REFERENCES

- [1] B. Chauncy, E. Gellert & K. N. Trivedi, Austral. J. Chemistry, 22, 427 (1969).
- [2] R. B. Herbert & C. J. Moody, Chem. Commun. 1970, 121.
- [3] T. R. Govindachari, B. R. Pai, S. Prabhakar & T. S. Savitri, Tetrahedron 21, 2573 (1965).
- [4] B. Chauncy & E. Gellert, Austral. J. Chemistry, 23, 2503 (1970).
- [5] E. Hardegger & H. Ott, Helv. 38, 313 (1955).
- [6] A. Novelli & A. de Santis, Bol. Soc. Quim. Peru 30, (4) 155 (1964); Chem. Abstr. 64, 5069a.
- [7] W. Wiegand, L. Faber & H. Budzikiewicz, Liebigs Ann. Chem. 733, 125 (1970).
- [8] W. Wiegand, L. Faber, H. Brockmann jr., H. Budzikiewicz & U. Krüger, Liebigs Ann. Chem. 721, 154 (1969).
- [9] W. Wiegand, L. Faber & Th. Breyhan, Arch. Pharmaz. 304, 188 (1971).

298. Zur Photochemie von $\Delta^{1,9}$ -10-Methyl-2-octalonvon Paul Margaretha¹⁾ und Kurt Schaffner

Département de Chimie Organique, Université de Genève, 1211 Genève 4

(13. XI. 73)

Summary. Some aspects of the photochemistry of $\Delta^{1,9}$ -10-methyl-2-octalone (**1**) were investigated. In view of the hitherto neglected substantial thermal reversibility of the photolytic double bond shift under previous conditions of analysis (1) the solvent-dependent selectivity in product formation in C_6H_6 and *t*-BuOH was confirmed, (2) the essentially exclusive double bond shift and its intermolecular nature in C_6F_6 were established, (3) the quantum yields of the major products in these three solvents were measured, and (4) the dependence of triplet quenching data on temperature, enone concentration and conversion was shown in C_6F_6 for the double bond shift. In conclusion, the previously published experimental basis for the postulate of two differentially reactive triplet states of **1** is insufficient, although the postulate itself is not necessarily disproved.

Das photochemische Verhalten der Titelverbindung **1** ist bereits von mehreren Arbeitsgruppen untersucht worden. Zimmerman & al. [1] bestimmten für die Umlagerung von **1** in das «Lumiketon» **2** Quantenausbeuten von $3,8 \cdot 10^{-3}$ in *t*-Butylalkohol und $6,1 \cdot 10^{-3}$ in Benzol. Schaffner & al. [2] hingegen fanden ausgeprägte Lösungsmittelleffekte auf die photochemische Umsetzung von **1**: in *t*-Butylalkohol wurden überwiegend **2** und **3**²⁾, in Benzol hingegen bevorzugt das β,γ -ungesättigte Keton

¹⁾ Beurlaubt vom Max-Planck-Institut für Kohlenforschung, Abt. Strahlenchemie, Mülheim a.d. Ruhr.

²⁾ Der prozentuale Anteil des Photoisomeren **3** ist vernachlässigbar klein.