

Green Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: J. Schwarz and B. Koenig, *Green Chem.*, 2016, DOI: 10.1039/C6GC01101B.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Metal-Free, Visible-Light-Mediated, Decarboxylative Alkylation of Biomass-Derived Compounds†

Johanna Schwarz and Burkhard König^a

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

This work describes a mild, environmentally friendly method to activate natural carboxylic acids for decarboxylative alkylation. After esterification of biomass-derived acids to *N*-(acyloxy)phthalimides, the active esters are cleaved reductively by photocatalysis to give alkyl radicals, which undergo C–C bond formation with electron-deficient alkenes. This reaction is catalyzed by the organic dye eosin Y and green light (535 nm) and the scope of acids includes abundant amino acids, α -oxy acids and fatty acids which are available from renewable resources.

Introduction

Carboxylic acids are among the most abundant, renewable feedstocks on our planet. They are non-toxic, stable and inexpensive and therefore valuable starting materials for “green chemistry”.¹ Due to shortage of fossil resources and rising energy demand, alternative sources of raw materials gain in importance.² In order to generate platform chemicals and high-value chemicals like pharmaceuticals from biomass-derived compounds, the carboxy group can be targeted as chemo- and regioselective leaving group for C–C bond formation reactions.³


In principle, decarboxylative reactions are known for a long time, but old protocols (e.g. according to Hunsdiecker⁴, Barton⁵ and Kolbe⁶) are not suitable for cross-coupling reactions between acids and other substrates under benign conditions. Over the last two decades, many transition metal-catalysed, decarboxylative C–C and C–X coupling methods were reported by Gooßen,⁷ Myers⁸ and many others.⁹ Even though these methods are versatile in application and often compatible with multiple step reactions, they require high temperatures as well as palladium or copper reagents. Moreover, they are usually limited to decarboxylation of C_{sp}²–COOH or C_{sp}–COOH bonds.

Recently, first photoredox-mediated processes were developed for the activation of the carboxy group. The usage of photocatalysts and visible light enables reactions under mild conditions and low energy consumption. During the last years, photocatalytic, decarboxylative reactions like arylations,¹⁰ vinylations,¹¹ allylations,¹² alkynylations,¹³ fluorinations¹⁴ and hydrodecarboxylations¹⁵ have been reported by MacMillan and other groups.¹⁶ Also different methods for decarboxylative alkylations have already been investigated.¹⁷ Here, alkyl radicals


which are generated from carboxylic acids upon extrusion of CO₂ react with activated alkenes, typically Michael acceptors. The alkyl radicals can either be formed by esterification of carboxylic acids and subsequent reductive cleavage of the ester bond (Scheme 1a)^{17c, 18} or by oxidative cleavage of the acid itself (Scheme 1b).^{17a} These methods have already been reported by different groups, but expensive and toxic transition metal catalysts are needed in all cases and the scope of carboxylic acids is limited.

Herein, we report for the first time a metal-free, photocatalytic, decarboxylative alkylation which is applicable for a broad variety of natural carboxylic acids involving amino acids, α -oxy acids and fatty acids (Scheme 2). Therefore, *N*-(acyloxy)-phthalimides were synthesized from the corresponding acid according to a simple method developed by Okada,^{18b, 19} analogously to Overman's approach. These active esters undergo C–C bond formation with electron-deficient alkenes under irradiation with green light in the presence of DIPEA and the organic, non-toxic and cheap dye eosin Y. Thus, valuable chemicals can be obtained from renewable biomass under eco-friendly and mild conditions.

a) Reductive decarboxylation according to Overman *et al.*^{17c}



b) Oxidative decarboxylation according to MacMillan *et al.*^{17a}




Scheme 1 Photocatalytic generation of alkyl radicals from carboxylic acids for subsequent coupling with electron-deficient alkenes.

^a Institute of Organic Chemistry, University of Regensburg, Universitätsstraße 31, 93053 Regensburg, Germany

E-mail: Burkhard.Koenig@chemie.uni-regensburg.de; Phone: (+49)-941-943-4575.

† Electronic Supplementary Information (ESI) available: synthesis of starting materials, spectroscopic investigations, analytical data and ¹H- and ¹³C-NMR spectra. See DOI: 10.1039/x0xx00000x




Scheme 2 Our work: general approach for the decarboxylative alkylation of natural compounds by reductive cleavage of *N*-(acyloxy)phthalimide **1**

Results and discussion

Synthesis and scope

For investigation of the reaction conditions of the decarboxylative alkylation, the *N*-(acyloxy)phthalimide of *N*-Boc-protected proline (**1a**) and *n*-butyl acrylate (**2a**) served as test substrates. A mixture of both compounds, the base DIPEA and a homogeneous photocatalyst was irradiated with LEDs under nitrogen atmosphere. For this reaction, no heterogeneous catalysts have been investigated, although an application would be conceivable.²⁰ First, the catalytic activity of the photocatalysts eosin Y (**A**) and [Ru(bpy)₃]Cl₂ (**B**) in different solvents was screened (Table 1). The inexpensive organic dye **A** enabled higher yields than metal catalyst **B** in

Table 1 Evaluation of catalyst and solvent.^a



Entry	Photocatalytic system	Solvent	Yield [%] ^b
1	[Ru(bpy) ₃]Cl ₂ (2 mol%, 455 nm)	DMSO	30
2	[Ru(bpy) ₃]Cl ₂ (2 mol%, 455 nm)	DMF	47
3	[Ru(bpy) ₃]Cl ₂ (2 mol%, 455 nm)	CH ₃ CN	60
4	[Ru(bpy) ₃]Cl ₂ (2 mol%, 455 nm)	CH ₂ Cl ₂	18
5	[Ru(bpy) ₃]Cl ₂ (2 mol%, 455 nm)	THF	7
6	Eosin Y (10 mol%, 535 nm)	DMSO	30
7	Eosin Y (10 mol%, 535 nm)	DMF	37
8	Eosin Y (10 mol%, 535 nm)	CH ₃ CN	65
9	Eosin Y (10 mol%, 535 nm)	CH ₂ Cl ₂	96
10	Eosin Y (10 mol%, 535 nm)	THF	53

^a Reactions were performed using 1 equiv. **1a**, 5 equiv. **2a** and 2 equiv. DIPEA.


^b Determined by GC analysis using naphthalene as internal standard.

almost every solvent (except DMF: Table 1, entries 2 and 7). As [Ru(bpy)₃]Cl₂ is known to be a stronger reductant than eosin Y,²¹ the reason for the better performance of the metal-free catalyst **A** is probably its better solubility and stability in these solvents. Decomposition of **B** was indicated by darkening of the reaction mixtures after several minutes. The best product yield of 96% was obtained with photocatalyst **A** and CH₂Cl₂ as solvent (Table 1, entry 9). For further optimization, different reaction conditions and control reactions were investigated (Table 2). Reducing the amount of catalyst to 5 mol% (Table 2, entry 1) or the amount of alkene from 5 to 2 eq. (Table 2, entry 4) resulted in lower yields of about 85%. Also a shorter reaction time (Table 2, entry 2) or the use of less base (Table 2, entry 3) had a clearly negative impact. However, running the reaction in the presence of air (Table 2, entry 5) gave the same yield as after degassing the reaction mixture (Table 1, entry 9). Measuring the oxygen concentration during the non-degassed reaction showed that the oxygen was consumed within two hours (see Supporting Information). Control experiments without light, DIPEA or catalyst led to no product formation at all, independent if oxygen was present or not (Table 2, entry 6 to 11).

Table 2 Optimization of reaction conditions.

Entry	Modification	Yield [%] ^a
1	5 mol% A	85
2	9 h irradiation time	64
3	1 equiv. DIPEA	26
4	2 equiv. 2a	83
5	Air atmosphere ^b	96
6	No base	0
7	No light	0
8	No photocatalyst	0
9	Air atmosphere, no base	0
10	Air atmosphere, no light	0
11	Air atmosphere, no photocatalyst	0

^a Determined by GC analysis using naphthalene as internal standard.

Table 3 Scope of electron-deficient alkenes.^a Product not isolated. Yield estimated by $^1\text{H-NMR}$.

With these optimized reaction conditions, the scope and limitations of the decarboxylative alkylation were explored. Therefore, the reaction of different electron-deficient alkenes with active ester **1a** was investigated (Table 3). The expected cross coupling products were observed in moderate to excellent yields when α,β -unsaturated esters and ketones were used as coupling partners. Unsubstituted or β -methylated aliphatic Michael acceptors gave the desired products **3a**, **3b** and **3e** in good yields of 73 to 80%. Benzylic, α,β -unsaturated esters **2g** and **2h** were suitable reaction partners in the same way. Also reactions with cyclic ketones gave products **3d** and **3e** in about 75% yield. However, introduction of a phenyl ring at the attacked carbon inhibited formation of product **3l** almost completely, presumably due to steric hindrance by the bigger substituent in β -position. In contrast, if the phenyl ring was positioned on the α -carbon of the Michael acceptor, the best yield of 92% (**3i**) was observed. This result indicates that after attack of the *N*-Boc proline fragment at the double bond, a radical is formed at the α -position, which is stabilized by an adjacent phenyl ring in the case of **3i**. Next to Michael acceptors, also the electrophilic, heteroaromatic styrene derivative vinylpyridine (**2k**) yielded the corresponding product **3k** in 40%.

After establishing the scope of activated alkenes, *N*-(acyloxy)phthalimide **1** was varied (Table 4). Here, benzyl methacrylate **2h** served as model alkene and again, the optimized conditions of Table 2, entry 5 were applied. A broad variety of esterified acids, including readily available, biomass-derived compounds could be employed. This photocatalytic


cross-coupling reaction works on the one hand for primary (**1c**, **1k**, **1n-s**), secondary (**1b**, **1c-j**, **1l**) as well as for tertiary carboxylic acids (**1i**) and on the other hand for different substrate classes like amino acids (**1b-g**), α -oxy acids (**1h-j**) and even fatty acids (**1n-s**).

The active esters of simple, Boc-protected, secondary amino acids proline (**1a**, Table 3), alanine (**1b**) or valine (**1d**) gave the highest yields (68-89%). The *N*-(acyloxy)phthalimide of the primary amino acid glycine showed slightly less product formation (**4c**, 62%). Also amino acid derivatives with aromatic side chains like phenylalanine (**1e**) or benzyl-protected serine (**1f**) and aspartic acid (**1g**) yielded the corresponding products **4e-g** in about 60%. As a typical natural α -hydroxy acid, lactic acid with different protecting groups was investigated. Here, the phenoxy- (**1i**) and methoxy-protected derivatives (**1j**) gave with 50 and 58% yield, respectively, less coupling product than cyclic α -oxy starting material **1h** (72%).

Applying active esters of simple carboxylic acids without any heteroatom in α -position to the carboxy moiety also resulted in product formation, although the yields decreased to 30-45% (**4k-s**). This decline in efficiency indicates that the stability of the alkyl radical generated after decarboxylation is the decisive point for successful product formation. In the case of amino and α -oxy compounds, the radical is stabilized by electron-donation from the lone pair of the heteroatom to the single-occupied p orbital of the neighboring radical centered carbon, which explains the higher yields.²² Considering the decarboxylative alkylation of *N*-(acyloxy)phthalimides of simple carboxylic acids, secondary (**1l**, 40%) or even more tertiary carboxylic acids (**1m**, 45%) are more prone to decarboxylate than primary, linear carboxylic acids (**1k**, **1n-s**; all about 30%). Again, tertiary alkyl radicals are more stable than secondary and especially primary radicals, which is in good accordance with the observed yields.^{22b} Nonetheless, several fatty acids, which are widespread in nature could be applied for this photocatalytic reaction. Saturated (**1n-q**), mono-unsaturated (**1r**) and di-unsaturated (**1s**) fatty acid derivatives with different chain lengths (12-18 C) all gave products **4n-s** in about 30% yield. The reaction with the active ester of cinnamic acid (**1t**) yielded no product, because the vinylic radical intermediate, which is formed upon decarboxylation is apparently not stable enough to undergo further reactions. In summary, we could show that cheap and abundant compounds from renewable biomass are suitable starting materials for this cross-coupling reaction.

Mechanistic investigations

Based on the report by Overman *et al.*^{17c}, the known photochemistry of eosin Y^{23} and our own results, we propose the following mechanism for the decarboxylative alkylation of *N*-(acyloxy)phthalimides with electron-deficient alkenes (Scheme 3): Irradiation of a photocatalyst (PC) with visible light generates the excited catalyst PC^* , which is reductively quenched by the sacrificial electron donor DIPEA to give $\text{DIPEA}^{+\bullet}$. Regeneration of the PC is presumably achieved by reduction of *N*-(acyloxy)phthalimide **1a**, which gives the corresponding radical anion. Splitting of the N–O bond and


Table 4 Scope of *N*-(acyloxy)phthalimides.View Article Online
DOI: 10.1039/C6GC01101B

subsequent elimination of CO₂ generates alkyl radical **1a**[•]. This radical can attack the double bond of Michael acceptor **2a** to form the intermediate **3a**[•]. It is assumed that hydrogen atom abstraction from DIPEA^{••} or the solvent finally enables formation of product **3a**.

This mechanistic proposal was confirmed by several spectroscopic studies. Single electron transfer (SET) from DIPEA to the excited PC should be exergonic according to the redox potentials of DIPEA (+0.72 V vs. SCE in CH₃CN)²⁴ and the excited

PC **A** (eosin Y[•]/eosin Y^{••}: +0.83 V vs. SCE in CH₃CN-H₂O, 1:1)^{21b} or **B** (Ru²⁺*/Ru^{•+}: +0.77 V vs. SCE in CH₃CN)²⁵ and is a well documented process in literature.^{21, 26} Furthermore, this step was confirmed by luminescence quenching experiments with both photocatalysts. In the case of eosin Y, fluorescence quenching could hardly be observed with DIPEA, *N*-(acyloxy)phthalimide **1a** or alkene **2a** (see Supporting Information for graphs and further details). This indicates that excited PC **A** does rather react from the triplet than from the singlet state which

Scheme 3 Proposed reaction mechanism for the photocatalytic, decarboxylative alkylation of *N*-(acyloxy)phthalimide **1a** with *n*-butylacrylate (**2a**).



was already shown by our group for another reaction.²⁷ For verification of this triplet reactivity, transient absorption measurements were performed, due to the short triplet lifetime of eosin Y ($\tau_T = 320 \pm 10$ ns).^{27a} Performing fluorescence experiments with [Ru(bpy)₃]Cl₂ (**B**) revealed that the emission of **B**^{*} is quenched by the electron donor DIPEA, which proves SET between both species (Figure 1). Upon titration of **B**^{*} with **1a** or **2a**, no quenching was observed (see Supporting Information).

The next step of the catalytic cycle was assumed to be regeneration of the PC by SET from PC^{•+} to **1a**. Cyclic voltammetry measurements of the active ester **1a** (-1.20 V vs.

Figure 1 Fluorescence quenching of [Ru(bpy)₃]Cl₂ (**B**, 15.0 μ M in CH₃CN) upon titration with DIPEA (100 mM in CH₃CN). DOI: 10.1039/C6GC01101B

SCE in CH₃CN, see Supporting Information) showed that reduction by the strongly reducing PC **B** (Ru⁺/Ru²⁺: -1.33 V vs. SCE in CH₃CN)^{21a} is thermodynamically feasible. Although redox potentials are not directly comparable with each other in the case of PC **A** (eosin Y^{•+}/eosin Y: -1.06 V vs. SCE in CH₃CN-H₂O, 1:1),^{21b} this catalyst is also known to be a powerful reductant. Eosin Y shows an absorption maximum at 527 nm which corresponds to an energy of 2.35 eV. After intersystem crossing to its reactive triplet state, 1.89 eV are still available for photocatalytic reactions.^{21b} To confirm the formation of radical **3a**[•] during the reaction, this intermediate was captured by the persistent radical TEMPO giving product **5** (Scheme 3), which was determined by LC-MS analysis (see Supporting Information). Furthermore, the quantum yield of the photocatalytic model reaction was determined (see Supporting Information). The low value of $\Phi = 2.6 \pm 0.5\%$ corresponds to the relatively long reaction times of 18 h and indicates that efficient radical chain processes are very unlikely in this mechanism.²⁸ For further insight, the stability of the photocatalyst eosin Y has been investigated by measuring UV-vis absorption spectra during the reaction (see Supporting Information). Over the reaction time of 18 h, the catalyst slowly degrades and a precipitate occurs, which slows down the product formation after some hours (see Supporting Information for the time course of the reaction). However, the remaining photocatalytic active species is sufficient to reach almost complete conversion within the reaction time.


Experimental

General procedure for the synthesis of *N*-(acyloxy)phthalimides **1**

Starting materials **1** were synthesized by a slightly modified procedure based on Reiser *et al.*^{18a} and Overman *et al.*^{17c} The respective carboxylic acid (8.00 mmol, 1.0 equiv.), *N*-hydroxyphthalimide (1.43 g, 8.80 mmol, 1.1 equiv.), *N,N'*-dicyclohexylcarbodiimide (1.98 g, 9.60 mmol, 1.2 equiv.) and 4-dimethylaminopyridine (0.98 g, 0.80 mmol, 0.1 equiv.) were mixed in a flask with a magnetic stirring bar. Dry THF (40 mL) was added and the orange reaction mixture was stirred for 15 h at rt. The white precipitate was filtered off and the solution was concentrated by evaporation of the solvent. Purification by column chromatography on flash silica gel (CH₂Cl₂ or CH₂Cl₂/CH₃OH = 9:1) gave the corresponding product.

General procedure for the photocatalytic decarboxylative alkylation

In a 5 mL crimp cap vial with a stirring bar, eosin Y (**A**, 19.4 mg, 0.03 mmol, 0.1 equiv.) and *N*-(acyloxy)phthalimide **1** (0.30 mmol, 1.0 equiv.) were added. After addition of DIPEA (102 μ L, 0.60 mmol, 2.0 equiv.), the corresponding olefin **2** (1.50 mmol, 5.0 equiv.) and dry CH₂Cl₂ (4 mL), the vial was capped to prevent evaporation. The reaction mixture was stirred and irradiated through the vials' plane bottom side using



green LEDs (535 nm) for 18 h at rt. The reaction mixture of two vials with the same content was combined and diluted with saturated aqueous solution of NaHCO₃ (20 mL). It was extracted with EA (3 x 20 mL) and the combined organic phases were washed with brine (20 mL), dried over Na₂SO₄ and concentrated in vacuum. Purification of the crude product was performed by automated flash column chromatography (PE/EA = 19:1 to 1:1) yielding the corresponding product **3** or **4** as colorless oil.

Conclusions

In conclusion, we have developed a metal-free, photocatalytic method for the decarboxylative alkylation of biomass-derived compounds. The advantage of this procedure over other methods reported in literature is the broad substrate scope including cheap and abundant α -amino acids, α -oxy acids and fatty acids as well as primary, secondary and tertiary substrates. In addition, the reactions can be carried out under mild, environmentally friendly conditions with the metal-free, organic dye eosin Y as photocatalyst. The carboxylic acids are activated by esterification to *N*-(acyloxy)phthalimides and then reductively cleaved upon irradiation with green light. This generates alkyl radicals, which undergo cross-coupling with electron-deficient alkenes. Thereby, largely new compounds for the synthesis of pharmaceuticals or fine chemicals are generated. The method contributes to the ongoing efforts replacing fossil resources by renewable feedstocks in the synthesis of chemical intermediates.

Acknowledgements

We thank the German Science Foundation (DFG, GRK 1626, Chemical Photocatalysis) for financial support for this work.

References

- 1 P. T. Anastas and J. C. Warner, *Green Chemistry: Theory and Practice*, Oxford University Press, New York, 1998.
- 2 (a) A. Pfennig, *Chem. Ing. Tech.*, 2007, **79**, 2009-2018; (b) R. Watson, in *Ressourcenverknappung, 50 Schlüsselideen der Zukunft*, Springer Berlin Heidelberg, 2014, pp. 24-27.
- 3 E. Scott, F. Peter and J. Sanders, *Appl. Microbiol. Biotechnol.*, 2007, **75**, 751-762.
- 4 H. Hunsdiecker and C. Hunsdiecker, *Chem. Ber.*, 1942, **75**, 291-297.
- 5 (a) D. H. R. Barton, D. Bridon, I. Fernandez-Picot and S. Z. Zard, *Tetrahedron*, 1987, **43**, 2733-2740; (b) D. H. R. Barton, D. Crich and W. B. Motherwell, *J. Chem. Soc., Chem. Commun.*, 1983, 939-941.
- 6 (a) H. Kolbe, *Liebigs Ann. Chem.*, 1849, **69**, 257-294; (b) H. Kolbe, *Liebigs Ann. Chem.*, 1848, **64**, 339-341.
- 7 (a) L. J. Gooßen, G. Deng and L. M. Levy, *Science*, 2006, **313**, 662-664; (b) N. Rodriguez and L. J. Goossen, *Chem. Soc. Rev.*, 2011, **40**, 5030-5048.
- 8 (a) A. G. Myers, D. Tanaka and M. R. Mannion, *J. Am. Chem. Soc.*, 2002, **124**, 11250-11251; (b) D. Tanaka, S. P. Romeril and A. G. Myers, *J. Am. Chem. Soc.*, 2005, **127**, 10323-10333.
- 9 R. Shang and L. Liu, *Sci. China Chem.*, 2011, **54**, 1670-1687.
- 10 (a) Z. Zuo and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2014, **136**, 5257-5260; (b) Z. Zuo, H. Cong, W. Li, J. Choi, G. C. Fu and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2016, **138**, 1832-1835; (c) Z. Zuo, D. T. Ahneman, L. Chu, J. A. Terrett, A. G. Doyle and D. W. C. MacMillan, *Science*, 2014, **345**, 437-440; (d) Y. Jin, M. Jiang, H. Wang and H. Fu, *Scientific Reports*, 2016, **6**, 20068.
- 11 (a) A. Noble and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2014, **136**, 11602-11605; (b) A. Noble, S. J. McCarver and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2015, **137**, 624-627.
- 12 (a) C. Hu and Y. Chen, *Org. Chem. Front.*, 2015, **2**, 1352-1355; (b) S. B. Lang, K. M. O'Nele, J. T. Douglas and J. A. Tunge, *Chem. Eur. J.*, 2015, **21**, 18589-18593; (c) S. B. Lang, K. M. O'Nele and J. A. Tunge, *J. Am. Chem. Soc.*, 2014, **136**, 13606-13609.
- 13 (a) J. Yang, J. Zhang, L. Qi, C. Hu and Y. Chen, *Chem. Comm.*, 2015, **51**, 5275-5278; (b) F. Le Vaillant, T. Courant and J. Waser, *Angew. Chem. Int. Ed.*, 2015, **54**, 11200-11204; (c) H. Huang, G. Zhang and Y. Chen, *Angew. Chem. Int. Ed.*, 2015, **54**, 7872-7876; (d) Q.-Q. Zhou, W. Guo, W. Ding, X. Wu, X. Chen, L.-Q. Lu and W.-J. Xiao, *Angew. Chem. Int. Ed.*, 2015, **54**, 11196-11199.
- 14 (a) S. Ventre, F. R. Petronijevic and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2015, **137**, 5654-5657; (b) X. Wu, C. Meng, X. Yuan, X. Jia, X. Qian and J. Ye, *Chem. Commun.*, 2015, **51**, 11864-11867.
- 15 (a) C. Cassani, G. Bergonzini and C.-J. Wallentin, *Org. Lett.*, 2014, **16**, 4228-4231; (b) J. D. Griffin, M. A. Zeller and D. A. Nicewicz, *J. Am. Chem. Soc.*, 2015, **137**, 11340-11348.
- 16 J. Xuan, Z.-G. Zhang and W.-J. Xiao, *Angew. Chem. Int. Ed.*, 2015, **54**, 15632-15641.
- 17 (a) L. Chu, C. Ohta, Z. Zuo and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2014, **136**, 10886-10889; (b) D. W. Manley, R. T. McBurney, P. Miller, R. F. Howe, S. Rhydderch and J. C. Walton, *J. Am. Chem. Soc.*, 2012, **134**, 13580-13583; (c) G. Pratsch, G. L. Lackner and L. E. Overman, *J. Org. Chem.*, 2015, **80**, 6025-6036; (d) Y. Miyake, K. Nakajima and Y. Nishibayashi, *J. Am. Chem. Soc.*, 2012, **134**, 3338-3341; (e) Y. Miyake, K. Nakajima and Y. Nishibayashi, *Chem. Comm.*, 2013, **49**, 7854-7856.
- 18 (a) G. Kachkovskyi, C. Faderl and O. Reiser, *Adv. Synth. Catal.*, 2013, **355**, 2240-2248; (b) K. Okada, K. Okamoto, N. Morita, K. Okubo and M. Oda, *J. Am. Chem. Soc.*, 1991, **113**, 9401-9402.
- 19 K. Okada, K. Okubo, N. Morita and M. Oda, *Tetrahedron Lett.*, 1992, **33**, 7377-7380.
- 20 (a) M. A. Fox and M. T. Dulay, *Chem. Rev.*, 1993, **93**, 341-357; (b) U. I. Gaya, *Heterogeneous Photocatalysis*

Using Inorganic Semiconductor Solids, Springer Netherlands, 2014; (c) B. Kraeutler and A. J. Bard, *J. Am. Chem. Soc.*, 1978, **100**, 5985-5992; (d) M. Schiavello, *Electrochim. Acta*, 1993, **38**, 11-14; (e) X. Ning, S. Meng, X. Fu, X. Ye and S. Chen, *Green Chem.*, 2016; (f) Q. Wang, J. Li, Y. Bai, J. Lian, H. Huang, Z. Li, Z. Lei and W. Shangguan, *Green Chem.*, 2014, **16**, 2728-2735; (g) J. Yuan, J. Wen, Y. Zhong, X. Li, Y. Fang, S. Zhang and W. Liu, *J. Mater. Chem.*, 2015, **3**, 18244-18255; (h) J.-J. Zhou, R. Wang, X.-L. Liu, F.-M. Peng, C.-H. Li, F. Teng and Y.-P. Yuan, *Appl. Surf. Sci.*, 2015, **346**, 278-283.

- 21 (a) J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102-113; (b) D. P. Hari and B. König, *Chem. Comm.*, 2014, **50**, 6688-6699.
- 22 (a) J. Hioe and H. Zipse, *Org. Biomol. Chem.*, 2010, **8**, 3609-3617; (b) M. L. Coote, C. Y. Lin and H. Zipse, *The Stability of Carbon-Centered Radicals*, John Wiley & Sons, 2009; (c) J. S. Wright, H. Shadnia and L. L. Chepelev, *J. Comput. Chem.*, 2009, **30**, 1016-1026.
- 23 M. Majek, F. Filace and A. J. v. Wangelin, *Beilstein J. Org. Chem.*, 2014, **10**, 981-989.
- 24 (a) V. V. Pavlishchuk and A. W. Addison, *Inorg. Chim. Acta*, 2000, **298**, 97-102; (b) G. J. Barbante, N. Kebede, C. M. Hindson, E. H. Doeven, E. M. Zammit, G. R. Hanson, C. F. Hogan and P. S. Francis, *Chem. Eur. J.*, 2014, **20**, 14026-14031.
- 25 C. Creutz and N. Sutin, *Inorg. Chem.*, 1976, **15**, 496-499.
- 26 (a) D. M. Schultz and T. P. Yoon, *Science*, 2014, **343**; (b) K. Zeitler, *Angew. Chem.*, 2009, **121**, 9969-9974; (c) J. Hu, J. Wang, T. H. Nguyen and N. Zheng, *Beilstein J. Org. Chem.*, 2013, **9**, 1977-2001.
- 27 (a) A. U. Meyer, T. Slanina, C.-J. Yao and B. König, *ACS Catalysis*, 2015, 369-375; (b) A. Penzkofer, A. Beidoun and M. Daiber, *J. Lumin.*, 1992, **51**, 297-314.
- 28 M. A. Cismesia and T. P. Yoon, *Chem. Sci.*, 2015, **6**, 5426-5434.
- 29

View Article Online
DOI: 10.1039/C6GC01101B