

RSC Advances



This article can be cited before page numbers have been issued, to do this please use: C. Luo, G. Parthasarathy, Y. Wu, W. Chen and C. H. Cheng, *RSC Adv.*, 2015, DOI: 10.1039/C5RA23065A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Copper Promoted Synthesis of Substituted Quinolines from Benzylic Azides and Alkynes

Ching-Zong Luo, Parthasarathy Gandeepan, Yun-Ching Wu, Wei-Chen Chen and Chien-Hong Cheng*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/


A novel copper promoted synthesis of substituted quinolines from various benzylic azides and internal alkynes has been demonstrated. The reaction features a broad substrate scope, high product yields and excellent regioselectivity. In contrast to the known two-step process of acid promoted [4+2] cycloaddition reaction and oxidation, the present methodology allows the synthesis of quinolines in a single step under neutral reaction conditions and can be applied to the synthesis of biologically active 6-chloro-2,3-dimethyl-4-phenylquinoline (antiparasitic agent) and 3,4-diphenylquinolin-2(1*H*)-one (p38 α MAP kinase inhibitor). A plausible reaction mechanism involves rearrangement of benzylic azide to *N*-arylimine (Schmidt reaction) followed by intermolecular [4+2] cyclo addition with internal alkynes.

Introduction


Quinoline and its derivatives are potential heterocyclic scaffold found in many natural products¹⁻³, and bio-active molecules.⁴⁻¹⁰ They are also key structures in materials, agrochemicals, dyestuffs, and pharmaceuticals.¹¹⁻¹⁵ In particular, the use of quinoline derivatives as ligands for transition metal complexes for organic synthesis and organic light-emitting materials has attracted great attention for their synthesis (Fig. 1).¹⁶⁻²¹

Regardless of many classical methods, such as Pfitzinger, Skraup, Friedlander, Doebner von Miller, Conrad-Limbach, and Combes reactions known for the synthesis of quinolines, most are limited by harsh reaction conditions, limited substrate scope and low yields.²² Among the various synthetic strategies,²³⁻²⁴ Lewis acid promoted tandem cyclization reaction of *N*-aryl imines with terminal alkynes under oxidative condition allows the access to a variety of 2,4-disubstituted quinolines.²⁵⁻²⁸ The proposed reaction mechanism involves the nucleophilic addition of a terminal alkyne to imine to form propargylamine intermediate, which undergoes intramolecular cyclization, followed by oxidation to afford quinoline products. Similar cyclization reactions using internal alkynes are difficult and hardly achieved.²⁹ These reactions are not suitable for the synthesis of quinolines without substitution at C-2 position (Scheme 1a).

Natural products



Pharmacologically active compounds



Materials





Fig. 1 Examples of useful quinoline cored compounds.

Recently, the in situ generation of *N*-aryliminium ions from benzylic azides by means of strong acids has been utilized for the synthesis of tetrahydroquinoline scaffolds, which can be oxidized to quinolines by 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ).³⁰⁻³¹ The drawbacks of these reactions are

Department of Chemistry, National Tsing Hua University, Hsinchu 30013, Taiwan. E-mail: chcheng@mx.nthu.edu.tw; Fax: +886-3-5724698; Tel: +886-3-5721454

Electronic Supplementary Information (ESI) available: General experimental procedures, characterization details and ¹H and ¹³C NMR spectra of new compounds. CCDC 1419576–1419579. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/x0xx00000x

(i) lower product yields, (ii) use of strong acid, (iii) poor regioselectivity and (iv) a two-step process (cyclization and oxidation) (Scheme 1b). Because of the useful applications of quinolines and the lack of simple and high yielding method for their synthesis, the search for new synthetic methodologies are still highly sought after. Our continuing interests in the quinoline synthesis^{32–33} lead us to envision that the *in situ* generation of *N*-aryliminium ion from benzylic azide using a Lewis acid catalyst which is also an oxidant under acid free condition provides a good opportunity to avoid the above mentioned problems at least in part. In this report, we disclose a simple and a novel method for the synthesis of various substituted quinolines from benzylic azides and internal alkynes under neutral reaction conditions using Cu(OTf)₂ as both a Lewis acid and oxidant (Scheme 1c).



Scheme 1 Synthetic strategies for substituted quinolines.

Results and discussion


Initially, several Cu^{II}-salts were screened for the reaction of benzyl azide (**1a**) and diphenylacetylene (**2a**) to form 3,4-diphenylquinoline (**3aa**) (Table 1). Treatment of **1a** (0.34 mmol), and **2a** (0.28 mmol) in the presence of Cu(OTf)₂ (0.56 mmol) in MeNO₂ at 100 °C for 24 h gave **3aa** in 36% yield (Table 1, entry 2). The product was unambiguously confirmed by its ¹H and ¹³C NMR, HRMS, and X-ray structure analysis.³⁴ Among the Cu(II)-salts and other Lewis acids tested, only Cu(BF₄)₂·6H₂O showed considerable potency to give **3aa** in 33% yield (See, Table SI for the detailed optimization studies). By reversing the ratio of **1a**

and **2a** from 1.2:1 to 1:1.2, **3aa** was obtained in 63% yield (entry 7). The yield of **3aa** further increased to 77% as the amount of **2a** was raised to 2 equiv. (entry 8). We then optimized the amount of Cu(OTf)₂, reaction time and temperature. The optimized reaction conditions with **1a** (0.28 mmol), **2a** (0.56 mmol), and Cu(OTf)₂ (0.70 mmol) in MeNO₂ (2 mL) at 80 °C for 24 h provided 3,4-diphenylquinoline (**3aa**) in 93% isolated yield. The choice of solvent is crucial to obtain a high product yield. Among the several solvents tested, MeNO₂ afforded the highest product yield, while PhNO₂ and CF₃CH₂OH also gave **3aa** in 60 and 75% yields, respectively (Table 1). The attempt to use Cu(OTf)₂ as catalyst by employing 0.5 equivalent of Cu(OTf)₂ with a stoichiometric amount of K₂S₂O₈ or (NH₄)₂S₂O₈ as oxidants gave the quinoline product in about 50% yield (See Table 1).


Table 1 Optimization studies for the synthesis of 3,4-diphenylquinoline from benzyl azide and diphenylacetylene

Entry	Lewis acid (equiv)	Solvent	Temp.(°C)/ Time (h)	Yield (%) ^a
1	Cu(OAc) ₂ (2)	MeNO ₂	100/24	--
2	Cu(OTf) ₂ (2)	MeNO ₂	100/24	36
3	Cu(OCOCF ₃) ₂ ·H ₂ O (2)	MeNO ₂	100/24	trace
4	Cu(BF ₄) ₂ ·6H ₂ O (2)	MeNO ₂	100/24	33
5	CuCl ₂ (2)	MeNO ₂	100/24	--
6	CuF ₂ (2)	MeNO ₂	100/24	--
7	Cu(OTf) ₂ (2)	MeNO ₂	100/24	63 ^b
8	Cu(OTf) ₂ (2)	MeNO ₂	100/24	77 ^c
9	Cu(OTf) ₂ (3)	MeNO ₂	100/24	92 ^c
10	Cu(OTf) ₂ (4)	MeNO ₂	100/24	87 ^c
11	Cu(OTf) ₂ (3)	MeNO ₂	80/24	89 ^c
12	Cu(OTf) ₂ (3)	MeNO ₂	60/24	80 ^c
13	Cu(OTf)₂ (2.5)	MeNO₂	80/24	93^c
14	Cu(OTf) ₂ (2.5)	MeNO ₂	80/15	72 ^c
15	Cu(OTf) ₂ (1), 1 atm O ₂	MeNO ₂	80/24	45 ^c
15	Cu(OTf) ₂ (3)	EtOAc	80/24	36 ^c
16	Cu(OTf) ₂ (3)	DCE	80/24	22 ^c
17	Cu(OTf) ₂ (3)	PhNO ₂	80/24	60 ^c
18	Cu(OTf) ₂ (3)	TFE	80/24	75 ^c
19	Cu(OTf) ₂ (0.5), K ₂ S ₂ O ₈	MeNO ₂	100/24	49 ^{c,d}
20	Cu(OTf) ₂ (0.5), (NH ₄) ₂ S ₂ O ₈	MeNO ₂	100/24	52 ^{c,d}

^aUnless otherwise mentioned, all reactions were performed with **1a** (0.34 mmol), **2a** (0.28 mmol, limiting reagent), and Lewis acid (as given in the table) in MeNO₂ (2 mL) at 100 °C for 24 h. Isolated yields based on the limiting reagents were given. Yield given in the parenthesis was isolated yield. DCE: 1,2-dichloroethane; TFE: 2,2,2-Trifluoroethanol. ^b**1a** (0.28 mmol, limiting reagent), and **2a** (0.34 mmol) were used. ^cReactions were performed using **1a** (0.28 mmol, limiting reagent), **2a** (0.56 mmol), and Lewis acid (as given in the table) in MeNO₂ (2 mL) at the given temperature and time. ^dCu(OTf)₂ (0.14 mmol) was used along with K₂S₂O₈ or (NH₄)₂S₂O₈ (0.32 mmol).



- | | | | |
|---|---|--|--|
| 1a ; R ¹ = H | 1h ; R ¹ = 4-I | 1o ; R ¹ = 2-Ph | 1v ; 1-(azidomethyl)
naphthalene
1w ; 2-(azidomethyl)
naphthalene |
| 1b ; R ¹ = 4-Me | 1i ; R ¹ = 4-Ph | 1p ; R ¹ = 2-Br | |
| 1c ; R ¹ = 4- <i>i</i> Pr | 1j ; R ¹ = 4-CF ₃ | 1q ; R ¹ = 2-I | |
| 1d ; R ¹ = 4-OMe | 1k ; R ¹ = 4-Ac | 1r ; R ¹ = 2,4-di-Me | |
| 1e ; R ¹ = 4-F | 1l ; R ¹ = 4-CO ₂ Me | 1s ; R ¹ = 2,4-di-F | |
| 1f ; R ¹ = 4-Cl | 1m ; R ¹ = 2-Me | 1t ; R ¹ = 3,5-di-Me | |
| 1g ; R ¹ = 4-Br | 1n ; R ¹ = 2-OMe | 1u ; R ¹ = 3-Me | |




Scheme 2 Scope of benzylic azides in the synthesis of substituted quinolines. Conditions: **1** (0.28 mmol), **2a** (0.56 mmol), and Cu(OTf)₂ (0.70 mmol) in MeNO₂ (2 mL) at 80 °C for 24 h. Isolated yields were determined based on **1**.

ARTICLE

Journal Name

Next, we applied the optimized reaction conditions to a variety of benzylic azides to probe the generality of the reaction (Scheme 2). First, we tested the *para* substituted benzylic azides **1b-l**. Both electron-donating-group (EDG) and electron-withdrawing-group (EWG) substituted substrates were effectively transformed into the respective quinoline products. Thus, the EDG, 4-Me and 4-*i*Pr substituted substrates offered products **3ba** and **3ca** in 92% and 90% yields, respectively, but the very electron-donating 4-methoxy substituted benzyl azide (**1d**) yielded only 36% of product **3da**. Halo-substituted benzylazides **1e-h** are compatible under the reaction conditions to give the expected products **3ea-ha** in good yields. EWGs containing benzylic azides **1i-l** provided the respective substituted quinolines **3ia-la** in 44-90% yields. The steric hindrance of *ortho* substitution at benzylic azides does not show consistent influence on the yield of this cyclization reaction. We studied the reactions of benzylic azides possessing different *ortho* substituents such as Me, OMe, Ph, Br, and I with alkyne **2a**. The results reveal that most of the reactions afforded the respective quinolines in high yields (Scheme 2, products **3ma-qa**). Tetra substituted quinolines **3ra-ta** were obtained in good yields by employing disubstituted benzylic azides **1r-t**. The reaction of 3-methylbenzylic azide gave an inseparable mixture of regioisomeric quinoline products **3ua + 3ua'** (6:4) in 72% yield. Synthesis of substituted benzo[*h*]quinoline (**3va**) and benzo[*f*]quinoline (**3wa**) were achieved from **1v** and **1w** in 74 and 84% yields, respectively. The structure of **3va** was further confirmed by X-ray analysis.³⁴ It is worth to mention that the halo substituted quinoline products **3ea-ha**, **3pa-qa** are useful for further functionalization via cross couplings.


This copper(II) promoted quinoline synthesis was further expanded to a range of symmetrical and unsymmetrical internal alkynes (Scheme 3). Thus, the reaction of *p*-ditolylacetylene (**2b**) with **1a** gave the expected product **3ab** in 87% yield. Similarly, *p*-F, *p*-Cl, *p*-Br substituted diarylacetylenes **2c-e** reacted with **1a** to provide the corresponding quinolines **3ac-ae** in moderate yields. Di(2-thienyl)acetylene also underwent the cyclization reaction to provide **3af** in 33% yield. Next, we tested a number of unsymmetrical alkynes **2g-l** with **1a** and found that the reactions are highly regioselective giving exclusively a single regioisomeric product for each of these reactions. Under the reaction conditions, 1-phenylpropyne (**2g**) and 1-butylnylbenzene (**2h**) provided the respective quinolines **3ag** and **3ah** in 87 and 96% yields, respectively. The regioselectivity of the products were confirmed by X-ray structure analysis.³⁴ Electron-deficient alkyne such as ethyl phenylpropiolate (**2i**) is also active affording the expected quinoline product **3ai** in 62% yield. Moreover, 3-tolylpropargyl alcohol (**2j**), benzyl phenyl acetylene (**2k**) and 3-tolylpropargyl amine derivative (**2l**) also operative to produce quinoline derivatives **3aj - 3al** in good to excellent yields.



Scheme 3 Scope of alkynes in the synthesis of substituted quinolines. Conditions: **1a** (0.28 mmol), **2** (0.56 mmol), and Cu(OTf)₂ (0.70 mmol) in MeNO₂ (2 mL) at 80 °C for 24 h. ^aCu(BF₄)₂·6H₂O (0.70 mmol) was used instead of Cu(OTf)₂.


To demonstrate the synthetic utility of the present copper promoted cyclization reaction, we synthesize biologically active compound 6-chloro-2,3-dimethyl-4-phenylquinoline (**4**) by the

present method. The quinoline compound is known to show effectiveness against *Leishmania donovani*, *Trypanosoma cruzi*, *T. b. rhodesiense*.³⁵ As shown in Scheme 4, compound **4** was conveniently synthesized from **1f** and **2g** in two steps with 62% overall yield.




Scheme 4 Synthesis of biologically active 6-chloro-2,3-dimethyl-4-phenylquinoline **4**.

We also demonstrated the synthesis of 3,4-diphenylquinolin-2(1*H*)-one (**5**) from 3,4-diphenylquinoline (**3aa**) (Scheme 5). Compound **5** is known to be a potential p38 α MAP kinase inhibitor.^{36,37}



Scheme 5 Synthesis of 3,4-diphenylquinolin-2(1*H*)-one **5**.

A plausible reaction mechanism for the copper promoted intermolecular cyclization of benzylic azides and internal alkynes is shown in Scheme 6. The reaction is initiated by Cu^{II} assisted rearrangement of benzylic azide into *N*-aryliminium ion **II** by the loss of N₂ and the migration of the phenyl group.^{38,39} Next, the intermolecular nucleophilic attack of alkyne to **II** forms a vinyl cation intermediate **III**. An intramolecular electrophilic aromatic substitution of intermediate **III** followed by oxidation afford the final quinoline product. It is worth to mention that the reaction offer high regioselectivity for unsymmetrical alkynes **2h-i** (Scheme 2) and good yields for electron withdrawing group substituted benzylic azides (Scheme 1). Presumably, the high regioselectivity is due to the better stabilization of the vinyl cation intermediate **III** by the phenyl ring than the alkyl or ester group on the alkyne substrate. During this copper promoted cyclization, substituted benzylic azides are turned into an electron-rich substituted aryl amine (see intermediate **III**). As a result, an electron withdrawing substituent on the phenylamide ring will not completely stop the electrophilic cyclization of **III**.



Scheme 6 Plausible reaction mechanism.

Conclusions

The copper(II)-promoted synthesis of 3,4-disubstituted quinolines by the intermolecular cyclization reaction of benzylic azides and internal alkynes have been demonstrated. It has been shown that the effectiveness of Cu^{II}-salt in the rearrangement of benzylic azide into *N*-arylimine (Schmidt reaction) which has previously only been achieved by strong acids. The catalytic reaction is highly efficient with a wide range of substituted benzylic azides. Excellent regioselectivity was observed for the reaction with unsymmetrical alkyne giving a single regioisomeric product in high yield. The present reaction system is further applied to the synthesis of two biologically active compounds.

Experimental section

General procedure for the synthesis of benzylic azides⁴⁰

Substituted benzylic bromide (1.0 equiv) and sodium azide (1.5 equiv) were dissolved in DMF (2.0 mL/mmol) and stirred at 30 °C for 12 h. At the end of the reaction, the mixture was diluted with water and extracted with diethyl ether. The combined solution was concentrated in vacuo and the mixture was purified by a silica gel column (hexane/EtOAc, 90:10) to afford benzylic azide.

Typical procedure for the synthesis of substituted quinoline **3aa**

A sealed tube containing Cu(OTf)₂ (254 mg, 0.70 mmol) and 3,4-diphenylacetylene **2a** (100 mg, 0.56 mmol) was evacuated and purged with nitrogen gas three times. Then, a solution of benzylic azide **1a** (38 mg, 0.28 mmol) in MeNO₂ (2.0 mL) was added to the system by syringe under a nitrogen atmosphere and the reaction was stirred at 80 °C for 24 h. At the end of the reaction, the mixture was diluted with CH₂Cl₂ (10 mL), filtered through a Celite pad, which was then washed three times with CH₂Cl₂ (2

X 20 mL). The combined filtrate was concentrated in vacuo and the mixture was purified by a silica gel column using hexane/EtOAc (95:5) as eluent to afford the desired pure product **3aa** in 91% (71 mg) yield.

Yellow solid: m.p. 135-137 °C; ¹H NMR (400 MHz, CDCl₃): δ 8.99 (s, 1 H), 8.18 (d, *J* = 8.4 Hz, 1 H), 7.74-7.64 (m, 2 H), 7.49-7.43 (m, 1 H), 7.37-7.30 (m, 3 H), 7.23-7.14 (m, 7 H); ¹³C NMR (100 MHz, CDCl₃): δ 151.8 (CH), 147.5 (C), 145.5 (C), 138.1 (C), 136.3 (C), 133.1 (C), 130.5 (2 CH), 130.1 (2 CH), 129.5 (CH), 129.1 (CH), 128.1 (2 CH), 128.0 (2 CH), 127.7 (CH), 127.2 (C), 127.0 (CH), 126.8 (CH), 126.6 (CH); HRMS (FAB) cal for C₂₁H₁₅N [M⁺] 281.1204, found 281.1204; IR (KBr): 2923, 2854, 1727, 1565, 1488, 1442, 1380, 1272, 1072, 1025, 763 and 701 cm⁻¹.

Procedure for the synthesis of biologically active compound **4**⁴¹

Compound **3fg** was synthesized in 75% yield from 1-(azidomethyl)-4-chlorobenzene (**1f**) and phenylpropyne (**2g**) using a procedure similar to that for the synthesis of quinoline **3aa**.

Compound **3fg** (110 mg, 0.43 mmol) was dissolved in THF (4.0 mL) and MeLi/LiBr (0.40 mL (2.2 M in Et₂O, 0.87 mmol) was added to the solution at -78 °C. The mixture was allowed to warm to room temperature for 24 h. At the end of the reaction, iodine (328 mg, 1.29 mmol) was added to the mixture at 0 °C and stirred for 1 h at the same temperature. The mixture was then quenched with a saturated sodium thiosulfate solution (10 mL). The resulted biphasic solution was extracted with EtOAc (3 X 30 mL). The combined organic solution was concentrated in vacuo and the mixture was purified by a silica gel column using hexane/EtOAc (95:5) as eluent to afford the desired pure product **4** in 82% (94 mg) yield.

Yellow solid: m.p. 125-127 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.93 (d, *J* = 8.8 Hz, 1 H), 7.54-7.44 (m, 4 H), 7.25 (d, *J* = 2.4 Hz, 1 H), 7.21-7.18 (m, 2 H), 2.72 (s, 3 H), 2.15 (s, 3 H); ¹³C NMR (100 MHz, CDCl₃): δ 159.3 (C), 145.5 (C), 144.4 (C), 136.9 (C), 131.2 (C), 130.1 (CH), 129.3 (2 CH), 128.9 (CH), 128.7 (2 CH), 128.5 (C), 128.0 (CH), 127.6 (C), 124.8 (CH), 24.5 (CH₃), 17.0 (CH₃); HRMS (ESI) [M+H]⁺ cal for C₁₇H₁₅ClN 268.0893, found 268.0885; IR (KBr): 3062, 2923, 2854, 1727, 1666, 1589, 1481, 1373, 1172, 1072, 948, 825 and 701 cm⁻¹.

Procedure for the synthesis of 3,4-diphenylquinolin-2(1H)-one **5**.³⁶

Compound **3aa** (100 mg, 0.36 mmol) was dissolved in CH₂Cl₂ (4.0 mL) and *meta*-chloroperoxybenzoic acid (134 mg, 0.54 mmol) was added to the solution at 0 °C and stirred for 4 h at the same temperature. After that the reaction was quenched with saturated sodium bicarbonate solution (5 mL). The resulted biphasic solution was extracted with CH₂Cl₂ (3 X 10 mL) and the combined organic layer was dried over MgSO₄ and concentrated by rotary evaporation. The crude product was dissolved in Ac₂O (3.0 mL) and heated at 75 °C for 18 h. At the end of the reaction, the mixture was quenched with saturated sodium bicarbonate solution and extracted with CH₂Cl₂ (3 X 10 mL). The combined solution was dried over MgSO₄ and concentrated in vacuo and the resulted residue was purified by a

silica gel column using hexane/EtOAc (95:5) as eluent to afford the desired pure product **5** in 52% (55 mg) yield. Yellow solid: m.p. 173-175 °C; ¹H NMR (400 MHz, *d*₆-DMSO): δ 12.0 (bs, 1 H), 7.49 (t, *J* = 7.6 Hz, 1 H), 7.39 (d, *J* = 8.0 Hz, 1 H), 7.32-7.22 (m, 3 H), 7.18-7.04 (m, 8 H), 6.99 (d, *J* = 8.0 Hz, 1 H); ¹³C NMR (100 MHz, *d*₆-DMSO): δ 161.2 (C), 148.1 (C), 138.2 (C), 136.1 (C), 135.7 (C), 131.9 (C), 130.6 (2 CH), 130.1 (CH), 129.5 (2 CH), 127.9 (2 CH), 127.5 (CH), 127.1 (2 CH), 126.8 (CH), 126.5 (CH), 121.7 (CH), 119.9 (C), 115.1 (CH); HRMS (ESI) [M+H]⁺ cal for C₂₁H₁₆ON 298.1232, found 298.1225; IR (KBr): 3162, 1727, 1643, 1596, 1481, 1442, 1288, 705 and 701 cm⁻¹.

Acknowledgements

We thank the Ministry of Science and Technology of the Republic of China (MOST-103-2633-M-007-001) for support of this research.

Notes and references

- 1 J. P. Michael, *Nat. Prod. Rep.*, 2005, **22**, 627-646.
- 2 J. P. Michael, *Nat. Prod. Rep.*, 2007, **24**, 223-246.
- 3 J. P. Michael, *Nat. Prod. Rep.*, 2008, **25**, 166-187.
- 4 V. R. Solomon and H. Lee, *Curr. Med. Chem.*, 2011, **18**, 1488-1508.
- 5 G. Roma, M. D. Braccio, G. Grossi, F. Mattioli and M. Ghia, *Eur. J. Med. Chem.*, 2000, **35**, 1021-1035.
- 6 Y.-L. Chen, K.-C. Fang, J.-Y. Sheu, S.-L. Hsu and C.-C. Tzeng, *J. Med. Chem.*, 2001, **44**, 2374-2377.
- 7 A. P. Gorka, A. d. Dios and P. D. Roepe, *J. Med. Chem.*, 2013, **56**, 5231-5246.
- 8 P. Zajdel, A. Partyka, K. Marciniak, A. J. Bojarski, M. Pawlowski and A. Wesolowska, *Future Med. Chem.*, 2014, **6**, 57-75.
- 9 S. Mukherjee and M. Pal, *Curr. Med. Chem.*, 2013, **20**, 4386-4410.
- 10 R. Musiol, *Curr. Pharm. Des.*, 2013, **19**, 1835-1849.
- 11 F. M. Hamer, *The Cyanine Dyes and Related Compounds*, Interscience Publishers, New York, London; Wormerveer printed, 1964.
- 12 K.-Y. Lu, H.-H. Chou, C.-H. Hsieh, Y.-H. O. Yang, H.-R. Tsai, H.-Y. Tsai, L.-C. Hsu, C.-Y. Chen, I. C. Chen and C.-H. Cheng, *Adv. Mater.*, 2011, **23**, 4933-4937.
- 13 A. R. Surrey and H. F. Hammer, *J. Am. Chem. Soc.*, 1946, **68**, 113-116.
- 14 J. Wiesner, R. Ortmann, H. Jomaa and M. Schlitzer, *Angew. Chem. Int. Ed.*, 2003, **42**, 5274-5293.
- 15 M. Takamura, K. Funabashi, M. Kanai and M. Shibasaki, *J. Am. Chem. Soc.*, 2001, **123**, 6801-6808.
- 16 S. Ernst and W. Kaim, *J. Am. Chem. Soc.*, 1986, **108**, 3578-3586.
- 17 P. J. Steel, *Coord. Chem. Rev.*, 1990, **106**, 227-265.
- 18 A. Mamo, S. Nicoletti and N. C. Tat, *Molecules*, 2002, **7**, 618-627.
- 19 Y.-Z. Hu, G. Zhang and R. P. Thummel, *Org. Lett.*, 2003, **5**, 2251-2253.
- 20 Y. Fukata, K. Asano and S. Matsubara, *Chem. Lett.*, 2013, **42**, 355-357.
- 21 P. W. Yawalkar, S. J. Dhoble, N. Thejo Kalyani, R. G. Atram and N. S. Kokode, *Luminescence*, 2013, **28**, 63-68.
- 22 A. Weissberg and E. C. Taylor, *The Chemistry of Heterocyclic Compounds, Quinoline, Parts I-III*; vol. 32, ed. G. Jones, Interscience Publications, Wiley, London, 1977.

- 23 J. B. Bharate, R. A. Vishwakarma and S. B. Bharate, *RSC Adv.*, 2015, **5**, 42020-42053.
- 24 V. Kouznetsov, L. Mendez and C. Gomez, *Curr. Org. Chem.*, 2005, **9**, 141-161.
- 25 Y. Zhang, P. Li and L. Wang, *J. Heterocycl. Chem.*, 2011, **48**, 153-157.
- 26 F. Xiao, Y. Chen, Y. Liu and J. Wang, *Tetrahedron*, 2008, **64**, 2755-2761.
- 27 K. Cao, F.-M. Zhang, Y.-Q. Tu, X.-T. Zhuo and C.-A. Fan, *Chem. Eur. J.*, 2009, **15**, 6332-6334.
- 28 A. Kumar and V. Rao, *Synlett*, 2011, **2011**, 2157-2162.
- 29 X. Zhang, B. Liu, X. Shu, Y. Gao, H. Lv and J. Zhu, *J. Org. Chem.*, 2012, **77**, 501-510.
- 30 J. Tummatorn, C. Thongsornkleeb and S. Ruchirawat, *Tetrahedron*, 2012, **68**, 4732-4739.
- 31 J. Tummatorn, P. Poonsilp, P. Nimmual, J. Janprasit, C. Thongsornkleeb and S. Ruchirawat, *J. Org. Chem.*, 2015, **80**, 4516-4525.
- 32 R. P. Korivi and C.-H. Cheng, *J. Org. Chem.*, 2006, **71**, 7079-7082.
- 33 P. Gandeepan, P. Rajamalli and C.-H. Cheng, *Asian J. Org. Chem.*, 2014, **3**, 303-308.
- 34 CCDC 1419576-1419579 contains the supplementary crystallographic data for this paper.
- 35 G. C. Muscia, M. Bollini, J. P. Carnevale, A. M. Bruno and S. E. Asís, *Tetrahedron Lett.*, 2006, **47**, 8811-8815.
- 36 J. DeRuiter, B. E. Swearingen, V. Wandrekar and C. A. Mayfield, *J. Med. Chem.*, 1989, **32**, 1033-1038.
- 37 C. Peifer, R. Urich, V. Schattel, M. Abadleh, M. Röttig, O. Kohlbacher and S. Laufer, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 1431-1435.
- 38 A. Murali, M. Puppala, B. Varghese and S. Baskaran, *Eur. J. Org. Chem.*, 2011, **2011**, 5297-5302.
- 39 P. Desai, K. Schildknegt, K. A. Agrios, C. Mossman, G. L. Milligan and J. Aubé, *J. Am. Chem. Soc.*, 2000, **122**, 7226-7232.
- 40 O. Berger, A. Kaniti, C. T. van Ba, H. Vial, S. A. Ward, G. A. Biagini, P. G. Bray and P. M. O'Neill, *ChemMedChem*, 2011, **6**, 2094-2108.
- 41 C. C. C. Johansson, N. Bremeyer, S. V. Ley, D. R. Owen, S. C. Smith and M. J. Gaunt, *Angew. Chem. Int. Ed.*, 2006, **45**, 6024-6028.


View Article Online
DOI: 10.1039/C5RA23065A

RSC Advances Accepted Manuscript

Graphical Abstract:

Copper Promoted Synthesis of Substituted Quinolines from Benzylic Azides and Alkynes

Ching-Zong Luo, Parthasarathy Gandeepan, Yun-Ching Wu, Wei-Chen Chen and Chien-Hong Cheng*



A novel method for the synthesis of substituted quinolines from benzylic azides and internal alkynes using $\text{Cu}(\text{OTf})_2$ is described. The reaction features a broad substrate scope, high product yields and excellent regioselectivity.