

Dalton Transactions

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: N. Nakagawa, E. J. Derrah, M. Schelwies, F. -. Rominger, O. Trapp and T. Schaub, *Dalton Trans.*, 2016, DOI: 10.1039/C5DT04870B.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Triphos derivatives and diphosphines as ligands in the ruthenium-catalysed alcohol amination with NH₃

N. Nakagawa,^a E. J. Derrah,^a M. Schelwies,^b F. Rominger,^c O. Trapp^c and T. Schaub^{*,a,b}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


www.rsc.org/

The ruthenium-triphos and diphosphine-catalysed amination of alcohols with ammonia is reported. Various types of triphos derivatives with electron-donating functional group were synthesized and used as ligands in the Ru-catalysed alcohol amination with NH₃. The triphos derivatives are effective for the formation of primary amines. On the other hand, if hemilabile diphosphines as tridentate ligands are used, mixtures of secondary- along with primary amines are obtained. It was found that even simple diphosphines can be used as ligands for the selective formation of the secondary amines. The diphosphine system allows a new entry to the Ru-catalysed formation of secondary amines.

Introduction

Amines are important building blocks in the manufacture of pharmaceuticals and agrochemicals.¹ Intensive research has focused on the selective formation of primary, secondary and tertiary amines.² One of the most straightforward and environmentally-friendly methods is the amination of alcohols using cheap and abundant ammonia (NH₃), which represents an atom efficient route towards this valuable class of organic compounds, along with water as the sole by-product.³ The amination of alcohols with ammonia is performed with heterogeneous catalysts on industrial scale. Using heterogeneous catalysts, it is in many cases difficult to control the selectivity as harsh reaction conditions are required.^{1c} During the last decade, the amination of alcohols has been further developed with a series of homogeneous catalysts.⁴ In particular, ruthenium catalysts were identified to exhibit a high performance for the selective formation of primary amines⁵ and secondary amines.⁶ The combination of a ruthenium pre-catalyst and the tridentate phosphine ligand 1,1,1-tris(diphenylphosphinomethyl)ethane **1** (= triphos) was first published as one of the candidates in the alcohol amination by Beller and co-workers^{7a} and investigated in detail by our laboratory (Scheme 1).^{7b} We disclosed that the catalytic active species is a cationic ruthenium-triphos complex through experimental and computational investigations. Our previous report on this ruthenium catalysis described only the use of

the commercial available triphos **1** and effects of other phosphine derivatives have still remained undisclosed and undeveloped. We herein report the synthesis and evaluation of triphos derivatives and the corresponding ruthenium complexes in order to gain an insight how variations of the triphos scaffold influence the ruthenium-catalysed amination of alcohols with NH₃. Our approach was to alter the coordination sphere of triphos-type Ru-complexes in order to change the selectivities in the corresponding amination reactions.



Scheme 1 Ruthenium-triphos-catalysed amination of 1-octanol with ammonia^{7b}

Results and discussion

Ligands

To evaluate the ligand influence on the catalytic performance of the ruthenium-catalysed amination of 1-octanol with NH₃, the following triphos derivatives were used: Triphos **1**, the newly synthesised phenyl-*p*-tolyl-mixed triphos derivative **2**, and the all-*p*-tolyl triphos derivative **3** (obtained via a modified procedure).⁸ We also examined diphosphines with an electron-donating functional group such as a pyridyl substituted derivative **4**, which is potentially hemilabile ligand,⁹ as well as oxygen-substituted ones **5**¹⁰ and **6**.

^a Catalysis Research Laboratory (CaRLa), Im Neuenheimer Feld 584, D-69120 Heidelberg, Germany.

^b Synthesis & Homogeneous Catalysis, BASF SE, Carl-Bosch-Straße 38, D-67056 Ludwigshafen, Germany.

^c Organisch-Chemisches Institut, Ruprecht-Karls-Universität Heidelberg, Im Neuenheimer Feld 270, D-69120 Heidelberg, Germany.

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information should be included here]. See DOI: 10.1039/x0xx00000x

ARTICLE

Journal Name





Fig. 1 Triphos **1**, triphos derivatives **2** and **3**, and diphosphines with electron-donating functional groups **4–6** used in this work.


Synthesis of triphos derivatives via modified protocol

C_{3v} symmetrical triphos-type ligands such as **3** can be obtained using 1,1,1-tris(chloromethyl)ethane as a starting material. A route to C_1 symmetric variants of the ligand with three different phosphine donors was reported by Huttner and Helmchen.⁸ However, since we were only interested altering one of the three donor groups of triphos, we chose **7** as key intermediate^{10a} in our synthesis (Scheme 2). The substitution reaction of $CH_2C(CH_3)(CH_2OMs)_2(CH_2Br)$ **7** with an excess amount of lithium diphenylphosphide, followed by borane protection afforded a monomesyl diphosphine-borane compound **6-PG** in 64 % yield (Scheme 2a). After deprotection with DABCO, the intermediate **6** was obtained in 66 % yield. We repeated this sequence using lithium di(*p*-tolyl)phosphide with monomesylate **6** and obtained the unsymmetrical triphos-borane **2-PG**. The structure of **2-PG** was confirmed by X-ray analysis (Fig. 2). Deprotection of **2-PG** afforded the unsymmetrical triphos **2** in 27 % overall yield. The same protocol was applied to the synthesis of all-*p*-tolyl triphos **3** (51 % overall yield, Scheme 2b).¹¹

(a) Synthesis of **2**



(b) Synthesis of **3**



Scheme 2 Optimised synthetic protocol for triphos derivatives





Fig. 2 ORTEP drawing of the borane-adduct **2-PG** where all non-hydrogen atoms are represented by Gaussian ellipsoids at the 50 % probability level. All hydrogen atoms have been omitted for clarity.

Synthesis of ruthenium complexes

Ruthenium complexes of the triphos derivatives **1-3** were prepared according to the reported protocol for Ru-triphos complex **A** (Scheme 3).¹² When using **2** as a ligand, a mixture of three isomers is formed. ³¹P-NMR indicates that the three isomers differ in the position of the tolyl moiety (see supporting information). Elemental analysis confirms the composition of the mixture of **B-1**, **B-2**, and **B-3**. In case of the *all*-tolyl substituted ligand **3**, only one complex (**C**) is formed like in the synthesis of **A**. The structure of **C** was determined by X-ray analysis (Fig. 3). The distance of Ru-P bonds in **C** is the range of 2.27–2.38 Å which is similar like in the reported structure for RuHCl(CO)(triphos) **A**.¹³



Scheme 3 Synthesis of triphos-type ruthenium complexes **A-C**






Fig. 3 ORTEP drawing of the ruthenium complex **C** where all non-hydrogen atoms are represented by Gaussian ellipsoids at the 50 % probability level and all hydrogen atoms have been removed for simplicity. Selected bond lengths (Å) and angles (°) of **C**: Ru–P1 = 2.2740(12); Ru–P2 = 2.3847(12); Ru–P3 = 2.3840(13); Ru–H = 1.88(3); Ru–Cl = 2.5000(13); P1–Ru–P2 = 88.31(4); P2–Ru–P3 = 86.94(4); P3–Ru–P1 = 88.67(4).

We also conducted the preparation of the corresponding ruthenium complexes with pyridyl substituted diphosphine **4**.⁹ Using this ligand, a mixture of three complexes was obtained: the two isomers **D-1** and **D-2** as well as the dichloro ruthenium species **D-3** (Scheme 4). The structure of **D-3** was confirmed by X-ray analysis (Fig 4). Regarding the structure of **D-3**, the carbonyl ligand is located *trans* to the pyridyl moiety. Although the reaction pathway towards **D-3** is not confirmed, we propose that the chlorine atom on **D-3** was derived from the CH₂Cl₂ solvent through the chlorine abstraction by ruthenium hydride species. Under these conditions, selective formation and complete separation of the three isomers turned out to be difficult. In addition, we failed to obtain the ruthenium-pyridyl diphosphine complex selectively in toluene as solvent. Therefore, we used the pyridyl diphosphine **4** directly in combination with [Ru(PPh₃)₃(CO)HCl] in the amination reaction to generate the desired complexes *in situ* (*vide infra*).





ARTICLE

Journal Name

Scheme 4 Preparation of pyridyl diphosphine ruthenium complexes **D-1**, **D-2**, and **D-3****Fig. 4** ORTEP drawing of the dichloro ruthenium complex **D-3** where all non-hydrogen atoms are represented by Gaussian ellipsoids at the 50 % probability level. All hydrogen atoms and coordinating solvent have been omitted for clarity. Selected bond lengths (Å) and angles (°) of **D-3**: Ru–P1 = 2.2970(14); Ru–P2 = 2.2844(15); Ru–N = 2.178(4); Ru–Cl1 = 2.4314(13); Ru–Cl2 = 2.4775(14); P1–Ru–P2 = 87.65(5); P1–Ru–N = 83.76(11); P2–Ru–N = 90.25(11).

We were also successful to synthesise a novel hydroxyl-containing ruthenium complex **E** from **5**¹⁰ (Scheme 5). The structure of **E** was confirmed by X-ray analysis (Fig 5). One PPh₃ ligand from the precursor remains on ruthenium which is consistent with reported RuHCl(CO)PPh₃(diphosphine) complexes.^{5e,5g} The hydroxyl group orients to the opposite direction from the ruthenium centre, thus it seems to be an unsuitable coordinating group to achieve a tridentate coordination mode. In order to facilitate a coordination of the oxygen atom, the ruthenium complex **E** was treated with potassium *tert*-butoxide (Scheme 5). The red powder obtained was assigned as the alkoxo ruthenium complex **F** by ¹H and ³¹P NMR spectra as well as LIFDI mass spectroscopy. It is conceivable that under a basic ammonia atmosphere of the alcohol amination the alkoxo complex **F** will be formed in-situ from **E**.

**Scheme 5** Synthesis of hydroxyl-containing diphosphine ruthenium complex **E** and reaction between complex **E** and *t*BuOK**Fig. 5** ORTEP drawing of hydroxyl-containing diphosphine ruthenium complex **E** where all non-hydrogen atoms are represented by Gaussian ellipsoids at the 50 % probability level. All hydrogen atoms and coordinating solvent have been omitted for clarity. Selected bond lengths (Å) and angles (°) of **E**: Ru–P1 = 2.4279(10); Ru–P2 = 2.3586(10); Ru–P3 = 2.3806(11); Ru–H = 1.62(5); Ru–Cl = 2.4725(11); P1–Ru–P2 = 94.38(3); P3–Ru–P1 = 100.04(4).**Catalytic investigations**

In order to investigate the ligand effects, the ruthenium-catalysed amination of 1-octanol with NH₃ was conducted for the above described ligands **1-6** (Table 1). As previously reported by our group,⁷ the ruthenium-triphos complex **A** showed a high catalytic performance towards octylamine **8** along with small amounts of the secondary and tertiary amines **9** and **10** (entry 1). Note that it is unnecessary to use the isolated ruthenium catalyst **A**: the *in-situ* prepared ruthenium-triphos catalyst showed the same activity and selectivity as **A** (entry 2). In the absence of any additional phosphine, the amination products **8** to **10** were not obtained (entry 3). When the *p*-tolyl-substituted triphos derivatives **2** and **3** were used instead of **1**, it was found that the selectivity towards the primary amine is slightly shifted to dioctylamine **9** (entries 4 and 5). The use of the pyridyl diphosphine **4**, a potentially hemilabile ligand, led to the formation of dioctylamine **9** (entry 6). The amination reactions using the hydroxyl-diphosphine complex **E** and its alkoxo complex **F** were sluggish and the dioctylamine **9** is formed in good selectivity (entries 7 and 8). These results can be interpreted as the complex **E** probably converts into the alkoxo complex **F** through deprotonation of the hydroxyl group under the basic conditions of the amination. When the mesyl diphosphine **6**, which is the synthetic intermediate of the ligands **2** and **3**, was used as a ligand, dioctylamine **9** is formed in good selectivity (entry 9). Probably the mesyl group (OMs) will be displaced by the amino group (NH₂) via nucleophilic substitution under the ammonia atmosphere of the amination reaction. To our surprise, even simpler diphosphines, such as dpppdmp (1,3-

bis(diphenylphosphino)-2,2-dimethylpropane, $\text{Ph}_2\text{PCH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{PPh}_2$ and dppp (1,3-bis(diphenylphosphino)propane, $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{PPh}_2$), were found to be good ligands for the preferable formation of dioctylamine **9** (entries 10 and 11).^{14,15} Finally, the highly selective formation of dioctylamine **9** was achieved with dppp at a higher substrate concentration (entries 12 and 13).

Table 1 Effect of altering the phosphine ligand

$\text{C}_8\text{H}_{17}\text{OH} + \text{NH}_3$ (7-8 bar)		Ru-phosphine catalyst (0.20 mol%) toluene 165 °C, 15 h		$\text{C}_8\text{H}_{17}\text{NH}_2$ (8) $(\text{C}_8\text{H}_{17})_2\text{NH}$ (9) $(\text{C}_8\text{H}_{17})_3\text{N}$ (10)		
Entry ^a	Ru-phosphine catalyst	GC area ratio (%)				Selectivity of mono:di (%)
		mono	di	tri	SM recov.	
1	A	96	3	<1	1	97:3
2 ^b	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3 + \mathbf{1}$	93	5	<1	2	95:5
3	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3$	<1	<1	<1	99	-
4	B	90	8	<1	1	92:8
5	C	86	11	1	2	88:12
6	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3 + \mathbf{4}$	14	66	3	17	18:82
7	E	10	47	2	41	17:83
8	F	9	51	2	38	15:85
9	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3 + \mathbf{6}$	10	88	<1	1	10:90
10	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3 + \text{dppdpmp}^c$	11	86	2	1	11:89
11	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3 + \text{dppp}^d$	35	63	<1	1	36:64
12 ^e	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3 + \text{dppp}^d$	<1	95	4	1	<1:99
13 ^f	$\text{RuHCl}(\text{CO})(\text{PPh}_3)_3 + \text{dppp}^d$	5	87	6	2	8:92
		[5] ^g	[88] ^g	[5] ^g	[2] ^g	

^a 1-Octanol (3.0 g, 23.6 mmol) and 17 mL of toluene was used. ^b 0.22 mol % of ligand was used. ^c dppdpmp = 1,3-bis(diphenylphosphino)-2,2-dimethylpropane. ^d dppp = 1,3-bis(diphenylphosphino)propane. ^e 6 mL of toluene was used as a solvent. ^f $p(\text{NH}_3)$ = 4 bar (~52 mmol). ^g The calibrated GC yields against hexadecane as an internal standard.

The reduced amount of toluene significantly affected the formation of the secondary amine. In order to get more insights in the reaction, we examined the reaction profile by analysing the reaction mixture at several reaction times (1-8 h).¹⁶ The obtained curves showed that there is no significant induction period, and that the amount of octylamine **8** and dioctylamine **9** increases as time elapses without decomposing or interconverting into the other products. We also confirmed the formation of small amount of octyloctan-1-imine **11**, which is assumed to be the intermediate before dioctylamine **9** is formed. The imine **11** is probably reactive to a Ru-H species to produce dioctylamine **9** through hydrogenation.





Fig. 6. Reaction profile for the Ru-dppp-catalysed amination of octanol with NH_3 . Reaction conditions: In a premix autoclave (60 mL stainless steel), 1-octanol (3.0 g, 24 mmol) (\blacklozenge), $p(\text{NH}_3)$ = 4 bar (0.9 g, 53 mmol of NH_3), $\text{RuHCl}(\text{CO})(\text{PPh}_3)_3$ (45 mg, 0.047 mmol), dppp (21 mg, 0.052 mmol), and toluene (6 mL). Observed products: octylamine **8** (\blacktriangle), dioctylamine **9** (\blacksquare), and octyloctan-1-imine **11** (\bullet). Trioctylamine **10** was observed in 2-4 % GC area ratio in each reaction time.

The possible mechanism is described in Scheme 6. As we postulated in the previous report,⁷ the selective mono-alkylation of ammonia using ruthenium-triphos complexes involves the formation of cationic ruthenium-triphos intermediates as active species (Scheme 6a). Among the triphos derivatives, the different ratio between the primary and secondary amines can originate from the difference of electron negativity of P atom. In the case of hemilabile diphosphines and simple diphosphines, the formation of the secondary amine is preferred under the same reaction conditions. This observation can be explained by considering a different mechanism. As Vogt and co-workers disclosed the mechanism on the amination of cyclohexanol using xantphos,^{4g} the dissociation of the remaining PPh_3 can play an important role (Scheme 6b). Although the mechanism of dialkylation is still unclear, it is possible to consider that dissociation of PPh_3 and chloride can provide more vacant sites on the ruthenium than in the triphos systems. The transient ketone or imine can coordinate at the vacant site and incorporate in the further alkylation. When octylamine **8** as a starting material is used instead of 1-octanol, the formation of the dioctylamine **9** was only observed in small amounts (eq 1). Therefore, it is conceivable that the ruthenium-diphosphine complex possesses a poor ability on dehydrogenation of the primary amine **8**. Presumably, in-situ generated ketone or imine (from the starting alcohol) are the reactants for the dialkylation.


ARTICLE

Journal Name


(a) With triphos-type ligands (based on Ref. 7)



(b) With diphosphine ligands (based on Ref. 4g)



Scheme 6 Simplified possible catalytic cycles



Conclusions

A series of ruthenium-complexes bearing triphos derivatives and diphosphines with the electron-donating functional group were prepared and examined for the amination of primary alcohols with NH_3 . Deviating the P-substituents on the triphos scaffold slightly affects the ratio of mono- and dialkylamines, but these triphos derivatives are basically effective for the formation of the primary amines. The reaction mechanism of mono-alkylation can involve the formation of the cationic ruthenium complex based on our previous investigations. On

the other hand, the use of hemilabile diphosphines leads to a mixture of the primary and secondary amines. It seems to be clearer after these results, that the origin of high selectivities towards the primary amines is related to the very stable ligand spheres of Ru/triphos and the Milstein type Ru/pincer systems.^{5a,h} The selective formation of the secondary amine was finally realized by using simple dppp as the ligand. The reaction mechanism of the ruthenium-diphosphine system can be influenced by dissociation of attaching PPh_3 which is different to the triphos-based systems. So far, the preparation of secondary amines from alcohols with NH_3 using homogeneous catalysts has been achieved by Milstein and co-worker with Ru catalyst,⁶ or by Fujita, Yamaguchi, and co-workers as well as in previous work from our laboratory using Ir catalysts.¹⁷ Compared with these studies, our current diphosphine system offers a simple protocol for the preparation of diamines with commercially available catalysts.

Experimental

General considerations

All reactions were carried out under a positive pressure of argon in an MBraun glovebox or using standard Schlenk line techniques. All nondeuterated solvents were dried using an MBraun SPS-800 solvent purification system and degassed prior to use. 3-methyl-3-oxetanemethanol (a precursor of **7**), and 1-octanol were purchased from Aldrich and distilled prior to use. Di-*p*-tolyl and diphenylphosphines were purchased from ABCR, $\text{RuHCl(CO)(PPh}_3)_3$ was supplied by BASF and used without further purification. All other products were purchased from Aldrich and used without further purification. Liquid reagents including deuterium solvents were distilled prior to use, and all others were used without further purification. ^1H , ^2H , ^{13}C , and ^{31}P NMR spectra were recorded on a Bruker Avance 200, 400, or 600 MHz spectrometer. ^1H and ^{13}C chemical shifts are reported relative to residual solvent signals of CD_2Cl_2 (5.32 and 54.0 ppm) and $\text{THF-}d_8$ (5.38 and 67.21 ppm). $^{31}\text{P}\{^1\text{H}\}$ chemical shifts are referenced to an external 85% solution of phosphoric acid. The ^{13}C NMR data were assigned by HSQC and HMBC spectra. FAB and HR mass spectrometry was measured at the Mass Spectrometry Facility (Institute of the Organic Chemistry, University Heidelberg). Gas chromatography was performed on an Agilent 6890N modular GC base equipped with a split-mode capillary injection system and a flame ionization detector using a BGB-5 capillary column (Agilent 122-1033; 30 m \times 0.32 mm \times 0.25 μm ; He flow 1.0 mL/min, program: initial 50 $^\circ\text{C}$ for 2 min, ramp 6 $^\circ\text{C}/\text{min}$, 300 $^\circ\text{C}$ for 10 min). Starting materials and products had the following retention times: octylamine (t_R = 14.54 min), 1-octanol (t_R = 15.16 min), dioctylamine (t_R = 31.34 min), and trioctylamine (t_R = 41.71 min). Elemental analysis were performed in the "Mikroanalytisches Laboratorium der Chemischen Institute der Universität Heidelberg". X-ray structures were solved by direct methods and refined against F^2 with a full-matrix least squares algorithm by using the SHELXTL (version 2014/7) software package.¹⁸ Intensities were corrected for Lorentz and polarisation effects.¹⁹ CCDC 1440322 (**2-PG**), 1440323 (**C**), 1440324 (**D-3**), and 1440325 (**E**) contain the supplementary crystallographic data for

this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Synthesis of $\text{H}_3\text{CC}(\text{CH}_2\text{OMes})_2(\text{CH}_2\text{Br})$ (7)

A 100 mL round-bottom Schlenk flask was charged with 25 mL dioxane and 3-methyl-3-oxetanemethanol (5.45 g, 53 mmol). Aqueous HBr (48 %, 7.2 mL, 64 mmol) was slowly added over a 5-7 min period to give a light yellow solution that was slightly warm to the touch. The solution was refluxed for 3.5 h and the solvent was removed under vacuum (60 °C) to give $\text{H}_3\text{CC}(\text{CH}_2\text{OH})_2(\text{CH}_2\text{Br})$ (9.38 g, 51 mmol, 96 % yield) as an orange-brown solid. The product was used in the next step without further purification.

^1H NMR (200.1 MHz, CD_2Cl_2 , δ): 3.63 (s, 4H, CH_2OH), 3.54 (s, 2H, CH_2Br), 2.95 (s, 2H, OH), 0.91 (s, 3H, CH_3).

A 500 mL round-bottom Schlenk flask was charged with $\text{H}_3\text{CC}(\text{CH}_2\text{OH})_2(\text{CH}_2\text{Br})$ (4.08 g, 22.3 mmol), triethylamine (4.96 g, 0.49 mol) and 200 mL CH_2Cl_2 . The colourless solution was cooled in an ice bath and neat methanesulfonyl chloride (5.36 g, 47 mmol) was added dropwise to give a colourless solution and white precipitate. The solution was stirred for 2 h at 0 °C. The solution was concentrated to ~100 mL under vacuum and water (100 mL) was added. The organic layer was extracted and washed with another 100 mL of water, dried over MgSO_4 , and the organic solvent was removed under vacuum to obtain $\text{H}_3\text{CC}(\text{CH}_2\text{OMes})_2(\text{CH}_2\text{Br})$ **7** (6.14 g, 18 mmol, 81 % yield) as a thick yellowish oil.

Synthesis of $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2\cdot\text{BH}_3)_2(\text{CH}_2\text{OMes})$ (6-PG)

A 500 mL round-bottom Schlenk flask was charged with HPPH_2 (7.99 g, 43 mmol) and 60 mL THF. The colourless solution was cooled in an ice bath and 2.5 M *n*-BuLi in hexane (19 mL, 47 mmol) was added drop wise via syringe to give a red/orange solution. The solution was stirred for 30 min before being transferred to a 100 mL addition funnel affixed to a 500 mL round-bottom Schlenk flask charged with $\text{H}_3\text{CC}(\text{CH}_2\text{OMes})_2(\text{CH}_2\text{Br})$ (7.28 g, 21 mmol) in 150 mL THF. The solution was cooled to -40 °C and the LiPPh_2 -solution was added slowly over a 2 h period. The temperature was maintained at -40 °C for 3 h before being slowly warmed up to room temperature and the orange mixture was stirred overnight. A solution of $\text{BH}_3\cdot\text{SMe}_2$ (35 mL, 71 mmol) in THF (2.0 M) was slowly added to the light yellow reaction mixture. The resulting mixture was stirred for 2 h and the solvent was removed under vacuum. 250 mL diethyl ether (250 mL) and 250 mL water were added to the white solid. The organic layer was separated and washed with 250 mL water, dried over MgSO_4 and the organic solvent was removed under vacuum. The white solid was loaded onto a silica column (10 x 4 cm) and elucidated with a 70:30 mixture of ether and petroleum ether. The first fraction was discarded. The second fraction was collected to give $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2\cdot\text{BH}_3)_2(\text{CH}_2\text{OMes})$ **6-PG** (7.76 g, 14 mmol, 65 % yield) as white powder after the solvent was removed under vacuum.

$^{31}\text{P}\{^1\text{H}\}$ NMR (81.0 MHz, CD_2Cl_2 , δ): 8.9 (br, 2P); $^{11}\text{B}\{^1\text{H}\}$ NMR (128.3 MHz, CD_2Cl_2 , δ): -37.5 (br, 2B); ^1H NMR (399.9 MHz, CD_2Cl_2 , δ): 7.78 – 7.73 (m, 4 H, H_o), 7.65 – 7.60 (m, 4H, H_o), 7.51 (m, 12H, H_m and H_p), 4.28 (s, 2H, CH_2OMes), 3.02 – 2.94 (m, 2H, CH_2P), 2.92 (s, 3H, SCH_3), 2.52 – 2.45 (m, 2H, CH_2P), ~1.15 (broad in baseline, 6H, BH_3), 0.86 (s, 3H, CH_3); $^{13}\text{C}\{^1\text{H}\}$ NMR (100.6 MHz, CD_2Cl_2 , δ): 132.7 (d, $^2J_{\text{CP}} = 9$ Hz, C_o), 132.2 (d, $^2J_{\text{CP}} = 9$ Hz, C_o), 131.8 (d, $^4J_{\text{CP}} = 2$ Hz, C_p), 131.7 (d, $^4J_{\text{CP}} = 2$ Hz, C_p), 130.8 (d, $^1J_{\text{CP}} = 33$ Hz, C_i), 130.2 (d, $^1J_{\text{CP}} = 32$ Hz, C_i), 129.4 (d, $^3J_{\text{CP}} = 2$ Hz, C_m), 129.3 (d, $^3J_{\text{CP}} = 2$ Hz, C_m), 76.7 (t, $^3J_{\text{CP}} = 5$ Hz, CH_2OMes), 40.1 (s, C), 37.6 (s, SCH_3), 34.9 (d, $^1J_{\text{CP}} = 6$ Hz, CH_2P), 34.6 (d, $^1J_{\text{CP}} = 6$ Hz, CH_2P), 23.5 (t, $^3J_{\text{CP}} = 4$ Hz, CH_3);

HR-MS (FAB): $[\text{M}]^+ - \text{BH}_4$; theoretical $\text{C}_{30}\text{H}_{34}\text{BO}_3\text{P}_2\text{S}$: 547.1797; experimental 547.1813.

Elemental Analysis Calc. (Found): C 64.08 % (64.5 %), H 6.81 % (6.95 %)

Synthesis of $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2(\text{CH}_2\text{OMes})$ (6)

A 100 mL Teflon capped Schlenk flask was charged with $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2\cdot\text{BH}_3)_2(\text{CH}_2\text{OMes})$ (3.16 g, 5.6 mmol), DABCO (1.89 g, 17 mmol) and 20 mL toluene. The headspace was evacuated and the flask was heated at 80 °C for 2 h under static vacuum. The solution was filtered through a small plug of silica (3 cm) on a glass filter frit and washed with toluene (3 x 100 mL). The solvent was removed from the supernatant to give $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2(\text{CH}_2\text{OMes})$ **6** (1.57 g, 2.9 mmol, 52 % yield) as a thick white oil.

$^{31}\text{P}\{^1\text{H}\}$ NMR (81.0 MHz, CD_2Cl_2 , δ): -27.2 (s); ^1H NMR (200.1 MHz, CD_2Cl_2 , δ): 7.47 – 7.38 (m, 8 H, CH), 7.35 – 7.27 (m, 12 H, CH), 4.11 (s, 2H, CH_2OMes), 2.70 (s, 3H, SCH_3), 2.51 – 2.31 (m, 2H, CH_2P), 1.00 (s, 3H, CH_3).

Synthesis of $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2\cdot\text{BH}_3)_2(\text{CH}_2\text{P}(p\text{-tol})_2\cdot\text{BH}_3)$ (2-PG)

A 100 mL round-bottom Schlenk flask was charged with $\text{HP}(p\text{-tol})_2$ (0.344 g, 2.1 mmol) and 25 mL THF. The colorless solution was cooled in an ice bath and 2.5 M *n*-BuLi in hexane (0.89 mL, 2.2 mmol) was slowly added via syringe to give a red/orange solution. The solution was stirred for 30 min before drop wise addition to a 25 mL THF solution of $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2(\text{CH}_2\text{OMes})$ (845 mg, 1.58 mmol) in a 100 mL round-bottom Schlenk flask cooled to -40 °C. After the addition the solution was slowly warmed to room temperature and stirred overnight. The reaction mixture was additionally heated to reflux for 2 h. A solution of $\text{BH}_3\cdot\text{SMe}_2$ (2.84 mL, 5.69 mmol) in THF (2.0 M) was slowly added to the reaction mixture. The resulting mixture was stirred overnight and the volatile compounds were removed under vacuum. 100 mL Et_2O and 100 mL water were added to the white solid. The organic layer was separated and the aqueous layer was extracted with Et_2O for 2 times. The combined organic layer was washed with 100 mL water for 2 times, and then dried over MgSO_4 . After the removal of Et_2O under vacuum, the white residue in 20 mL Et_2O was diffused into hexane to give a white powder. The obtained powder was filtered off and washed 2 times with 10 mL hexane. After drying in vacuum,

ARTICLE

Journal Name

$\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2\bullet\text{BH}_3)_2(\text{CH}_2\text{P}(p\text{-tol})_2\bullet\text{BH}_3)$ **2-PG** (0.710 g, 1.02 mmol, 65 % yield) was obtained as a white powder.

$^{31}\text{P}\{^1\text{H}\}$ NMR (81.0 MHz, CD_2Cl_2 , δ): 9.2 (br, 2P), 7.4 (br, 1P); ^1H NMR (200.1 MHz, CD_2Cl_2 , δ): 7.66 – 7.37 (m, 24 H, CH), 7.24 – 7.19 (m, 4 H, CH), 2.93 – 2.82 (m, 6H, CH_2P), 2.36 (s, 6H, CH_3), ~1.17 (broad in baseline, 9H, BH_3), 0.79 (s, 3H, CH_3).

Elemental Analysis Calc. (Found): C 74.39 % (73.38 %), H 7.55 % (7.55 %)

Synthesis of $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2[\text{CH}_2\text{P}(p\text{-tol})_2]$ (**2**)

A 100 mL Teflon capped Schlenk flask was charged with $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2\bullet\text{BH}_3)_2[\text{CH}_2\text{P}(p\text{-tol})_2\bullet\text{BH}_3]$ (**11a**, 0.689 g, 0.99 mmol), DABCO (0.390 g, 3.47 mmol), and 20 mL toluene. The headspace was evacuated and the flask was heated at 80 °C for 2 h under static vacuum. The solution was filtered through a small plug of silica (3 cm) on a glass filter frit and washed with toluene (2 x 100 mL). The solvent was removed from the supernatant to give $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2[\text{CH}_2\text{P}(p\text{-tol})_2]$ **2** (0.663 g, 0.99 mmol, quantitative yield) as a thick white oil.

$^{31}\text{P}\{^1\text{H}\}$ NMR (81.0 MHz, CD_2Cl_2 , δ): -28.8 (t, $^4J_{\text{PP}} = 2.6$ Hz, 1P) -26.2 (d); ^1H NMR (200.1 MHz, CD_2Cl_2 , δ): 7.38 – 7.19 (m, 26 H, CH), 7.10 – 7.06 (m, 2 H, CH), 2.43 – 2.34 (m, 6H, CH_2P), 2.31 (s, 6H, CH_3), 0.93 (s, 3H, CH_3).

Synthesis of $\text{RuHCl}(\text{CO})\{\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2[\text{CH}_2\text{P}(p\text{-tol})_2]\}$ (**B**)

A round-bottom Schlenk flask was charged with $\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2[\text{CH}_2\text{P}(p\text{-tol})_2]$ (0.663 g, 1.0 mmol), $\text{RuHCl}(\text{CO})(\text{PPh}_3)_3$ (0.881 g, 0.92 mmol) and 40 mL toluene. The colorless solution with an off white precipitate was heated at reflux for 2 h to give a light yellow precipitate. The solution was filtered through a filter frit, the yellow residue washed three times with 10 mL toluene and dried in vacuum to give $\text{RuHCl}(\text{CO})\{\text{H}_3\text{CC}(\text{CH}_2\text{PPh}_2)_2[\text{CH}_2\text{P}(p\text{-tol})_2]\}$ **B** (0.544 g, 0.66 mmol, 72 % yield) as a isomeric mixture.

$^{31}\text{P}\{^1\text{H}\}$ NMR (81.0 MHz, CD_2Cl_2 , δ): 48.5 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 18$ Hz, 1P), 48.4 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 18$ Hz, 1P), 46.2 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 17$ Hz, 1P), 13.5 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 32$ Hz, 1P), 13.3 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 32$ Hz, 1P), 12.2 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 32$ Hz, 1P), 0.79 (dd, $^2J_{\text{PP}} = 32$ Hz, $^2J_{\text{PP}} = 18$ Hz, 1P), 0.58 (dd, $^2J_{\text{PP}} = 32$ Hz, $^2J_{\text{PP}} = 18$ Hz, 1P), -1.4 (dd, $^2J_{\text{PP}} = 32$ Hz, $^2J_{\text{PP}} = 18$ Hz, 1P); ^1H NMR (200.1 MHz, CD_2Cl_2 , δ): 7.83 – 6.53 (m, 28H, CH), 2.36 – 2.19 (m, 12H, CH_2 and CH_3), 1.54 – 1.51 (m, 3H, CH_3), -5.68 – -6.37 (m, 1H, RuH).

IR (KBr): 512, 697, 739, 837, 1093, 1434, 1483, 1923 (m, $\nu_{\text{Ru-H}}$), 1970 (vs, ν_{CO}), 2918, 2954, 3051 cm^{-1} .

Elemental Analysis Calc. (Found): C 64.58 % (64.16 %), H 5.42 % (5.32 %)

Improved synthesis of $\text{H}_3\text{CC}[\text{CH}_2\text{P}(p\text{-tol})_2]_3$ (**3**)

A 500 mL round-bottom Schlenk flask was charged with $\text{HP}(p\text{-tol})_2$ (3.06 g, 14 mmol) and 80 mL THF. The colorless solution was cooled in an ice bath and 2.5 M *n*-BuLi in hexanes (6 mL, 15 mmol) was slowly added via syringe to give a red/orange solution. The solution was stirred for 30 min before drop wise addition to a THF (100 mL) solution of $\text{H}_3\text{CC}(\text{CH}_2\text{OMes})_2(\text{CH}_2\text{Br})$ (**7**, 1.21 g, 3.6 mmol) in a 500 mL round-bottom Schlenk flask cooled to -40 °C. The resulting mixture was slowly warmed to room temperature, then heated at reflux overnight. A solution of $\text{BH}_3\bullet\text{SMe}_2$ (8.0 mL, 16 mmol) in THF (2.0 M) was slowly added to the colorless solution. The resulting mixture was stirred for 2 h and the solvent was removed under vacuum. 100 mL Ether and 100 mL water were added to the white solid. The organic layer was separated and washed with 150 mL water, dried over MgSO_4 and the organic solvent was removed under vacuum. The residue was suspended in 20 mL diethylether, 20 mL hexane was added, the product filtered off and washed two times with 20 mL hexane. After drying in vacuum $\text{H}_3\text{CC}[\text{CH}_2\text{P}(p\text{-tol})_2\bullet\text{BH}_3]_3$ (**3-PG**, 1.92 g, 2.6 mmol, 72 % yield) was obtained as a white powder. A 100 mL Teflon capped Schlenk flask was charged with $\text{H}_3\text{CC}[\text{CH}_2\text{P}(p\text{-tol})_2\bullet\text{BH}_3]_3$ (**3-PG**, 1.92 g, 2.6 mmol), DABCO (1.58 g, 14.1 mmol), and 20 mL toluene. The headspace was evacuated and the flask was heated at 80 °C for 2 days under static vacuum. The solution was filtered through a small plug of silica (3 cm) on a glass filter frit and the product extracted from the residue by washing it twice with 100 mL toluene. The solvent was removed from the combined organic layers to obtain a thick oil. After trituration of the oil with diethylether, a white powder was formed, which was filtered off and dried in vacuum to give $\text{H}_3\text{CC}[\text{CH}_2\text{P}(p\text{-tol})_2]_3$ **3** (1.03 g, 0.18 mmol, 72 % yield; 51 % yield overall) as a sticky white solid. All spectroscopic data was consistent with those reported in the literature.⁸

Synthesis of $\text{RuHCl}(\text{CO})\{\text{H}_3\text{CC}[\text{CH}_2\text{P}(p\text{-tol})_2]_3\}$ (**C**)

A round-bottom Schlenk flask was charged with $\text{H}_3\text{CC}[\text{CH}_2\text{P}(p\text{-tol})_2]_3$ (500 mg, 0.71 mmol), $\text{RuHCl}(\text{CO})(\text{PPh}_3)_3$ (612 mg, 0.64 mmol) and 40 mL toluene. The colorless solution with an off white precipitate heated at reflux for 2 h to give a light yellow precipitate. 10 mL pentane was added to the reaction mixture, the precipitate was filtered off via a sinter frit and washed three times with 10 mL pentane. After drying in vacuum, $\text{RuHCl}(\text{CO})\{\text{H}_3\text{CC}[\text{CH}_2\text{P}(p\text{-tol})_2]_3\}$ **C** (499 mg, 0.57 mmol, 80 % yield) was obtained as a yellow powder.

$^{31}\text{P}\{^1\text{H}\}$ NMR (81.0 MHz, CD_2Cl_2 , δ): 46.5 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 18$ Hz, 1P), 12.6 (dd, $^2J_{\text{PP}} = 40$ Hz, $^2J_{\text{PP}} = 32$ Hz), -0.8 ($^2J_{\text{PP}} = 32$ Hz, $^2J_{\text{PP}} = 18$ Hz); ^1H NMR (200.1 MHz, CD_2Cl_2 , δ): 7.73 – 6.58 (m, 24 H, CH), 2.52 – 2.08 (m, 15 H, CH_2 and CH_3), 1.50 – 1.46 (m, 3H, CH_3), -6.02 (ddd, $^2J_{\text{HP}} = 93.6$ Hz, $^2J_{\text{HP}} = 18.8$ Hz, $^2J_{\text{HP}} = 14.9$ Hz, 1H, RuH).

IR (KBr): 521, 558, 624, 713, 735, 804, 838, 1020, 1092, 1190, 1397, 1440, 1499, 1599, 1895 (m, $\nu_{\text{Ru-H}}$), 1979 (vs, ν_{CO}), 2866, 2920, 2948, 3019 cm^{-1} .

Elemental Analysis Calc. (Found): C 65.93 % (65.75 %), H 5.99 % (6.01 %)

Reaction of RuHCl(CO)(PPh₃)₃ with H₃CC(CH₂PPh₂)₂(2-pyridyl)

A round-bottom Schlenk flask was charged with H₃CC(CH₂PPh₂)₂(2-pyridyl) **4** (160 mg, 0.31 mmol, 1.2 equiv.), (PPh₃)₃RuHCl(CO) (250 mg, 0.26 mmol, 1 equiv.) and dichloromethane (40 mL). The light yellow solution was stirred at room temperature for 5 days to light yellow solution. The solution was filtered to remove a small amount of white precipitate and the volume was reduced to 2 mL. A fine yellow precipitate was obtained with the addition of hexane (~20 mL). The yellow solid contained a mixture of **D-1**, **D-2**, and **D-3**. Recrystallization by slow solvent diffusion of pentane (20 mL) into a concentrated DCM solution (2 mL) gave a small amount of **21** (~20 mg) in reasonable purity (~95 %). A second recrystallization attempt of the same sample gave a mixture of **D-1**, **D-2** and **D-3**. Single crystal of **D-3** were isolated from the sample and used for x-ray diffraction analysis.

MS (LIFDI, crude mixture): *m/z* (%): 896.2 (100 %, **D-1,2**⁺ – Cl), 669.0 (25 %, **D-3**⁺ – Cl).

D-1: ³¹P{¹H} NMR (121.4 MHz, CD₂Cl₂, δ): 44.9 (dd, ²J_{PP} = 280 Hz, ²J_{PP} = 20 Hz), 28.6 (dd, ²J_{PP} = 280 Hz, ²J_{PP} = 20 Hz), 3.85 (t, ²J_{PP} = 20 Hz); ¹H NMR (300.1 MHz, CD₂Cl₂, δ): -6.55 (ddd, ²J_{HP} = 18 Hz, ²J_{HP} = 22 Hz, ²J_{HP} = 108 Hz), all other peaks are overlapping with those of **D-2** and **D-3**.

D-2: ³¹P{¹H} NMR (121.4 MHz, CD₂Cl₂, δ): 45.9 (dd, ²J_{PP} = 230 Hz, ²J_{PP} = 20 Hz), 34.2 (dd, ²J_{PP} = 230 Hz, ²J_{PP} = 20 Hz), 9.8 (t, ²J_{PP} = 20 Hz); ¹H NMR (300.1 MHz, CD₂Cl₂, δ): -5.19 (dt, ²J_{HP} = 17 Hz, ²J_{HP} = 92 Hz), all other peaks are overlapping with those of **D-1** and **D-3**.

D-3: ³¹P{¹H} NMR (121.4 MHz, CD₂Cl₂, δ): 37.9 ppm (s); ¹H NMR (300.1 MHz, CD₂Cl₂, δ): all peaks are overlapping with those of **D-1** and **D-2**.

Synthesis of RuHCl(CO)(PPh₃)[H₃CC(CH₂PPh₂)₂(CH₂OH)] (**E**)

In a glovebox, a round-bottom Schlenk flask was charged with H₃CC(CH₂PPh₂)₂(CH₂OH) **5** (0.502 g, 1.1 mmol), RuHCl(CO)(PPh₃)₃ (952 mg, 1.0 mmol) and 40 mL toluene. The colorless solution with an off white precipitate heated at reflux for 2 h, then the volume of toluene was reduced under vacuum to ca. 10 mL. In the glovebox, 10 mL pentane was added to the reaction mixture, forming a pale yellow powder. The precipitate was filtered off via a sinter frit and washed three times with 10 mL pentane. After drying in vacuum, RuHCl(CO)(PPh₃)[H₃CC(CH₂PPh₂)₂(CH₂OH)] **E** (0.610 g, 0.69 mmol, 69 % yield) was obtained as a pale yellow powder. The ³¹P NMR indicated that the isolated compounds were the mixture of regioisomers of **E**.

IR (KBr): 517, 695, 742, 806, 839, 1054, 1091, 1158, 1188, 1312, 1434, 1481, 1586, 1924 (vs, ν_{CO} and ν_{Ru-H}), 2928, 3053 cm⁻¹

Elemental Analysis Calc. (Found): C 65.19 % (62.58 %), H 5.24 % (5.15 %)

³¹P{¹H} NMR (162.1 MHz, CD₂Cl₂, δ): 44.2 (dd, ²J_{PP} = 282 Hz, ²J_{PP} = 20 Hz), 43.6 (dd, ²J_{PP} = 282 Hz, ²J_{PP} = 20 Hz), 27.4 (dd, ²J_{PP} = 282 Hz, ²J_{PP} = 20 Hz), 26.9 (dd, ²J_{PP} = 282 Hz, ²J_{PP} = 20 Hz), 3.0 (t, ²J_{PP} = 20 Hz), 0.4 (t, ²J_{PP} = 20 Hz). All peaks are derived from the regioisomers of **E**; ¹H NMR (400.3 MHz, CD₂Cl₂, δ): 8.05 – 6.80 (m, 35H, CH), 3.30 – 0.18 (m, 9H, CH₂ and CH₃), -6.73 (ddd, ²J_{HP} = 109 Hz, ²J_{HP} = 23 Hz, ²J_{HP} = 17 Hz, 1H, RuH), -6.83 (ddd, ²J_{HP} = 109 Hz, ²J_{HP} = 23 Hz, ²J_{HP} = 17 Hz, 1H, RuH). All peaks are derived from the regioisomers of **E**.

Reaction and RuHCl(CO)(PPh₃)[H₃CC(CH₂PPh₂)₂(CH₂OH)] (**E**) with *t*-BuOK

In a glovebox, a round-bottom Schlenk flask was charged with RuHCl(CO)(PPh₃)[H₃CC(CH₂PPh₂)₂(CH₂OH)] **E** (0.114 g, 0.13 mmol), *t*-BuOK (14.4 mg, 0.13 mmol) and 5 mL toluene, forming a red solution at room temperature in 2 h. All volatile compounds were removed under vacuum, then 0.5 mL toluene and 5 mL pentane was added to a red powder. The resulting suspension was passed through a pad of celite, then a red filtrate was dried under vacuum to give alkoxo ruthenium complex **F** (81.3 mg, 0.096 mmol, 75 % yield) as a deep red powder. The ¹H and ³¹P NMR spectra of the obtained powder suggested that it was constituted of mainly alkoxo ruthenium complex **F** with some impurities including small amounts of the unreacted **E**.

Characteristic peaks of **F** are described below:

³¹P{¹H} NMR (243.0 MHz, *d*₈-THF, δ): 35.7 (dd, ²J_{PP} = 276 Hz, ²J_{PP} = 20 Hz), 33.5 (dd, ²J_{PP} = 276 Hz, ²J_{PP} = 20 Hz), 13.7 (dd, ²J_{PP} = 20 Hz); ¹H NMR (600.2 MHz, *d*₈-THF, δ): -4.65 (ddd, ²J_{HP} = 113 Hz, ²J_{HP} = 26 Hz, ²J_{HP} = 13 Hz, 1H, RuH). CH₃ peaks from *t*-Bu group were not observed in ¹H NMR spectra.

IR (KBr): 515, 695, 742, 836, 999, 1092, 1186, 1434, 1480, 1586, 1896 (m, ν_{Ru-H}), 1910 (vs, ν_{CO}), 2674, 2754, 2921, 2951, 3051 cm⁻¹.

MS (LIFDI): *m/z* (%): 848.01 (100 %, **F**⁺).

A representative procedure for catalyst screening described in Table 1

Entry 1: In an argon-filled glovebox, to a premix autoclave (60 mL, stainless steel) equipped with a magnetically coupled propeller blade stirrer was added RuHCl(CO)(triphos) **A** (36 mg, 0.047 mmol), toluene (17 mL), and 1-octanol (3.0 g, 24 mmol), then the autoclave was sealed. After introducing NH₃ gas (7-8 bar) at room

ARTICLE

Journal Name

temperature, the autoclave was heated to 165 °C for 15 h with vigorous stirring (700–800 rpm) without reintroducing NH₃. After cooling to the room temperature, the pressurized NH₃ gas in the autoclave was released in a fume hood. An aliquot of the reaction mixture was taken for the GC analysis, and the conversion of 1-octanol and the yield of amine products were calculated based on GC area %.

Entry 2: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol) and triphos (33 mg, 0.052 mmol) were used.

Entry 3: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol) was used.

Entry 4: RuHCl(CO){H₃CC(CH₂PPh₂)₂[CH₂P(*p*-tol)₂]} (38 mg, 0.047 mmol) was used.

Entry 5: RuHCl(CO){H₃CC[CH₂P(*p*-tol)₂]₃} (40 mg, 0.047 mmol) was used.

Entry 6: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol) and CH₃C(CH₂PPh₂)₂(2-pyridyl) **4** (26 mg, 0.046 mmol) were used.

Entry 7: RuHCl(CO)(PPh₃)[H₃CC(CH₂PPh₂)₂(CH₂OH)] **E** (41 mg, 0.047 mmol) was used.

Entry 8: The proposed alkoxo-ruthenium complex **F** (20 mg, 0.023 mmol), toluene (8.5 mL), and 1-octanol (1.5 g, 12 mmol) were used.

Entry 9: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol) and CH₃C(CH₂PPh₂)₂(CH₂OMs) **6** (28 mg, 0.052 mmol) were used.

Entry 10: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol) and dppdmp (23 mg, 0.052 mmol) were used.

Entry 11: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol) and dppp (21 mg, 0.052 mmol) were used.

Entry 12: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol), dppp (21 mg, 0.052 mmol), and toluene (6 mL) were used.

Entry 13: RuHCl(CO)(PPh₃)₃ (45 mg, 0.047 mmol), dppp (21 mg, 0.052 mmol), toluene (6 mL), and p(NH₃) = 4 bar (53 mmol of NH₃) were used.


Acknowledgements

CaRLa (Catalyst Research Laboratory) is co-financed by Rupprechts-Karls-University Heidelberg (Heidelberg University) and BASF SE.

Notes and references

- (a) B. R. Brown, *The Organic Chemistry of Aliphatic Nitrogen Compounds*, Oxford University, New York, 1994; (b) S. A. Lawrence, *Amines: Synthesis, Properties and Applications*, Cambridge University Press, Cambridge, 2005; (c) P. Roose, K. Eller, E. Henkes, R. Rossbacher and H. Höke, *Amines, Aliphatic: Ullmann's Encyclopedia of Industrial Chemistry*, Wiley-VCH, Weinheim, 2015.
- R. N. Salvatore, C. H. Yoon and K. W. Jung, *Tetrahedron*, 2001, **57**, 7785.
- (a) J. Kim, H. J. Kim and S. Chang, *Eur. J. Org. Chem.*, 2013, 3201; (b) J. L. Klinkenberg and J. F. Hartwig, *Angew. Chem. Int. Ed.*, 2011, **50**, 86; (c) S. Bähn, S. Imm, L. Neubert, M. Zhang, H. Neumann and M. Beller, *ChemCatChem*, 2011, **3**, 1853.

- Recent reviews, see: (a) S. Bähn, S. Imm, L. Neubert, M. Zhang, H. Neumann and M. Beller, *ChemCatChem*, 2011, **3**, 1852; (b) C. Gunanathan and D. Milstein, *Chem. Rev.*, 2014, **114**, 12024; (c) K. Shimizu, *Catal. Sci. Technol.*, 2015, **5**, 1412; (d) Q. Yang, Q. Wang and Z. Yu, *Chem. Soc. Rev.*, 2015, **44**, 2305.
- (a) C. Gunanathan and D. Milstein, *Angew. Chem. Int. Ed.*, 2008, **47**, 8661; (b) D. Pingen, C. Müller and D. Vogt, *Angew. Chem. Int. Ed.*, 2010, **49**, 8130; (c) S. Imm, S. Bähn, L. Neubert, H. Neumann and M. Beller, *Angew. Chem. Int. Ed.*, 2010, **49**, 8126; (d) G. Walther, J. Deutsch, A. Martin, F. E. Baumann, D. Fridag, R. Franke and A. Köckritz, *ChemSusChem*, 2011, **4**, 1052; (e) W. Baumann, A. Spannenberg, J. Pfeffer, T. Haas, A. Köckritz, A. Martin and J. Deutsch, *Chem. – Eur. J.*, 2013, **19**, 17702; (f) D. Pingen, O. Diebolt and D. Vogt, *ChemCatChem*, 2013, **5**, 2905; (g) D. Pingen, M. Lutz and D. Vogt, *Organometallics*, 2014, **33**, 1623; (h) X. Ye, P. N. Plessow, M. K. Brinks, M. Schelwies, T. Schaub, F. Rominger, R. Paciello, M. Limbach and P. Hofmann, *J. Am. Chem. Soc.*, 2014, **136**, 5923.
- E. Balaraman, D. Srimani, Y. D. Posner and D. Milstein, *Catal. Lett.*, 2015, **145**, 139.
- (a) S. Imm, S. Bähn, M. Zhang, L. Neubert, H. Neumann, F. Klasovsky, J. Pfeffer, T. Haas and M. Beller, *Angew. Chem. Int. Ed.*, 2011, **50**, 7599; (b) E. J. Derrah, M. Hanauer, P. N. Plessow, M. Schelwies, M. K. da Silva and T. Schaub, *Organometallics*, 2015, **34**, 1872; (c) T. Schaub, B. Buschhaus, M. K. Brinks, M. Schelwies, R. Paciello, J. P. Melder and M. Merger, WO2012119927.
- (a) H. Heide, G. Huttner and G. Helmchen, *Z. Naturforsch.*, 1993, **48b**, 1681; (b) A. Muth, O. Walter, G. Huttner, A. Asam, L. Zsolnai and C. Emerich, *J. Organomet. Chem.*, 1994, **468**, 149.
- S. Doherty, E. G. Robins, M. Nieuwenhuyzen, P. A. Champkin and W. Clegg, *Organometallics*, 2002, **21**, 4147.
- (a) T. Seitz, A. Muth and G. Huttner, *Chem. Ber.*, 1994, **127**, 1837; (b) Y.-T. Hsieh, C.-M. Cheng, M.-S. Peng and T.-S. Liu, *J. Chem. Soc. Dalton. Trans.*, 1994, 3499.
- Attempts to install other diorganophosphino moiety or heteroatom such as P(*o*-tolyl)₂, P(*p*-F-C₆H₄)₂, PCy₂, P*i*Pr₂, and NH₂, were unsuccessful by using our approach.
- K.-M. Sung, S. Huh and J.-M. Jun, *Polyhedron*, 1999, **18**, 469.
- RuHCl(CO)(triphos) **A** was characterised by X-ray analysis, see: S. Savourey, G. Lefèvre, J.-C. Berthert, P. Thuéry, C. Genre and T. Contat, *Angew. Chem. Int. Ed.*, 2014, **53**, 10466.
- Preparation and characterization of RuHCl(CO)(dppp)(PPh₃) from RuHCl(CO)(PPh₃)₃ and dppp was reported, see: (a) A. Santos, J. López, J. Montoya, P. Noheda, A. Romero and A. M. Echavarren, *Organometallics*, 1994, **13**, 3605; (b) S. Huh, Y. Cho and J.-M. Jun, *Polyhedron*, 1994, **13**, 1887.
- Cyclohexanol and ammonia undergo the selective mono-alkylation with dppp ligand. See ref. 4e.
- In Fig 6, p(NH₃) = 4.1 – 4.5 bar was used, which corresponded to approximately 53 mmol of NH₃ in the autoclave.
- (a) R. Yamaguchi, S. Kawagoe, C. Asai and K. Fujita, *Org. Lett.*, 2008, **10**, 181; (b) R. Kawahara, K. Fujita and R. Yamaguchi, *J. Am. Chem. Soc.*, 2010, **132**, 15108. (c) S. Wöckel, M. Schelweis, M. K. Brinks, F. Rominger, P. Hofmann and M. Limbach, *Org. Lett.*, 2013, **15**, 266.
- Program SADABS 2012/1 for absorption correction, see: G. M. Sheldrick, *Bruker Analytical X-Ray Division*, Madison, Wisconsin, 2012.
- L. González-Sebastián, M. Flores-Alamo and J. J. García, *Organometallics*, 2012, **31**, 8200.



RSC graphical abstract: 4 cm * 8 cm as a maxim side