


RSC Advances



This article can be cited before page numbers have been issued, to do this please use: M. Suchy, A. X. Li, M. Milne, R. Bartha and R. H. E. Hudson, *RSC Adv.*, 2016, DOI: 10.1039/C6RA11741D.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

DOTMA-based amides (DOTMAMs) as a platform for the development of PARACEST MRI contrast agents

M. Suchý,^{a,c} A.X. Li,^c M. Milne,^a R. Bartha^{c,d} and R.H.E. Hudson^{a,b*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

A synthetic methodology leading to previously unknown DOTMA-based secondary amides (DOTMAMs) has been developed. Alkylation of cyclen with *L*-lactic acid-derived pseudohalides was used as a key step affording the alkyl- and aryl-decorated DOTMAMs. Amino acid-decorated DOTMAMs were obtained *via* peptide coupling between DOTMA and protected amino acids. Metallation of the DOTMAMs ligand with Tm³⁺ gave complexes exhibiting proximal amide proton based paramagnetic CEST effects at < -50 ppm relative to water.

Introduction

Nowadays magnetic resonance imaging (MRI) represents an indispensable tool in clinical diagnostics. MRI is primarily based on the detection of water protons in the body and oftentimes is used in combination with an exogenous agent that enhances the image contrast.¹ Such agents are termed 'contrast agents' (CAs) and the development of CAs suitable for MRI is an active area of research. Several types of CAs are currently available,² among them paramagnetic chemical exchange saturation transfer (PARACEST) MRI CAs³ represent a relatively new group characterized by exceptional sensitivity to various parameters relevant to clinical diagnostics, such as temperature,⁴ pH,⁵ redox state,⁶ concentration of primary metabolites⁷ or enzymatic activity.⁸ Despite the potential usefulness of PARACEST CAs, their low *in vivo* sensitivity requires the administration of high concentrations (20–100 mM) of CAs;⁹ thus, it is important to employ ligands that stably bind the potentially toxic lanthanide metal, such as those based on the macrocyclic chelator DOTA, and employ strategies to increase the observable signals.¹⁰

Upon administration into a biological system PARACEST MRI CAs experience a significant decrease in sensitivity attributable

to magnetization transfer (MT) from endogenous macromolecules.^{11,12} The MT effect is a competing mechanism capable of saturating the bulk water spins and consequently lowers the contrast efficiency of PARACEST MRI CAs within the MT frequency range (*ca.* -100 to 100 ppm). Development of PARACEST MRI CAs exhibiting highly shifted CEST signals lying toward or beyond the outer limit of the MT frequency range represents a way to address the low sensitivity associated with these agents. In this regard, we recently demonstrated Tm³⁺ DOTAM-*t*-butyl (**1**, Figure 1),^{13,†} possessed an amide proton-based CEST effect clearly lying outside of the MT frequency range. The presence of the highly shifted CEST effect ($\delta = -102$ ppm) was attributed to the predominance of the twisted square antiprism (TSAP) isomer of CA **1** in solution¹³ which was favoured by the sterically demanding *t*-butylamides.¹⁴

Unlike the Tm³⁺ complexes derived from DOTA (**2**, Figure 1) and unsubstituted DOTAM (**3**, Figure 1) which exist in solution predominantly (*ca.* 90%) in the square antiprism isomer (SAP), the Tm³⁺ complex derived from DOTMA^{15,§} (**4**, Figure 1) favours the TSAP isomer in solution (*ca.* 90%).¹⁶ This conformational bias is attributed to the steric effect due to the presence of the additional four methyl groups on the side arms of DOTMA (**4**). We hypothesized tetraamides derived from DOTMA, herein referred to as DOTMAMs,^{§§} may possess highly shifted CEST signals similar to that observed for **1**, and have a greater preference for the TSAP isomer as observed for the carboxylic acid DOTMA, thus potentially resulting in a more intense, highly-shifted CEST effect.

While the syntheses of various ligands based on the DOTA (**2**) or DOTAM (**3**) scaffold via peralkylation of cyclen (**5**, Figure 1) are well documented in the literature,¹⁷ only a small number of DOTMA-based ligands are presently known.^{16,18} Furthermore, only the primary amide derived from DOTMA

^a Department of Chemistry The University of Western Ontario, London ON N6A 5B7 Canada

^b Centre for Advanced Materials and Biomaterials Research, The University of Western Ontario

^c Center for Functional and Metabolic Mapping, Robarts Research Institute, The University of Western Ontario, London ON N6A 5K8 Canada

^d Department of Medical Biophysics, The University of Western Ontario

* Robert H. E. Hudson, phone +519-661-2111, ext. 86349; e-mail: rhudson@uwo.ca

† Electronic Supplementary Information (ESI) available: Full experimental details, NMR and MS spectra associated with synthetic intermediates and final products. CEST spectra and Ω -plots associated with CAs **6–8**, **10** and **11**. See DOI: 10.1039/x0xx00000x

ARTICLE

Journal Name

(DOTMAM)¹⁹ has been described. This ligand was accessed by the peralkylation of cyclen with racemic 2-propanionamide resulting in a mixture of stereoisomeric ligands of which only the Eu³⁺ chelate was studied.¹⁹

Our interest in Tm³⁺ chelates of DOTMAMs (with a secondary amide) required the development of a new synthetic methodology. Herein, we report the preparation of the hitherto unreported Tm³⁺ complex of known ligand **6** and well as the synthesis of five new ligands and the Tm³⁺ complexes thereof (**7-11**, Figure 1) as well as evaluation of their CEST properties.





Figure 1. Chemical structures of Tm³⁺ DOTAM-*t*-Bu (**1**), DOTA (**2**), DOTAM (**3**), DOTMA (**4**), cyclen (**5**) and the DOTMA-derived complexes **6-11**.

Scheme 1. Preparation of electrophiles **15a-15c**, **16a**, **16b**, **17a-19a**



Allen's group has recently reported an alkylation of *N,O*-macrocycles related to cyclen by OTs-(*S*)-lactamide.²² The conversion of lactamide **15** to OTs-(*S*)-benzyl lactamide (**15c**,

Results and discussion

Our studies began with the synthesis of corresponding electrophiles, which we envisioned using for the peralkylation of cyclen (**5**). A DCC-mediated peptide coupling between (*S*)-2-chloropropionic acid (**12**, Scheme 1) and benzylamine afforded corresponding secondary chloroderivative **13** (Scheme 1) in good yield using a modified literature procedure.²⁰ Despite the use of 2-chloropropionic acid derivatives previously for the preparation of DOTMA,¹⁵ the reactivity of electrophile **13** was found to be insufficient to achieve the peralkylation of cyclen. Therefore, we turned our attention to the use of (*S*)-lactic acid derived pseudohalides as potential electrophiles. Coupling of (*S*)-lactic acid (**14**) with a small variety of amines,²¹ namely benzylamine, *t*-butylamine, aniline, and (*L*)-Lys(Boc)-OMe•HCl, proceeded smoothly and afforded the lactamides **15-18** (Scheme 1) in good yields (60-93%).


Firstly, conversion of the secondary OH group to the corresponding triflate (OTf-lactamide) was attempted. Among the few known preparations of DOTMA-based ligands, peralkylation of cyclen with OTf-(*S*)-lactic acid esters is used as a key step almost exclusively.^{16,18} Treatment of lactamide **15** with trifluoroacetic acid anhydride (Tf₂O)^{18a} afforded the unstable OTf-lactamide **15b** (two signals observed in ¹⁹F NMR spectrum). However, treatment of lactamide **16** with Tf₂O produced the more stable OTf-lactamide **16b** (one signal at δ -74.9 ppm in ¹⁹F NMR spectrum after storage for 24 hours at room temperature). Unfortunately, OTf-lactamide **16b** was not stable under the forcing conditions required for peralkylation of cyclen (heat, excess of base) resulting in the formation of complex mixture devoid of the desired product of peralkylation.

Scheme 1) proceeded smoothly (82% yield).²³ Unfortunately, the alkylation of cyclen with **15c** resulted in a complex mixture

containing only small amount (< 10%) of the desired product of peralkylation, obtained after careful HPLC purification.

The problem was solved by using OMs-(S)-lactamides **15a-18a** (Scheme 1) obtained in excellent yields (> 90%) after treatment of lactamides **15-18** with MsCl under basic conditions (Scheme 1).²⁴ With electrophiles **15a-18a** in hand we investigated the alkylation of cyclen. Mixtures of di-, tri- and tetrasubstituted cyclens have been obtained, when electrophiles **15a-17a** were used. The desired ligands were purified by HPLC, affording DOTMAM-Bn, DOTMAM-*t*-Bu and DOTMAM-Ph in low to moderate yields (14-42%) as described in the Supporting Information and shown in Scheme 2.


Scheme 2. Preparation of CAs 6-9



This methodology also proved successful for the preparation of the DOTMAM ligand with a primary amide group (Scheme 2). Thus **6** was obtained in 51% yield (see the Supporting Information for details) by peralkylation of cyclen with OMs-(S)-lactamide (**19a**, Scheme 1), followed by purification by flash column chromatography (FCC) on Al₂O₃.[†] Prolonged heating (48 h at 90 °C) of above mentioned ligands with TmCl₃·H₂O (in dioxane/water mixture, pH ~ 7) afforded the desired complexes **6-8** (Scheme 3), interestingly no metallation (complex **9**) was observed when DOTMAM-Ph was treated with TmCl₃·H₂O under identical conditions (Scheme 2).

Peralkylation of cyclen with OMs-lactamide **18a** did not lead to the formation of the desired DOTMAM-Lys(Boc)-OMe; therefore, we explored an alternative synthetic route to obtain the amino acid-decorated DOTMAM derived ligands. An HBTU-mediated coupling^{‡,25} between DOTMA tetrasodium salt (**4**, Scheme 3) prepared according to the literature procedures^{16,18a} and (L)-Lys(Boc)-OMe·HCl afforded the desired DOTMAM-(L)-Lys(Boc)-OMe (in 41% yield after the HPLC purification). Subsequent removal of protecting groups and metallation with TmCl₃·H₂O furnished the complex **10** in 27% overall yield (based on DOTMA tetrasodium salt) as shown in Scheme 3 and described in the Supporting Information. Conjugation of **4** with (L)-Phe-OEt·HCl, followed by removal of ester protecting groups by saponification and metallation with TmCl₃·H₂O gave the complex **11** in 25% overall yield (Scheme 3; see the Supporting Information for experimental details).

Scheme 3. Preparation of CAs 10 and 11



With complexes **6-8**, **10** and **11** prepared, their potential as PARACEST MRI CAs was investigated. The CEST spectra associated with CAs **6-8**, **10** and **11** were acquired as described in the Supporting Information. Temperature and pH sensitivity of the CAs **6-8**, **10** and **11** was also investigated. When possible, the results were compared with those obtained previously for related DOTAM-based complexes.^{13,26} The results of these studies are summarized in Table 1 and discussed below.

Table 1. CEST properties for CAs **6-8**, **10** and **11**.*

CA	CEST effect (ppm) signal intensity (%)	Related DOTAM complex; CEST effect (ppm)/signal intensity (%)
6	-54 (6); -63 (7)	-46 (32); -51 (33) ^{13b5}
7	-65 (6)	-51 (27) ^{13b5}
8	-115 (3)	-68 (10); -102 (21) ^{13a†}
10	-53 (20); -62 (10)	-50 (15) ^{26#}
11	-51 (1)	not available

* CEST spectra for **6-8**, **10** and **11** were collected on a 9.4 T clinical MRI scanner at 15 mM of complex, 37 °C, pH 7.5 using a saturation pulse of 14 μT for 3.95 s. ⁵ CEST spectra collected at: 14 T, 10 mM of complex, 37 °C, pH 7.0 using a saturation pulse of 650 Hz for 2 s. [†] CEST spectra collected at: 14 T, 10 mM complex (in 90% D₂O), 37 °C, pH 7.2 using a saturation pulse of 20 μT for 2 s. [#] CEST spectra collected at: 14 T, 10 mM complex, 37 °C, pH 7.0 using a saturation pulse of 20 μT for 10 s.

CA **6**, derived from the unsubstituted DOTMAM ligand, and CAs **7** and **8**, decorated with alkyl groups, all featured amide-proton based CEST effect with intensities that are 4-7 fold smaller (Table 1) than signals produced by the corresponding DOTAM-based counterparts (Tm³⁺ DOTAM, Tm³⁺ DOTAM-Bn and Tm³⁺ DOTAM-*t*-Bu).¹³ A CEST

signal at lower chemical shift was not observed for CA **8** (Table 1), presumably it was too weak to be detected. On the other hand, the CEST signals associated with CAs **6–8** were shifted farther from bulk water signal (by 8–14 ppm, Table 1) as compared to the analogous DOTAM CAs. The signal reduction of DOTAM-based CAs is caused by slower exchange rates (by *ca.* 2 fold) compared to related DOTAM-based complexes (244 μ s for **7** versus 125 μ s for Tm³⁺ DOTAM-Bn; 579 μ s for **8** versus 273 μ s for Tm³⁺ DOTAM-*t*-Bu).²⁷ Should these findings with respect to pendant arm amide proton exchange rates extend to a slowing of water exchange at the metal center, examination of Eu³⁺ DOTAM-based complexes (requiring slow exchange rates for bound water)³ may lead to the formation of PARACEST MRI CAs. These studies are beyond the scope of the present work.

Tm³⁺ DOTMAM-(L)-Lys-OH (**10**) possessed two CEST effects (Table 1), both showed a dependence on pH and temperature (Figure 2). The origin of the two peaks is possibly due to the presence of *s-cis* and *s-trans* amide conformations as suggested by the similarity in chemical shifts of the CEST signals to those present in the unsubstituted DOTMAM (**6**).²⁸




Figure 2. CEST spectra associated with Tm³⁺ DOTMAM-(L)-Lys-OH (**10**), measured at 9.4 T, 15 mM of complex, using a saturation pulse of 14 μ T for 3.95 s. The CEST effects are modulated by both pH (top) and temperature (bottom).

Curiously, only a very weak CEST effect (Table 1) was observed for Tm³⁺ DOTMAM-(L)-Phe-OH (**11**). The intensity of the signal was found to increase with increasing pH (see the Supporting Information), implying that the exchange kinetics associated with CA **11** is out of the optimal range (too slow)⁸ for the observation of the CEST effect at pH 7; presence of base increases the exchange rate of the side chain amide protons resulting in a somewhat stronger signal at pH 8 (see Supporting Information).²⁹ Our findings indicate that further variation of the amino acid residues³⁰ present in DOTMAM-derived CAs might lead to complexes endowed with interesting CEST properties.

Examination of the ¹H NMR spectra for Yb³⁺ DOTMAM, Yb³⁺ DOTMAM-Bn and Yb³⁺ DOTMAM-*t*-Bu (supporting information) indicate these complexes exist predominantly in as a single isomer. Comparison of the mostly highly shifted signal, the axial cyclen proton designated H₄, is commonly done discriminate coordination complex geometry (i.e. SAP versus TSAP),³¹ and estimate population distributions of the two isomers, but is unenlightening in the current case. Comparison of Eu³⁺ to Yb³⁺ metallated ligands show the following trends. The chemical shift of H₄ for Eu³⁺DOTAM (SAP isomer ~40 ppm; TSAP isomer ~5 ppm)¹⁹ less highly shifted than the Eu³⁺DOTAM complex (SAP isomer ~44–59 ppm; TSAP isomer ~15–20 ppm). The Yb³⁺DOTAM complex exists as a single isomer (SAP) displaying a chemical shift of H₄ = 95 ppm. The Yb³⁺DOTMAM complex prepared during this work shows H₄ = 124 ppm, which follows the trend of increasing chemical shift in going from the DOTAM to DOTMAM ligand observed for the Eu³⁺ complexes. The Yb³⁺DOTMAM-Bn shows a similar chemical shift for H₄ (127 ppm). It is notable that both the Tm³⁺DOTMAM and Tm³⁺DOTMAM-Bn complexes both give rise to relatively low-shifted amide-proton CEST signals. This is in contrast to Yb³⁺DOTMAM-*t*-Bu which possesses a lower shifted H₄ (104 ppm) yet the Tm³⁺ complex displays the CEST signal with the greatest hyperfine shift. It is tempting to suggest that the lower H₄ chemical shift for Tm³⁺DOTMAM-*t*-Bu indicates that it exists in a different coordination geometry, possibly TSAP, which is responsible for the high shifting CEST signal and would be consistent with our previous work.¹³

Experimental

General Experimental Procedures

Reagents were commercially available and all solvents were HPLC grade except for water (18.2 M Ω cm millipore water), CH₂Cl₂ and DMF (dried over Al₂O₃, in a solvent purification system). Solvents were removed under reduced pressure in a rotary evaporator, aqueous solutions were lyophilized and organic extracts were dried over Na₂SO₄. Flash column chromatography (FCC) was carried out using silica gel (SiO₂), mesh size 230 – 400 Å and basic alumina (Al₂O₃), pH 9.5–10.5, mesh size 10 – 100 Å. Thin-layer chromatography (TLC) was carried out on Al backed silica gel or alumina plates with compounds visualised by I₂ vapours, anisaldehyde stain, 5% ninhydrin stain, phosphomolybdic acid stain, and UV light. Melting points (mp) were obtained on Fisher–Johns apparatus

and are uncorrected. Specific rotations $[\alpha]_D$ were determined by polarimeter at ambient temperature using a 1 ml, 10 cm path length cell; the units are $10^{-1} \text{ deg cm}^2 \text{ g}^{-1}$ and the concentrations are reported in g/100 mL. Ultra performance liquid chromatography (UPLC) was performed using a BEH C18 column (particle size 1.7 μm ; 1.0 id \times 100 mm) and HR-ESI-MS detector. Mobile phase: Method A: 100% H_2O – 100% MeCN (both solvents containing 0.1% HCOOH) over 5 min, then 100% MeCN for 2 min, linear gradient, flow rate 0.1 mL/min. HPLC purification was performed using a Delta-Pak C₁₈ 300 Å column (particle size 15 μm ; 8 \times 100 mm Radial-Pak cartridge). Mobile phase for Method B (DOTMAM-Bn) was 0 min, 90% H_2O – 10% MeCN (for each method both solvents containing 0.1% TFA) to 13 min, 20% H_2O – 80% MeCN; Method C (DOTMAM-*t*-Bu, **7**): 0 min, 90% H_2O – 10% MeCN to 11 min, 13% H_2O – 87% MeCN; Method D (DOTMAM-Ph): 0 min, 90% H_2O – 10% MeCN to 13 min, 100% MeCN; Method E [DOTMAM-(L)-Lys(Boc)-OMe, DOTMAM-(L)-Phe-OEt]: 0 min, 90% H_2O – 10% MeCN to 10 min, 100% MeCN; Method F (**8**): 0 min, 90% H_2O – 10% MeCN to 11 min, 20% H_2O – 80% MeCN; linear gradient and 3 mL/min flowrate were used for each method. Size exclusion chromatography (SEC) was carried out on Bio-Gel P2, 45-90 μm mesh resin (8 g, per 0.1 mmol of compound). Ten fractions (10 ml each) were collected and identified with I_2 vapours and UV light. Absence of free Tm^{3+} was verified by xylenol orange test.³² NMR spectra were recorded on a 400 MHz spectrometer for ^1H NMR spectra δ values were recorded as follows: CDCl_3 (7.27 ppm); CD_3OD (3.31; 4.87 ppm), D_2O (4.75 ppm) for ^{13}C (125 MHz) δ CDCl_3 (77.0 ppm); CD_3OD (49.2 ppm). Mass spectra (MS) were obtained using electron impact (EI) or electrospray ionisation (ESI). Chemical exchange saturation transfer (CEST) spectra were acquired using a 9.4 T small animal clinical MRI scanner as follows: NMR tubes with solutions of the complexes (15 mM, pH 7.5) were imaged at 37 °C [the temperature was monitored and controlled by blowing hot air using a Model 1025 Small Animal Monitoring and Gating System (SA Instruments, Inc., Stony Brook, NY)] using a fast spin echo pulse sequence (FOV: $12.8 \times 12.8 \text{ mm}^2$, matrix: 32×32 , TR = 4000 ms, 4 echoes, and TE = 10 ms), preceded by a frequency selective saturation pulse ($B_1 = 14 \mu\text{T}$, saturation range = -150 to 50 ppm in steps of 1 ppm, saturation time = 3.95 s). CEST spectra were generated using the average signal intensity from each tube. Similar methodology was used to evaluate the temperature (15 mM, pH 7.5, temperature range 33-39 °C) and pH (15 mM, 37 °C, pH range 6.0-8.0) sensitivity of selected agents. The exchange rates of the amide protons with bulk water were measured by Ω -plot method as described previously.³³

Preparation of (S)-N-benzyl-2-chloropropanamide

The reaction was carried out as described in the literature,³⁴ starting from 430 μL (5 mmol) of S-chloropropionic acid (**12**). The product was isolated by FCC on 70 g SiO_2 , eluting with hexanes/EtOAc (3:1), obtained (S)-N-benzyl-2-chloropropanamide (**13**, 813 mg, 82%), colorless solid. $[\alpha]_D -1.3$ (c 1, MeOH); lit.³⁴ $[\alpha]_D -3.8$ (c 1, MeOH). ^1H NMR (CDCl_3) δ 7.37 (m, 2H); 7.31 (m, 3H); 6.88 (br s, D_2O exch, 1H); 4.49 (m,

3H); 1.79 (d, $J = 7.0 \text{ Hz}$, 3H). ^{13}C NMR (CDCl_3) δ 169.4, 137.4, 128.6, 127.5 (2 \times C), 55.6, 43.7, 22.5. HRMS (EI) m/z ; found 197.0602 $[\text{M}]^+$ (calcd 197.0607 for $\text{C}_{10}\text{H}_{12}\text{ClNO}$); LRMS (EI) m/z (rel. abundance): 197 $[\text{M}]^+$ (23), 162 (85), 106 (14).

Coupling of (S)-lactic acid with aniline, benzylamine and *t*-butylamine

Separate mixtures containing (S)-lactic acid (**14**, 450 mg, 5 mmol for benzylamine or 180 mg, 2 mmol), HOBt (1.35 g, 10 mmol for benzylamine or 540 mg, 4 mmol), DCC (1.55 g, 7.5 mmol for benzylamine or 619 mg, 3 mmol) and benzylamine (660 μL , 6 mmol), aniline (220 μL , 2.4 mmol) or *t*-butylamine (220 μL , 2.4 mmol) in dry CH_2Cl_2 (15 mL for benzylamine or 6 mL) were stirred for 24 h at room temperature (rt). The solids were filtered off, the filters were washed with ice cold CH_2Cl_2 . Resulting solutions were washed with saturated NaHCO_3 solution (30 mL for benzylamine or 20 mL), the aqueous layers were extracted with CH_2Cl_2 (2 \times 30 mL for benzylamine or 2 \times 20 mL). Combined organic extracts were dried, were concentrated, the residues were dissolved in small amount of cold acetone (ca. 5 – 10 mL) and precipitated dicyclohexylurea was filtered off using a Pasteur pipette with a cotton plug. The volatiles were evaporated and the residues were subjected to FCC on 40 g SiO_2 , eluting with hexanes/EtOAc (3:1, reaction with benzylamine); 30 g SiO_2 , eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (98:2, reaction with aniline) or 25 g SiO_2 , eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (95:5). Evaporation of the eluates afforded (S)-N-benzyl lactamide (**15**, 727 mg, 81%), (S)-N-*t*-butyl lactamide (**16**, 175 mg, 60%) or (S)-N-phenyl lactamide (**17**, 266 mg, 81%).

(S)-N-benzyl lactamide (**15**), colorless oil. $[\alpha]_D -6.4$ (c 1, MeOH); lit.³⁵ $[\alpha]_D -6.8$ (c 6.9, CHCl_3). ^1H NMR (CDCl_3) δ 7.29 (m, 5H); 7.11 (br s, D_2O exch, 1H); 4.41 (d, $J = 6.0 \text{ Hz}$, 2H); 4.24 (q, $J = 7.0 \text{ Hz}$, 1H); 1.42 (d, $J = 7.0 \text{ Hz}$, 3H). ^{13}C NMR (CDCl_3) δ 174.7, 137.8, 128.7, 127.6, 127.5, 68.3, 43.0, 21.2. HRMS (EI) m/z ; found 179.0952 $[\text{M}]^+$ (calcd 179.0946 for $\text{C}_{10}\text{H}_{13}\text{NO}_2$); LRMS (EI) m/z (rel. abundance): 179 $[\text{M}]^+$ (33), 161 (15), 134 (10), 91 (100), 65 (12).

(S)-N-*t*-butyl lactamide (**16**), colorless solid. $[\alpha]_D -28.0$ (c 1, MeOH). ^1H NMR (CDCl_3) δ 6.47 (br s, D_2O exch, 1H); 4.09 (q, $J = 6.5 \text{ Hz}$, 1H); 3.52 (br s, D_2O exch., 1H); 1.38 (d, $J = 6.5 \text{ Hz}$, 3H); 1.36 (s, 9H). ^{13}C NMR (CDCl_3) δ 174.5, 68.3, 50.7, 28.6, 21.1. HRMS (EI) m/z ; found 145.1107 $[\text{M}]^+$ (calcd 145.1103 for $\text{C}_7\text{H}_{15}\text{NO}_2$); LRMS (EI) m/z (rel. abundance): 145 $[\text{M}]^+$ (95), 130 (100), 101 (67).

(S)-N-phenyl lactamide (**17**), colorless oil. $[\alpha]_D -41.0$ (c 0.5, MeOH); lit.³⁶ $[\alpha]_D -23.5$ (c 0.48, CHCl_3). ^1H NMR (CDCl_3) δ 8.67 (br s, D_2O exch., 1H); 7.52 (m, 2H); 7.30 (m, 2H); 7.12 (m, 1H); 4.31 (q, $J = 6.5 \text{ Hz}$, 1H); 4.14 (br s, D_2O exch, 1H); 1.48 (d, $J = 6.5 \text{ Hz}$, 3H). ^{13}C NMR (CDCl_3) δ 173.2, 137.0, 129.0, 124.6, 119.9, 68.7, 21.0. HRMS (EI) m/z ; found 165.0797 $[\text{M}]^+$ (calcd 165.0790 for $\text{C}_9\text{H}_{11}\text{NO}_2$); LRMS (EI) m/z (rel. abundance): 165 $[\text{M}]^+$ (50), 121 (21), 93 (100), 65 (13).

Coupling of (S)-lactic acid with H-(L)-Lys(Boc)-OMe•HCl

ARTICLE

Journal Name

A stirred mixture containing (S)-lactic acid (**14**, 180 mg, 2 mmol), *H*-(L)-Lys(Boc)-OMe · HCl (594 mg, 2 mmol), HOBt (270 mg, 2 mmol) and Et₃N (280 µL, 2 mmol) in dry CH₂Cl₂ (15 mL) was cooled to 0 °C, followed by an addition of DCC (825 mg, 4 mmol). The stirring continued for further 30 min at 0 °C, the cooling bath was removed and the stirring was continued for further 48 h at rt. The solids were filtered off, the filter was washed with ice cold CH₂Cl₂. Resulting solution was washed with saturated NaHCO₃ solution (30 mL), the aqueous layer was extracted with CH₂Cl₂ (30 mL). Combined organic extract was dried, was concentrated, the residue was dissolved in small amount of cold acetone (ca. 5 – 10 mL) and precipitated dicyclohexylurea was filtered off using a Pasteur pipette with a cotton plug. The volatiles were evaporated and the residue was subjected to FCC on 50 g SiO₂, eluting with CH₂Cl₂/MeOH (95:5). Evaporation of the eluate afforded (S)-*N*-(L)-Lys(Boc)-OMe lactamide (**18**, 621 mg, 93%) as colorless oil. [α]_D -11.9 (c 1.4, MeOH). ¹H NMR (CDCl₃) δ 7.20 (d, D₂O exch., *J* = 8.5 Hz, 1H); 4.71 (br s, D₂O exch, 1H); 4.57 (m, 1H); 4.24 (q, *J* = 6.5 Hz, 1H); 3.73 (s, 3H); 3.08 (m, 2H); 1.86 (m, 1H); 1.71 (m, 1H); 1.49 (m, 2H); 1.43 (m, 12H); 1.35 (m, 2H). ¹³C NMR (CDCl₃) δ 174.8, 172.8, 156.1, 79.2, 68.3, 52.4, 51.5, 40.1, 31.9, 29.4, 28.4, 22.4, 21.0. HRMS (ESI) *m/z*: found 333.2025 [M + H]⁺ (calcd. 333.2026 for C₁₅H₂₉N₂O₆).

Reaction of (S)-*N*-*t*-butyl lactamide with Tf₂O

A round bottom flask containing (S)-*N*-*t*-butyl lactamide (**16**, 202 mg, 1.39 mmol) was flushed with N₂, followed by the addition of dry CH₂Cl₂ (2 mL) and dry pyridine (120 µL, 1.46 mmol). The solution was cooled to 0 °C, followed by a dropwise addition (over ca. 1 min) of Tf₂O (230 µL, 1.39 mmol). The stirring continued for further 1 h at 0 °C, the solvent was evaporated and the residue was subjected to FCC on 25 g SiO₂, eluting with hexanes/EtOAc (1:1). Evaporation of the eluate afforded (S)-*O*-Tf-*N*-*t*-butyl lactamide (**16b**, 223 mg, 34%) as colorless solid. [α]_D -16.8 (c 0.5, MeOH). ¹H NMR (CDCl₃) δ 5.94 (br s, D₂O exch, 1H); 5.14 (q, *J* = 7.0 Hz, 1H); 1.70 (d, *J* = 7.0 Hz, 3H); 1.38 (s, 9H). ¹³C NMR (CDCl₃) δ 166.0, 123.1, 120.0, 116.8, 113.6, 83.4, 52.1, 28.4, 19.1. ¹⁹F NMR (CDCl₃) δ -74.9. HRMS (EI) *m/z*: found 277.0591 [M]⁺ (calcd 277.0596 for C₈H₁₄F₃NO₄S); LRMS (EI) *m/z* (rel. abundance): 277 [M]⁺ (16), 262 (100), 222 (44), 177 (20), 112 (42), 69 (90).

Reaction of (S)-*N*-benzyl lactamide with TsCl

A solution of (S)-*N*-benzyl lactamide (**15**, 295 mg, 1.65 mmol) in dry CH₂Cl₂ (6 mL) was cooled to 0 °C, followed by the addition of Et₃N (460 µL, 3.29 mmol), DMAP (24 mg, 0.2 mmol) and TsCl (471 mg, 2.47 mmol). The mixture was stirred for 1 h at 0 °C, then for 24 h at rt. The solvent was evaporated and the residue was subjected to FCC on 30 g SiO₂, eluting with hexanes/EtOAc (3:1). Evaporation of the eluate afforded (S)-*O*-Ts-*N*-benzyl lactamide (**15c**, 450 mg, 82%) as colorless solid. [α]_D -53.9 (c 1, MeOH). ¹H NMR (CDCl₃) δ 7.77 (d, *J* = 8.0 Hz, 2H); 7.33 (m, 5H); 7.20 (d, *J* = 6.5 Hz, 2H); 6.62 (br s, D₂O exch, 1H); 4.92 (q, *J* = 7.0 Hz, 1H); 4.41 (d, *J* = 6.0 Hz, 2H); 2.46 (s, 3H); 1.47 (d, *J* = 7.0 Hz, 3H). ¹³C NMR (CDCl₃) δ 168.6, 145.4, 137.3, 132.6, 130.0, 128.6, 127.7, 127.4, 77.3, 43.1, 21.6, 18.7.

HRMS (EI) *m/z*: found 179.0952 [M]⁺ (calcd 179.0946 for C₁₀H₁₃NO₂); LRMS (EI) *m/z* (rel. abundance): 179 [M]⁺ (33), 161 (15), 134 (10), 91 (100), 65 (12). HRMS (ESI) *m/z*: found 334.1124 [M + H]⁺ (calcd. 334.1113 for C₁₇H₂₀NO₄S).

Reaction of (S)-lactamides 15-19 with MsCl

Separate suspensions or solutions of (S)-*N*-benzyl lactamide (**15**, 788 mg, 4.4 mmol), (S)-*N*-*t*-butyl lactamide (**16**, 175 mg, 1.21 mmol), (S)-*N*-phenyl lactamide (**17**, 266 mg, 1.61 mmol), (S)-*N*-(L)-Lys(Boc)-OMe lactamide (**18**, 279 mg, 0.84 mmol) or (S)-lactamide (**19**, 891 mg, 10 mmol) in dry CH₂Cl₂ [20 mL in the case of (S)-*N*-benzyl lactamide (**15**); 6 mL in the case of (S)-*N*-*t*-butyl lactamide (**16**); 8 mL in the case of (S)-*N*-phenyl lactamide (**17**); 4 mL in the case of (S)-*N*-(L)-Lys(Boc)-OMe lactamide (**18**); 15 mL in the case of (S)-lactamide (**19**)] and Et₃N [1.12 mL, 8 mmol in the case of (S)-*N*-benzyl lactamide (**15**); 310 µL, 2.19 mmol in the case of (S)-*N*-*t*-butyl lactamide (**16**); 410 µL, 2.93 mmol in the case of (S)-*N*-phenyl lactamide (**17**); 580 µL, 4.2 mmol in the case of (S)-*N*-(L)-Lys(Boc)-OMe lactamide (**18**); 2.54 mL, 18.2 mmol in the case of (S)-lactamide (**19**)] were cooled to 0 °C, followed by a dropwise addition (over ca. 1 min) of MsCl [430 µL, 5.58 mmol in the case of (S)-*N*-benzyl lactamide (**15**); 120 µL, 1.53 mmol in the case of (S)-*N*-*t*-butyl lactamide (**16**); 160 µL, 2.04 mmol in the case of (S)-*N*-phenyl lactamide (**17**); 90 µL, 1.07 mmol in the case of (S)-*N*-(L)-Lys(Boc)-OMe lactamide (**18**); 980 µL, 12.7 mmol in the case of (S)-lactamide (**19**)]. The cooling baths were removed and the mixtures were stirred at rt as follows: 2 h in the case of (S)-*N*-benzyl lactamide (**15**) and (S)-*N*-(L)-Lys(Boc)-OMe lactamide (**18**); 4 h in the case of (S)-*N*-*t*-butyl lactamide (**16**) and (S)-*N*-phenyl lactamide (**17**); 24 h in the case of (S)-lactamide (**19**). The volatiles were evaporated and the residues were subjected to FCC as follows: 40 g SiO₂, eluting with hexanes/EtOAc (1:1) in the case of (S)-*N*-benzyl lactamide (**15**) and (S)-*N*-phenyl lactamide (**17**); 30 g SiO₂, eluting with hexanes/EtOAc (1:1) in the case of (S)-*N*-*t*-butyl lactamide (**16**); 30 g SiO₂, eluting with CH₂Cl₂/MeOH (95:5) in the case of (S)-*N*-(L)-Lys(Boc)-OMe lactamide (**18**); 60 g SiO₂, eluting with CH₂Cl₂/MeOH (95:5) in the case of (S)-lactamide (**19**). Evaporation of the eluates afforded (S)-*O*-Ms-*N*-benzyl lactamide (**15a**, 1.05 g, 93%), (S)-*O*-Ms-*N*-*t*-butyl lactamide (**16a**, 253 mg, 94%), (S)-*O*-Ms-*N*-phenyl lactamide (**17a**, 373 mg, 95%) and (S)-*O*-Ms-*N*-(L)-Lys(Boc)-OMe lactamide (**18a**, 322 mg, 93%). In the case of (S)-lactamide (**19**) was the eluate concentrated to ca. one quarter of its original volume (a precipitate started to form), hexanes were added and the mixture was set aside for 2 h at -10 °C. Separated precipitate was filtered off, was washed with hexanes and was dried to afford (S)-*O*-Ms-lactamide (**19a**, 1.34 g, 80%).

(S)-*O*-Ms-*N*-benzyl lactamide (**15a**), colorless solid. [α]_D -39.8 (c 1, MeOH). ¹H NMR (CDCl₃) δ 7.34 (m, 5H); 6.64 (br s, D₂O exch, 1H); 5.14 (q, *J* = 7.0 Hz, 1H); 4.53 (dd, *J* = 15.0, 6.0 Hz, 1H); 4.47 (dd, *J* = 15.0, 6.0 Hz, 1H); 3.08 (s, 3H); 1.67 (d, *J* = 7.0 Hz, 3H). ¹³C NMR (CDCl₃) δ 168.7, 137.4, 128.6, 127.5, 127.4, 76.5, 43.2, 38.4, 18.9. HRMS (EI) *m/z*: found 258.0806 [M + H]⁺ (calcd 258.0800 for C₁₁H₁₆NO₄S); LRMS (EI) *m/z* (rel.

abundance): 258 [M⁺] (10), 161 (62), 133 (23), 106 (44), 91 (100).

(*S*)-*O*-Ms-*N*-*t*-butyl lactamide (**16a**), colorless solid. [α]_D -68.8 (c 0.5, MeOH). ¹H NMR (CDCl₃) δ 6.09 (br s, D₂O exch., 1H); 4.95 (q, *J* = 7.0 Hz, 1H); 3.10 (s, 3H); 1.60 (d, *J* = 7.0 Hz, 3H); 1.38 (s, 9H). ¹³C NMR (CDCl₃) δ 167.7, 51.6, 38.8, 28.5, 18.8. HRMS (EI) *m/z*; found 223.0874 [M]⁺ (calcd 223.0878 for C₈H₁₇NO₄S); LRMS (EI) *m/z* (rel. abundance): 223 [M⁺] (29), 208 (100), 180 (16), 136 (15), 123 (50), 84 (79).

(*S*)-*O*-Ms-*N*-phenyl lactamide (**17a**), colorless solid. [α]_D -64.6 (c 1, MeOH). ¹H NMR (CDCl₃) δ 8.13 (br s, D₂O exch., 1H); 7.55 (m, 2H); 7.35 (m, 2H); 7.17 (m, 1H); 5.21 (q, *J* = 7.0 Hz, 1H); 3.16 (s, 3H); 1.72 (d, *J* = 7.0 Hz, 3H). ¹³C NMR (CDCl₃) δ 166.6, 136.6, 129.1, 125.2, 120.2, 76.6, 38.9, 18.8. HRMS (EI) *m/z*; found 243.0562 [M]⁺ (calcd 243.0565 for C₁₀H₁₃NO₄S); LRMS (EI) *m/z* (rel. abundance): 243 [M⁺] (83), 120 (100), 92 (43).

(*S*)-*O*-Ms-*N*-(*L*)-Lys(Boc)-OMe lactamide (**18a**), pale yellow oil. [α]_D -36.5 (c 1, MeOH). ¹H NMR (CDCl₃) δ 6.89 (d, D₂O exch., *J* = 8.0 Hz, 1H); 5.08 (q, *J* = 7.0 Hz, 1H); 4.63 (br s, D₂O exch., 1H); 4.58 (m, 1H); 3.74 (s, 3H); 3.14 (s, 3H); 3.10 (m, 2H); 1.90 (m, 1H); 1.72 (m, 1H); 1.63 (d, *J* = 7.0 Hz, 3H); 1.49 (m, 2H); 1.42 (s, 9H); 1.35 (m, 2H). ¹³C NMR (CDCl₃) δ 172.2, 168.6, 156.0, 79.1, 76.3, 52.5, 51.8, 40.0, 38.7, 31.8, 29.4, 28.4, 22.3, 19.2. HRMS (ESI) *m/z*; found 411.1786 [M + H]⁺ (calcd. 411.1801 for C₁₆H₃₁N₂O₈S).

(*S*)-*O*-Ms-lactamide (**19a**), colorless crystals; mp 110–112 °C. [α]_D -50.9 (c 1, MeOH). ¹H NMR (CD₃OD) δ 5.02 (q, *J* = 7.0 Hz, 1H); 3.16 (s, 3H); 1.57 (d, *J* = 7.0 Hz, 3H). ¹³C NMR (CD₃OD) δ 174.6, 77.0, 38.6, 19.6. HRMS (EI) *m/z*; found 167.0247 [M]⁺ (calcd 167.0252 for C₄H₉NO₄S); LRMS (EI) *m/z* (rel. abundance): 167 [M⁺] (10), 123 (56), 109 (12), 65 (29).

Alkylation of cyclen with (*S*)-*O*-Ms-lactamides **15a**–**19a**

K₂CO₃ [320 mg, 2.31 mmol or 767 mg, 5.55 mmol in the case of (*S*)-*O*-Ms-lactamide (**19a**)] was added to separate solutions of cyclen [**5**, 43 mg, 0.25 mmol or 103 mg, 0.6 mmol in the case of (*S*)-*O*-Ms-lactamide (**19a**)] and (*S*)-*O*-Ms-*N*-benzyl lactamide (**15a**, 257 mg, 1 mmol), (*S*)-*O*-Ms-*N*-*t*-butyl lactamide (**16a**, 223 mg, 1 mmol), (*S*)-*O*-Ms-*N*-phenyl lactamide (**17a**, 243 mg, 1 mmol) or (*S*)-*O*-Ms-lactamide (**19a**, 401 mg, 2.4 mmol) in MeCN [1.5 mL or 5 mL in the case of (*S*)-*O*-Ms-lactamide (**19a**)]. The mixtures were stirred for 48 h at 60 °C, were cooled to rt, were diluted with brine (40 mL) and were extracted with EtOAc (2 × 20 mL). In the case of (*S*)-*O*-Ms-lactamide (**19a**) the solids were filtered off with suction and the filtrate was washed with MeCN. Combined organic extracts were dried and were concentrated along with the filtrate obtained in the case of (*S*)-*O*-Ms-lactamide (**19**). The residues obtained after extraction were dissolved in MeOH (3 mL) and were subjected to semi-preparative HPLC purification as described in General experimental procedures. The fractions containing the desired product were combined and concentrated to leave DOTMAM-Bn × 4CF₃COO[−] (115 mg,

42%), DOTMAM-*t*-Bu × 4CF₃COO[−] (67 mg, 24%) or DOTMAM-Ph × 4CF₃COO[−] (43 mg, 14%). The residue obtained in the case of (*S*)-*O*-Ms-lactamide (**19a**) was subjected to FCC on 25 g Al₂O₃, eluting first with CH₂Cl₂/MeOH/NH₄OH (80:19:1, 3 fractions ca. 30 mL each) later with MeOH (8–10 fractions, ca. 30 mL each). The later fractions were concentrated to give DOTMAM (141 mg, 51%).

DOTMAM-Bn × 4CF₃COO[−], colorless hygroscopic solid. HPLC, Method B, *t*_R 9.1 min. ¹H NMR (D₂O) δ 6.82–6.71 (br m, 20H); 4.17–2.47 (br m, 28H); 1.48 (br s, 6H); 0.92 (br s, 6H). HRMS (ESI) *m/z*; found 817.5093 [M + H]⁺ (calcd. 817.5129 for C₄₈H₆₅N₈O₄).

DOTMAM-*t*-Bu × 4CF₃COO[−], colorless hygroscopic solid. HPLC, Method C, *t*_R 7.9 min. ¹H NMR (D₂O) δ 4.12–2.60 (br m, 20H); 1.49 (br s, 6H); 1.22 (br s, 36H); 1.08 (br s, 6H). HRMS (ESI) *m/z*; found 681.5723 [M + H]⁺ (calcd. 681.5755 for C₃₆H₇₃N₈O₄).

DOTMAM-Ph × 4CF₃COO[−], colorless hygroscopic solid. HPLC, Method D, *t*_R 7.8 min. ¹H NMR (D₂O) δ 7.32–6.88 (br m, 20H); 4.74–2.66 (br m, 20H); 1.68–1.17 (br m, 12H). HRMS (ESI) *m/z*; found 761.4512 [M + H]⁺ (calcd. 761.4503 for C₄₄H₅₇N₈O₄).

DOTMAM, colorless solid. ¹H NMR (CD₃OD) δ 3.63 (q, *J* = 7.0 Hz, 4H); 3.00 (t, *J* = 13.5 Hz, 4H); 2.82 (t, *J* = 12.5 Hz, 4H); 2.38 (d, *J* = 14.0 Hz, 4H); 2.15 (d, *J* = 13.5 Hz, 4H); 1.19 (d, *J* = 7.0 Hz, 12H). ¹³C NMR (CD₃OD) δ 179.5, 57.3, 47.7, 46.2, 7.6. HRMS (ESI) *m/z*; found 457.3230 [M + H]⁺ (calcd. 457.3251 for C₂₀H₄₁N₈O₄).

Coupling of DOTMA · 4Na⁺ with *H*-(*L*)-Lys(Boc)-OMe · HCl and *H*-(*L*)-Phe-OEt · HCl

A stirred separate suspensions of DOTMA×4Na⁺ (**4**, 82 mg, 0.15 mmol), prepared according to previously described procedures^{37,38} and *H*-(*L*)-Lys(Boc)-OMe·HCl (178 mg, 0.6 mmol) or *H*-(*L*)-Phe-OEt·HCl (138 mg, 0.6 mmol) in DIPEA (210 μL, 1.2 mmol) and dry DMF (1 mL) were cooled to 0 °C, followed by the addition of HBTU (228 mg, 0.6 mmol). The mixtures were stirred for 10 min at 0 °C, then for 24 h at 60 °C, then they were cooled to rt and were diluted with brine (50 mL), followed by the extraction with EtOAc (2 × 25 mL). Combined organic extracts were washed with brine (3 × 50 mL), were dried and were concentrated. The residues were dissolved in MeOH (3 mL) and were subjected to semi-preparative HPLC purification as described in General experimental considerations. The fractions containing the desired product were combined and concentrated to leave DOTMAM-(*L*)-Lys(Boc)-OMe × 4CF₃COO[−] (117 mg, 41%) or DOTMAM-(*L*)-Phe-OEt × 4CF₃COO[−] (195 mg, 81%).

DOTMAM-(*L*)-Lys(Boc)-OMe × 4CF₃COO[−], colorless solid. HPLC, Method E, *t*_R 7.5 min. ¹H NMR (CD₃OD) δ 4.78–3.94 (br m, 4H); 3.76 (br s, 12H); 3.47–2.55 (br m, 24H); 1.86–1.34 (br m, 72H). HRMS (ESI) *m/z*; found 1429.9065 [M + H]⁺ (calcd. 1429.9133 for C₆₈H₁₂₅N₁₂O₂₀).

DOTMAM-(L)-Phe-OEt \times 4CF₃COO⁻, pale brown solid. HPLC, Method E, *t*_R 7.8 min. ¹H NMR (CD₃OD) δ 7.31-7.02 (br m, 20 H); 5.06 (br m, 4H); 4.46-4.12 (br m, 12H); 3.41-2.31 (br m, 24H); 1.48 (br s, 6H); 1.35 (m, 12H); 1.08 (br s, 6H). HRMS (ESI) *m/z*: found 1161.6594 [M + H]⁺ (calcd. 1161.6600 for C₆₄H₈₉N₈O₁₂).

Deprotection of DOTMAM-(L)-Lys(Boc)-OMe

A solution of DOTMAM-(L)-Lys(Boc)-OMe \times 4CF₃COO⁻ (117 mg, 0.062 mmol) in TFA (1 mL) was stirred for 20 min at rt. The volatiles were evaporated, the residue was dissolved in MeOH (600 μ L), followed by the addition of NaOH solution (2.5 M, 1 mL, 2.5 mmol). Resulting mixture was stirred for 2 h at 60 $^{\circ}$ C, MeOH was evaporated, the mixture was cooled to 0 $^{\circ}$ C and the pH was adjusted to ca. 7 (1 M HCl). Resulting aqueous solution was subjected to SEC as described in General experimental procedures. The fractions containing the product were combined and concentrated to leave DOTMAM-(L)-Lys-OH (50 mg, 83%) as colorless solid. ¹H NMR (D₂O) δ 4.09 (m, 4H); 3.78 (m, 4H); 3.09-2.88 (br m, 24H); 1.76-1.30 (br m, 36H). HRMS (ESI) *m/z*: found 973.6386 [M + H]⁺ (calcd. 973.6410 for C₄₄H₈₅N₁₂O₁₂).

Saponification of DOTMAM-(L)-Phe-OEt

A solution of NaOH (64 mg, 1.6 mmol) in H₂O (1 mL) was added to a solution of DOTMAM-(L)-Phe-OEt \times 4CF₃COO⁻ (117 mg, 0.1 mmol) in THF (1 mL). The mixture was vigorously stirred for 24 h at 60 $^{\circ}$ C, was diluted with small amount of H₂O (ca. 3 mL) and was cooled to 0 $^{\circ}$ C. The pH was adjusted to ca. 4-5 (1 M HCl), the mixture was set aside for 4 h at 3 $^{\circ}$ C, the precipitate was filtered off with suction, was washed with water and was dried to leave DOTMAM-(L)-Phe-OH (57 mg, 54%) as pale brown solid. ¹H NMR (CD₃OD) δ 7.19 (m, 20 H); 4.71 (br m, 4H); 3.58-2.82 (br m, 28H); 1.14 (br m, 12H). HRMS (ESI) *m/z*: found 1049.5399 [M + H]⁺ (calcd. 1049.5348 for C₅₆H₇₃N₈O₁₂).

Metallation of DOTMAM, DOTMAM-Bn, DOTMAM-*t*-Bu, DOTMAM-(L)-Lys-OH and DOTMAM-(L)-Phe-OH with TmCl₃•H₂O

Separate solutions of TmCl₃•H₂O [34 mg, 0.13 mmol in the case of DOTMAM; 13 mg, 0.047 mmol in the case of DOTMAM-Bn; 6 mg, 0.021 mmol in the case of DOTMAM-*t*-Bu; 18 mg, 0.066 mmol in the case of DOTMAM-(L)-Lys-OH and DOTMAM-(L)-Phe-OH] in water [2 mL in the case of DOTMAM, DOTMAM-(L)-Lys-OH and DOTMAM-(L)-Phe-OH; 2.5 mL in the case of DOTMAM-Bn; 1 mL in the case of DOTMAM-*t*-Bu] were added to separate solutions of DOTMAM (46 mg, 0.1 mmol), DOTMAM-Bn \times 4CF₃COO⁻ (54 mg, 0.043 mmol), DOTMAM-*t*-Bu \times 4CF₃COO⁻ (22 mg, 0.019 mmol), DOTMAM-(L)-Lys-OH \times 4CF₃COO⁻ (54 mg, 0.055 mmol) and DOTMAM-(L)-Phe-OH \times 4CF₃COO⁻ (57 mg, 0.055 mmol) in dioxane [2 mL in the case of DOTMAM, DOTMAM-(L)-Lys-OH and DOTMAM-(L)-Phe-OH; 2.5 mL in the case of DOTMAM-Bn; 1 mL in the case of DOTMAM-*t*-Bu]. The mixtures were stirred for 48 h at 90 $^{\circ}$ C, while the pH was maintained at ca. 6.5-7.0 (1 M NaOH solution). Reaction mixtures were transferred into centrifuge tubes and were lyophilized. The residues obtained from DOTMAM-Bn and

DOTMAM-*t*-Bu were dissolved in MeOH/water (1.5 mL each) and were subjected to semipreparative HPLC purification as described in General experimental procedures. The fractions containing the desired product were combined, were concentrated, were dissolved in water (1 mL) and were neutralized (pH ca. 7.0, 1 M NaOH solution). Resulting solutions were subjected to SEC as described in General experimental procedures. The fractions containing the desired product were combined and concentrated to leave Tm³⁺ DOTMAM-Bn (**7**, 21 mg, 50%) and Tm³⁺ DOTMAM-*t*-Bu (**8**, 7 mg, 40%). The residues obtained from DOTMAM, DOTMAM-(L)-Lys-OH and DOTMAM-(L)-Phe-OH were dissolved in water (2 mL) and were subjected to SEC as described in General experimental procedures. The fractions containing the desired product were combined and concentrated to leave Tm³⁺ DOTMAM (**6**, 32 mg, 49%), Tm³⁺ DOTMAM-(L)-Lys-OH (**10**, 49 mg, 78%) and Tm³⁺ DOTMAM-(L)-Phe-OH (**11**, 38 mg, 58%).

Tm³⁺ DOTMAM (**6**), colorless solid. HRMS (ESI) *m/z*: found 623.2344 [M - 2H]⁺ (calcd. 623.2358 for C₂₀H₃₈N₈O₄Tm).

Tm³⁺ DOTMAM-Bn (**7**), colorless solid. HPLC, Method C, *t*_R 6.1 min. HRMS (ESI) *m/z*: found 983.4254 [M - 2H]⁺ (calcd. 983.4236 for C₄₈H₆₂N₈O₄Tm).

Tm³⁺ DOTMAM-*t*-Bu (**8**), colorless solid. HPLC, Method F, *t*_R 6.6 min. HRMS (ESI) *m/z*: found 847.4824 [M - 2H]⁺ (calcd. 847.4862 for C₃₆H₇₀N₈O₄Tm).

Tm³⁺ DOTMAM-(L)-Lys-OH (**10**), colorless solid. HRMS (ESI) *m/z*: found 1139.5552 [M - 2H]⁺ (calcd. 1139.5523 for C₄₄H₈₂N₁₂O₁₂Tm).

Tm³⁺ DOTMAM-(L)-Phe-OH (**11**), colorless solid. HRMS (ESI) *m/z*: found 1237.4275 [M - 3H + Na]⁺ (calcd. 1237.4239 for C₅₆H₆₉N₈O₁₂NaTm).

Conclusions

In summary, we have developed a new synthetic methodology and prepared the first examples of DOTMAM ligands possessing secondary amides including amino acids. This was achieved by tetraalkylation of cyclen with OMs-lactamides as a key step for the preparation of simple alkyl- and aryl-decorated ligands, while peptide coupling of DOTMA tetrasodium salt with protected amino acids furnished amino acid-decorated DOTMAM ligands. Furthermore, the Tm³⁺ complexes of these ligands were prepared and their CEST properties were investigated. Although the CEST effects were more highly shifted than comparable Tm³⁺DOTAM-based complexes, the intensity of the CEST effects were observed to be weaker. This has been ascribed to less favourable (slower) amide proton exchange rates. The L-lysine conjugated agent **10** displayed temperature- and pH-dependent CEST effects in the physiological range. Based on similarity to the analogous DOTAM-based agent, it is expected to have reasonable biocompatibility³⁹ and may be suitable for future *in vivo*

studies.^{9,39} Future work should investigate Eu³⁺ DOTMAM-alkyl decorated complexes as well as a wider selection of Tm³⁺ DOTMAM-amino acid decorated complexes for useful CEST properties.

Examination of the Yb³⁺ complexes for DOTMAM, DOTMAM-Bn and DOTMAM-*t*-Bu indicated an overwhelming predominance of a single coordination isomer in solution although the *t*-butyl amide appeared different than the other two. The Tm³⁺ complexes of these ligands also show the greatest difference in the chemical shift of the amide-proton CEST signal. On the basis of the cyclen H₄ proton chemical shift we speculate that the TSAP isomer is most favoured in this series by the combination of the *t*-butyl amide substituent and DOTMAM ligand, although further studies are needed to elucidate the geometry unambiguously.

Acknowledgements

Financial support from the Canadian Institutes of Health Research (CIHR) and the Natural Sciences and Engineering Research Council of Canada (NSERC) is gratefully acknowledged. We thank Prof. J. M. Chong (University of Waterloo) for the assistance with optical rotation measurements.

Notes and references

‡ DOTAM refers to the tetraamide derivative of DOTA (1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid).

§ The term DOTMA is used to refer to the *RRRR*-enantiomer of α , α' , α'' , α''' -tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetate in this manuscript.

§§ The term DOTMAM is used to denote DOTMA-based amides in analogy to DOTAM which is used to denote DOTA-based amides. This abbreviation was first used by Merbach and coworkers with respect to the primary amide derivative of DOTMA. See reference 19.

† The outcome of FCC purification depended strongly on quality of Al₂O₃, the best results were obtained with 10-100 mesh, pH 9.5-10.5.

‡ The coupling between DOTMA•4Na⁺ and benzylamine (7% yield) was found to be inferior to that obtained by tetraalkylation of cyclen with OM_s-(*S*)-benzyl lactamide.

‡ The CEST effect associated with CA 11 was too weak to measure the exchange rate.

¹Boros, E.; Gale, E. M.; Caravan, P. *Dalton Trans.* **2015**, 44, 4804-4818.

²(a) Yoo, B.; Pagel, M. D. *Front. Biosci.* **2008**, 13, 1733-1752. (b) Geraldine, C. F. G. C.; Laurent, S. *Contrast Med. Mol. Imaging* **2009**, 4, 1-23.

³(a) Zhang, S.; Merrit, M.; Woessner, D. E.; Lenkinski, R. E.; Sherry, A. D. *Acc. Chem. Res.* **2003**, 36, 783-790. (b) Viswanathan, S.; Kovacs, Z.; Green, K. N.; Ratnakar, S. J.; Sherry, A. D. *Chem. Rev.* **2010**, 110, 2960-3018.

⁴Li, A. X.; Wojciechowski, F.; Suchý, M.; Jones, C. K.; Hudson, R. H. E.; Menon, R. S.; Bartha, R. *Magn. Reson. Med.* **2008**, 59, 374-381.

⁵Suchý, M.; Li, A. X.; Milne, M.; Bartha, R.; Hudson, R. H. E. *Contrast Med. Mol. Imaging* **2012**, 7, 441-449.

⁶Ratnakar, S. J.; Viswanathan, S.; Kovacs, Z.; Jindal, A. K.; Green, K. N.; Sherry, A. D. *J. Am. Chem. Soc.* **2012**, 134, 5798-5800.

⁷Zhang, S.; Trokowski, R.; Sherry, A. D. *J. Am. Chem. Soc.* **2003**, 125, 15288-15289.

⁸Yoo, B.; Pagel, M. D. *J. Am. Chem. Soc.* **2006**, 128, 14032-14033.

⁹(a) Jones, C. K.; Li, A. X.; Suchý, M.; Hudson, R. H. E.; Menon, R.; Bartha, R. *Magn. Reson. Med.* **2010**, 63, 1184-1192. (b) Li, A. X.; Suchý, M.; Li, C.; Gati, J. S.; Meakin, S.; Hudson, R. H. E.; Menon, R.; Bartha, R. *Magn. Reson. Med.* **2011**, 66, 67-72. (c) McVicar, N.; Li, A. X.; Suchý, M.; Hudson, R. H. E.; Menon, R.; Bartha, R. *Magn. Reson. Med.* **2013**, 70, 1016-1025.

¹⁰E. Terreno, D. Delli Castelli, A. Viale and S. Aime *Chem. Rev.* **2010**, 110, 3019-3042.

¹¹Li, A. X.; Hudson, R. H. E.; Barret, J. W.; Jones, C. K.; Pasternak, S. H.; Bartha, R. *Magn. Reson. Med.* **2008**, 60, 1197-1206.

¹²R.M. Henkelman, G.J. Stanisz and S.J. Graham *NMR Biomed.* **2001**, 14, 57-64.

¹³(a) Stevens, T. K.; Milne, M.; Elmhrikki, A. A. H.; Suchý, M.; Bartha, R.; Hudson, R. H. E. *Contrast Med. Mol. Imaging* **2013**, 8, 287-290. (b) Elmhrikki, A. A. H.; Milne, M.; Suchý, M.; Bartha, R.; Hudson, R. H. E. *Can. J. Chem.* **2013**, 91, 211-219.

¹⁴Mani, T.; Tircsó, G.; Zhao, P.; Sherry, A. Dean; Woods, M. *Inorg. Chem.* **2009**, 48, 10338-10345.

¹⁵H. G. Brittain and J. F. Desreux *Inorg. Chem.* **1984**, 23, 4459-4466.

¹⁶Aime, S.; Botta, M.; Garda, Z.; Kucera, B. E.; Tircso, G.; Young, V. G.; Woods, M. *Inorg. Chem.* **2011**, 50, 7955-7965.

¹⁷Reviewed by: Suchý, M.; Hudson, R. H. E. *Eur. J. Org. Chem.* **2008**, 4847-4865.

¹⁸(a) Kang, S. I.; Ranganathan, R. S.; Emswiler, J. E.; Kumar, K.; Gougoutas, J. Z.; Malley, M. F.; Tweedle, M. F. *Inorg. Chem.* **1993**, 32, 2912-2918. (b) Wiener E. C.; Abadjian, M. C.; Sengar, R.; van der Elst, L.; van Niekerk, C.; Grothjan, D. B.; Leung, P. Y.; Schulte, C.; Moore, C. E.; Rheingold, A. L. *Inorg. Chem.* **2014**, 53, 6554-6568.

¹⁹F.A. Dunand, R.S. Dickins, D. Parker and A.E. Merbach *Chem. Eur. J.*, 2001, 7, 5160.

²⁰Meng, L.; Zuo, H.; Vijaykumar, B. V. D.; Dupati, G.; Choi, K. M.; Jang, K.; Yoon, Y. J.; Shin, D. S. *Bull. Korean Chem. Soc.* **2013**, 34, 585-589.

²¹Savinov, S. N.; Austin, D. J. *Org. Lett.* **2002**, 4, 1415-1418.

²²Averill, D. J.; Allen, M. J. *Inorg. Chem.* **2014**, 53, 6257-6263.

²³Steiner, M. A.; Gatfield, J.; Brisbare-Roch, C.; Dietrich, H.; Treiber, A.; Jenck, F.; Boss, C. *ChemMedChem* **2013**, 8, 898-903.

²⁴Szymański, W.; Westerbeek, A.; Janssen, D. B.; Feringa, B. L. *Angew. Chem. Int. Ed.* **2011**, 50, 10712-10715.

ARTICLE

Journal Name

²⁵De León-Rodríguez, L. M.; Viswanathan, S.; Sherry, A. D. *Contrast Med. Mol. Imaging* **2010**, *5*, 121-125.

²⁶Suchý, M.; Li, A. X.; Bartha, R.; Hudson, R. H. E. *Bioorg. Med. Chem.* **2008**, *16*, 6156-6166.

²⁷Exchange rates were measured according to Dixon, W. T.; Ren, J.; Lubag, A. J. M.; Ratnakar, J.; Vinogradov, E.; Hancu, I.; Lenkinski, R. E.; Sherry A. D. *Magn. Reson. Med.* **2010**, *63*, 625-632.

²⁸ Zhang, S.; Michaudet, L.; Burgess, S.; Sherry, A. D. *Angew. Chem. Int. Ed.*, **2002**, *41*, 1919-1921.

²⁹Opina, A. C. L.; Wu, Y.; Zhao, P.; Kiefer, G.; Sherry, A. D. *Contrast Med. Mol. Imaging* **2011**, *6*, 459-464.

³⁰Wojciechowski, F.; Suchý, M.; Li, A. X.; Azab, H. A.; Bartha, R.; Hudson, R. H. E. *Bioconjugate Chem.* **2007**, *18*, 1625-1636.

³¹ Marques, M. P. M.; Geraldès, C.; Sherry, A. D.; Merbach, A. E. ; Powell, H.; Pubanz, D.; Aime, S.; Botta, M. *J. Alloys Compd.*, **1995**, *225*, 303-307.

³² Barge, A.; Cravotto, G.; Fedeli, F. *Contrast Med. Mol. Imaging* **2006**, *1*, 184-188.

³³ Dixon, W. T.; Ren, J.; Lubag, A. J. M.; Ratnakar, J.; Vinogradov, E.; Hancu, I.; Lenkinski, R. E.; Sherry A. D. *Magn. Reson. Med.* **2010**, *63*, 625-632.

³⁴ Meng, L.; Zuo, H.; Vijaykumar, B. V. D.; Dupati, G.; Choi, K. M.; Jang, K.; Yoon, Y. J.; Shin, D. S. *Bull. Korean Chem. Soc.* **2013**, *34*, 585-589.

³⁵ Savinov, S. N.; Austin, D. J. *Org. Lett.* **2002**, *4*, 1415-1418.

³⁶ Bai, J.; Zambroń, B. K.; Vogel, P. *Org. Lett.* **2014**, *16*, 604-607.

³⁷ Kang, S. I.; Ranganathan, R. S.; Emswiler, J. E.; Kumar, K.; Gougoutas, J. Z.; Malley, M. F.; Tweedle, M. F. *Inorg. Chem.* **1993**, *32*, 2912-2918.

³⁸ Aime, S.; Botta, M.; Garda, Z.; Kucera, B. E.; Tircso, G.; Young, V. G.; Woods, M. *Inorg. Chem.* **2011**, *50*, 7955-7965.

³⁹Suchý, M.; Milne, M.; Elmehriki, A. A. H.; McVicar, N.; Li, A. X.; Bartha, R.; Hudson, R. H. E. *J. Med. Chem.*, **2015**, *58*, 6516-6532.

