Palladium-catalysed dimerization of vinylarenes using indium triflate as an effective co-catalyst

Teruhisa Tsuchimoto, Susumu Kamiyama, Ryoju Negoro, Eiji Shirakawa* and Yusuke Kawakami Graduate School of Materials Science, Japan Advanced Institute of Science and Technology, Asahidai, Tatsunokuchi, Ishikawa 923-1292, Japan. E-mail: shira@jaist.ac.jp; Fax: 81 761 51 1631; Tel: 81 761 51 1631

Received (in Cambridge, UK) 3rd February 2003, Accepted 26th February 2003 First published as an Advance Article on the web 11th March 2003

A palladium–indium triflate catalyst was found to be much more active for the dimerization of vinylarenes compared with generally used cationic palladium(II) catalysts.

Palladium complexes are powerful catalysts for carbon-carbon bond forming reactions because palladium(II) intermediates generated by oxidative addition of organic electrophiles to palladium(0) complexes can be utilized for wide scope of transformations such as transmetalation and insertion of unsaturated bonds.1 However, organic electrophiles that can add oxidatively to palladium(0) complexes have been limited mainly to aryl/alkenyl halides/triflates.¹ Although the expansion of the scope by enhancing electrophilicity of organic compounds with Lewis acids has made a certain success, only substrates having a carbon-heteroatom bond have been used.^{2,3} On the other hand, simple alkenes activated by Lewis acids are widely known to accept addition of relatively weak nucleophiles such as arenes, i.e., the Friedel-Crafts alkylation.⁴ In this context, we have reported that metal triflates are efficient activators of alkynes and catalyse the Friedel-Crafts alkenylation of arenes.5 Thus we envisaged that metal triflates should activate vinylarenes to accept nucleophilic attack of palladium(0) complexes, giving oxidative adduct equivalents that accept insertion of another vinylarene as shown in Scheme 1. Here we report the palladium-indium triflate-catalysed dimerization of vinlyarenes through nucleophilic attack of palladium(0) complexes to vinylarenes activated by indium triflate.



Scheme 1

We first examined the catalytic activity of Pd-In(OTf)₃ for the dimerization of α -methylstyrene (1a) (Scheme 2 and Table 1). Treatment of **1a** with 1 mol % of $Pd(OAc)_2^6$ and 5 mol % of In(OTf)₃ in 1,4-dioxane at 20 °C for 3 h gave a mixture of (E)-4-methyl-2,4-diphenyl-2-pentene (2a) and 4-methyl-2,4-diphenyl-1-pentene (3a) in 78% yield with complete consumption of 1a (entry 1).⁷ Use of a catalytic amount of a phosphine such as PPh₃ or 1,3-bis(diphenylphosphino)propane (dppp) raised the yields in compensation for the reaction rate (entries 2-4). An in situ generated palladium(0) complex⁸ coordinated by 2 equivalents of PPh₃ was also effective, whereas cationic Pd(II)-PPh₃ complexes, which are known to catalyse the dimerization of alkenes,9 were totally inactive (entries 5-7). The use of Zr(OTf)₄ instead of In(OTf)₃ slightly lowered the yield, whereas metal chlorides failed to catalyse the dimerization (entries 8–11). Lack of a metal triflate or a palladium complex resulted in no or slow reaction, respectively (entries 12 and 13).



Table 1 Dimerization of α -methylstyrene (1a) catalysed by Pd–Lewis acid^{*a*}

Entry	Pd cat.	Additive (mol%)	Lewis Acid	Conv. (%) ^b	Yield (%) of 2a + 3a ^b
1	$Pd(OAc)_2$	none	In(OTf)3	>99	78
2	$Pd(OAc)_2$	$PPh_3(1)$	In(OTf) ₃	>99	86
3	$Pd(OAc)_2$	$PPh_3(2)$	In(OTf) ₃	91	83
4	$Pd(OAc)_2$	dppp (1)	In(OTf) ₃	92	92
5	Pd(0) complex ^c	$PPh_3(2)$	In(OTf) ₃	70	41
6	$[Pd(PPh_3)_2](BF_4)_2$	none	none	< 1	< 1
7	$[(\pi - C_3H_5)Pd(cod)]BF_4$	PPh ₃ (2)	none	< 1	< 1
8	$Pd(OAc)_2$	$PPh_3(1)$	Zr(OTf) ₄	99	67
9	$Pd(OAc)_2$	$PPh_3(1)$	InCl ₃	11	< 1
10	$Pd(OAc)_2$	$PPh_3(1)$	AlCl ₃	< 1	< 1
11	$Pd(OAc)_2$	$PPh_3(1)$	TiCl ₄	< 1	< 1
12	$Pd(OAc)_2$	$PPh_3(1)$	none	< 1	< 1
13	none	none	$In(OTf)_3$	46	30
T 1		1 4 1	(10 T)		<u>.</u>

^{*a*} The reaction was carried out in 1,4-dioxane (1.0 mL) at 20 °C for 3 h using **1a** (1.0 mmol) in the presence of a palladium catalyst (10 µmol), an additive and a Lewis acid (50 µmol). ^{*b*} Determined by GC. ^{*c*} A palladium(0) complex generated *in situ* by the reaction of $[(\pi$ -C₃H₅)PdCl]₂ with NaCH(CO₂Me)₂ was used.⁸

The Pd(OAc)₂/PPh₃/In(OTf)₃ (1/1/5) catalyst was successfully applied to the dimerization of various vinylarenes (Scheme 3 and Table 2). In addition to simple vinylarenes such as styrene (**1b**) and 2-vinylnaphthalene, styrenes bearing an electronwithdrawing group at the *para* position also dimerized to give 1,3-diaryl-1-butenes in high yields (entries 1–6). A styrene having a carboxyl group underwent the dimerization without protection (entry 4). Even aryl and benzyl halides, which are known to add to palladium(0) complexes, were tolerated (entries 5–7).¹ For the reaction of an electron-rich vinylarene, higher dilution that may prevent polymerization was required (entry 8). *o*-Divinylbenzene (**4**) underwent an intramolecular reaction to give 3-methylindene (**5**) in 73% yield, where CH₃CN was a superior solvent to 1,4-dioxane (Scheme 4).

We had assumed that **1a** is more susceptible than **1b** to activation by Lewis acids because its methyl group would stabilize the resulting benzyl cation, whereas cationic H–Pd complexes, active species in the cationic palladium(π)-catalysed dimerization of alkenes,⁹ prefer less bulky **1b** to **1a**. Thus, we examined the reaction with a 1:1 mixture of **1a** and **1b** using a Pd–In(OTf)₃ catalyst or a cationic Pd(π) catalyst and found that **1a** dimerized predominantly in the presence of the Pd–In(OTf)₃ catalyst, whereas a cationic complex catalysed the reaction of **1b** exclusively (Scheme 5). The different substrate preference in addition to the result that a palladium(0) is as effective as Pd(OAc)₂ (entries 3 and 5 of Table 1) should imply that a palladium(0) complex is included as an active species¹⁰ in

852



Scheme 3

Table 2 Dimerization of vinylarenes catalysed by Pd–In(OTf)₃^a

Entry	Ar	Temp (°C)	Time (h)	Yield (%) ^b
1	Ph	20	3	98
2	2-Naphtyl	50	3	79
3	$4-CF_3-C_6H_4$	50	3	86
4	4-HOCO-C ₆ H ₄	20	24	70
5	4-Cl-C ₆ H ₄	50	2.5	96
6	4-Br-C ₆ H ₄	20	3	70
7	4-ClCH ₂ -C ₆ H ₄	50	14	73
8^c	$4-CH_3-C_6H_4$	20	2	85

^{*a*} The reaction was carried out in 1,4-dioxane (1.0 mL) using **1** (1.0 mmol) in the presence of Pd(OAc)₂ (10 μmol), PPh₃ (10 μmol) and In(OTf)₃ (50 μmol). ^{*b*} Isolated yield. ^{*c*} 1,4-Dioxane (5.0 mL) was used.



Scheme 5

the present reaction not only in use of a palladium(0) complex as a precursor but also with $Pd(OAc)_2$.¹¹

In conclusion, we have disclosed that the novel catalyst system consisted of a palladium complex, PPh_3 and $In(OTf)_3$ is quite effective for the dimerization of vinylarenes. Studies on the details of the reaction mechanism as well as applications of the catalyst to various unreactive compounds are in progress in our laboratory.

We are grateful to the Ministry of Education, Sports, Culture, Science, and Technology, Japan Government, for Grant-in-Aid for Scientific Research, No. 13750794 for financial support. T.T. thanks the Teijin Ltd. for financial support through Award in Synthetic Organic Chemistry, Japan.

Notes and references

- 1 J. Tsuji, *Palladium Reagents and Catalysts*, John Wiley & Sons, Chichester, 1995, p. 125.
- 2 Lewis acids have been utilized for the palladium-catalysed carboncarbon bond forming reactions of heteroatom-containing compounds. BEt₃ for allylic alcohols: Y. Tamaru, Y. Horino, M. Araki, S. Tanaka and M. Kimura, *Tetrahedron Lett.*, 2000, **41**, 5705; Y. Horino, M. Naito, M. Kimura, S. Tanaka and Y. Tamaru, *Tetrahedron Lett.*, 2001, **42**, 3113; M. Kimura, Y. Horino, R. Mukai, S. Tanaka and Y. Tamaru, *J. Am. Chem. Soc.*, 2001, **123**, 10401; Ti(O-i-Pr)₄ for allylic alcohols: K. Itoh, N. Hamaguchi, M. Miura and M. Nomura, *J. Chem. Soc., Perkin Trans. 1*, 1992, 2883; SnCl₂ for allylic alcohols: J. P. Takahara, Y. Masuyama and Y. Kurusu, *J. Am. Chem. Soc.*, 1992, **114**, 2577; BX₃ (X = F, C₆F₅) or AlX₃ (X = Cl, Me) for α,β-unsaturated carbonyl compounds: S. Ogoshi, T. Yoshida, T. Nishida, M. Morita and H. Kurosawa, *J. Am. Chem. Soc.*, 2001, **123**, 1944; RZnI (R = Me, Et) for thioimidates: I. Ghosh and P. A. Jacobi, *J. Org. Chem.*, 2002, **67**, 9304.
- 3 The palladium-catalysed carbon-heteroatom bond forming reactions with a Lewis acid have been reported. ZnCl₂ for α,β-unsaturated carbonyl compounds: E. Keinan and N. Greenspoon, J. Am. Chem. Soc., 1986, **108**, 7314; Ti(O-i-Pr)₄ for allylic alcohols: S.-C. Yang and C.-W. Hung, J. Org. Chem., 1999, **64**, 5000; S.-C. Yang and W.-H. Chung, Tetrahedron Lett., 1999, **40**, 953; Me₃SiOTf for α,β-unsaturated carbonyl compounds or arenecarbaldehydes: S. Ogoshi, S. Tomiyasu, M. Morita and H. Kurosawa, J. Am. Chem. Soc., 2002, **124**, 11598.
- 4 G. A. Olah, R. Krishnamurti and G. K. S. Prakash in *Comprehensive Organic Synthesis*, ed. B. M. Trost, I. Fleming and G. Pattenden, Pergamon Press, New York, 1991, vol. 3, ch. 1.8, p. 293.
- 5 T. Tsuchimoto, T. Maeda, E. Shirakawa and Y. Kawakami, *Chem. Commun.*, 2000, 1573.
- 6 It is well-known that palladium(π) complexes are reduced to palladium(0) species in the presence of an alkene. For example, see p 37 of reference 1.
- 7 The reproducibility on the ratio of **2a and 3a was poor.**
- 8 T. Hayashi, M. Kawatsura and Y. Uozumi, *Chem. Commun.*, 1997, 561; M. Kawatsura, Y. Uozumi and T. Hayashi, *Chem. Commun.*, 1998, 217.
- 9 [Pd(PPh₃)₂](BF₄)₂: Z. Jiang and A. Sen, *Organometallics*, 1993, **12**, 1406 and references therein; [(π-C₃H₅)Pd(solvent)₂]BF₄-PPh₃: A. Sen and T.-W. Lai, *Organometallics*, 1983, **2**, 1059 and references therein.
- 10 No palladium(0) species is thought to be involved in the catalytic cycle of the previously reported palladium(II)-catalysed dimerization of alkenes. See reference 9.
- 11 There might be a possibility that a complex having a Pd–In bond catalyses the reaction. For late transition metal–indium bonds, see: R. A. Fischer and J. Behm, *J. Organomet. Chem.*, 1991, **413**, C10; R. A. Fischer and J. Weiβ, *Angew. Chem. Int. Ed.*, 1999, **38**, 2830.