Cite this: Org. Biomol. Chem., 2011, 9, 2062

## COMMUNICATION

## Benzyl radical addition reaction through the homolytic cleavage of a benzylic C–H bond<sup>†</sup>‡

Masafumi Ueda, Eiko Kondoh, Yuta Ito, Hiroko Shono, Maiko Kakiuchi, Yuki Ichii, Takahiro Kimura, Tetsuya Miyoshi, Takeaki Naito and Okiko Miyata\*

*Received 10th December 2010, Accepted 25th January 2011* DOI: 10.1039/c0ob01148g

Direct generation of a benzyl radical by C–H bond activation of toluenes and the addition reaction of the resulting radical to an electron deficient olefin were developed. The reaction of dimethyl fumarate with toluene in the presence of  $Et_3B$ as a radical initiator at reflux afforded 2-benzylsuccinic acid dimethyl ester in good yield.

Free radical reactions represent one of the most straightforward tools for carbon–carbon bond formation in organic synthesis.<sup>1,2</sup> Consequently, extensive studies have been demonstrated for obtaining novel radical reactions. Among them, the radical reaction mediated by triethylborane, which is recognized as an excellent radical initiator, mediator and terminator, has recently led to many novel and useful synthetic applications.<sup>3</sup> However, Et<sub>3</sub>B-mediated radical addition to  $\alpha$ , $\beta$ -unsaturated ester 1 generally does not give the corresponding adduct **3** in the absence of a hydrogen donor such as tributyltin hydride (Scheme 1).<sup>3a,4</sup>



Scheme 1 Triethylborane-mediated radical addition to a  $\alpha$ , $\beta$ -unsaturated ester.

This drawback is attributed to the observation that the intermediate enoxy radical **A** generated by the 1,4-radical addition cannot be trapped by  $Et_3B$ . Therefore,  $Et_3B$  does not function as a radical terminator. We postulated that the enoxy radical **A** and the  $\alpha$ carbonyl radical **B** have the ability to extract a benzylic hydrogen atom to generate stable benzyl radical derivatives, which can be used for the addition reaction. Although various strategies have been developed for the generation of a carbon radical center, C– H bond activation is a particularly promising and economically attractive method for the direct generation of a carbon radical.<sup>5-9</sup> Albini and co-workers have reported the generation of a benzyl radical from toluene by photosensitization; however, the addition reaction afforded a mixture of many products.<sup>10</sup> Herein, we report a novel approach for the generation of a benzyl radical and the addition reaction to an electron deficient olefin in connection with our recent efforts in radical chemistry.<sup>11</sup>

For our initial optimization studies, dimethyl fumarate **4** as a radical acceptor and toluene as benzyl radical source and solvent were selected (Table 1). When the radical reaction was carried out with 2 equiv. of Et<sub>3</sub>B in toluene (concentration 0.05 M) at room temperature, the dimethyl fumarate **4** was rapidly consumed but no isolable products were afforded, probably due to the slow hydrogen abstraction which led to the polymerization of an intermediate radical (entry 1). The reaction was run at reflux to improve the efficiency of hydrogen abstraction from toluene by an  $\alpha$ -carbonyl radical. The desired benzyl adduct **5a** was obtained as expected, albeit in 41% yield (entry 2). Upon decreasing the concentration to 0.025 M and the amount of Et<sub>3</sub>B to 1 equiv., the yield increased to 53% (entry 4). The reaction in the absence of Et<sub>3</sub>B proceeded very slowly to give **5a** in low yield (entry 6). This result indicates that Et<sub>3</sub>B plays an important role at the radical initiation step. The

 Table 1
 Benzyl radical addition to dimethyl fumarate 4

MeC	D <sub>2</sub> C CO <sub>2</sub> Me	Et₃B PhMe Me	eO <sub>2</sub> C	∠CO₂Me
	4		5a	`Ph
Entry	Concentration (M)	Et <sub>3</sub> B (equiv.)	Temp.	Yield (%) <sup>a</sup>
1	0.05	2	rt	
2	0.05	2	reflux	41
3	0.025	2	reflux	48
4	0.025	1	reflux	53
5	0.025	0.5	reflux	48
6	0.025	none	reflux	8 (83) <sup>b</sup>
7	0.01	1	reflux	74

<sup>*a*</sup> Isolated yield. <sup>*b*</sup> Yield in parentheses is for the recovered starting material **4**.

Kobe Pharmaceutical University, Motoyamakita, Higashinada, Kobe, 658-8558, Japan. E-mail: miyata@kobepharma-u.ac.jp; Fax: +81-78-441-7556 † This work is dedicated to the memory of Athel Beckwith, a teacher and scientist from whom we learned how to study chemistry by example. His pioneering advances in radical chemistry laid the foundation for much of the current radical clock methodology.

<sup>‡</sup> Electronic supplementary information (ESI) available: Experimental procedures and characterization data. See DOI: 10.1039/c0ob01148g

 Table 2
 Radical addition to dimethyl fumarate with various toluene derivatives

MeO <sub>2</sub> C´	∽∽CO₂Me 4	<u>ArCH₂R<sup>1</sup>, Et₃B</u> 180 °C 0.5-3 h	MeO <sub>2</sub> C R <sup>1</sup>	CO <sub>2</sub> Me
Entry	<b>R</b> <sup>1</sup>	<b>R</b> <sup>2</sup>	Product	Yield (%) <sup>a</sup>
1 <sup>b</sup>	Н	4-OMe	5b	88
2 <sup>c</sup>	Н	3-OMe	5c	77
3 <sup>c</sup>	Н	2-OMe	5d	77
$4^c$	Н	3,4-di-OMe	5e	39
5 <sup>b</sup>	Н	4-Me	5f	68
6 <sup>b</sup>	Н	3-Me	5g	52
$7^c$	Н	2-Me	5h	19
$8^d$	Н	4-Cl	5i	63
$9^d$	Н	4-Br	5j	66
$10^{d}$	Н	4-F	5k	36
$11^{d}$	Н	4-I	51	30
$12^{d}$	Н	3-C1	5m	57
13 <sup>c</sup>	Н	2-Cl	5n	43
$14^{d}$	Н	$4-CO_2Me$	50	64
15 <sup>b</sup>	Bn	Н	5p	$65 (6:1)^d$
16 <sup>e</sup>	Me	Н	5q	$28 (1:1)^d$

<sup>*a*</sup> Isolated yield. <sup>*b*</sup> Reactions carried out with 0.3 equiv. of Et<sub>3</sub>B. <sup>*c*</sup> Reactions carried out with 0.9 equiv. of Et<sub>3</sub>B. <sup>*a*</sup> Ratio was determined by <sup>1</sup>H NMR. The relative stereoconfiguration was not determined. <sup>*c*</sup> Reactions carried out with 0.6 equiv. of Et<sub>3</sub>B.

best result was obtained when the concentration was 0.01 M with 1 equiv. of Et<sub>3</sub>B (entry 7). The product, 2-benzylsuccinate, represents an important compound because of its utility as an intermediate in the synthesis of biologically active compounds and functional materials.<sup>12</sup> Although several synthetic methods including multistep processes have been reported,<sup>13</sup> the direct introduction of the benzyl group to fumarate has been less frequently reported.<sup>14</sup> Thus, our direct benzylation reaction is characterized as atom economical, straightforward and a valuable reaction.

To examine the scope of this direct benzylation, various toluene derivatives were subjected to the radical reaction (Table 2). We were pleased to observe that electron rich or poor toluene derivatives were readily accommodated, producing the expected adducts 5bo. It should be noted that the reaction proceeded with even less than 1 equivalent of  $Et_3B$ , owing to the higher reaction temperature employed. Reactions of 2-, 3- and 4-methoxytoluenes proceeded effectively with good yields (entries 1-3), while a lower yield was observed in the reaction with 3,4-dimethoxytoluene (entry 4). pand *m*-Xylene were also suitable reagents to use; however, *o*-xylene gave a low yield of product 5h because of its steric properties (entries 5-7). A halogen and an ester substituent showed tolerance towards this radical reaction (entries 8-14). Notably, the bromo and the iodo groups, which did not affect the course of the reaction, could serve as a handle for further transformations. Interestingly, the addition of dibenzyl produced the adduct 5p in 68% yield with a 6:1 mixture of stereoisomers, although a reduced yield of 5q and no stereoselectivity were observed with ethylbenzene (entries 15 and 16).

A competition experiment was conducted to study the influence of the electronic nature of the benzyl radical on reactivity towards the electron deficient olefin (Scheme 2). As a result of the



Scheme 2 Competition experiment.

reaction with an equimolar mixture of 4-methoxytoluene and 4-chlorotoluene, the 4-methoxybenzyl radical exhibited higher reactivity in the formation of **5b**. This indicated that the methoxy group as an electron donating group increased the SOMO energy level of the nucleophilic benzyl radical; thus, interaction of the LUMO of the olefin with the SOMO of the benzyl radical would be enhanced.

We have also extended the acceptor tolerance to 1,2-dicyanoand 1,2-diketo olefins (Scheme 3). Fumaronitrile 6 showed excellent reactivity giving the desired product 7 in quantitative yield. 1,2-Diketone 8 was also found to react with 4methoxytoluene to afford the adduct 9, albeit in relatively lower yield due to the competition with the ethyl radical addition reaction.



Scheme 3 4-Methoxybenzyl radical addition to the electron deficient olefins.

A possible reaction pathway is presented in Scheme 4. The first step involves the generation of an ethoxy radical and an ethylperoxy radical by the oxidative decomposition of  $Et_3B$  with molecular oxygen. The radicals then abstract a hydrogen atom from the toluene derivatives to produce a benzyl radical **C**. The benzyl radical can add to fumarate to form the  $\alpha$ -carbonyl radical **D**. The trapping of the radical **D** by toluene derivatives gave product **5** and the regeneration of the benzyl radical which is used for the radical chain cycle.

In conclusion, we have demonstrated the direct generation of a benzyl radical by C–H bond activation of toluenes and the addition reaction of the resulting radical to an electron deficient olefin.

## Acknowledgements

This work was supported in part by Grants-in Aid from the Ministry of Education, Culture, Sports, Science and Technology



Scheme 4 Possible reaction pathway

of Japan, and the Science Research Promotion Fund of the Japan Private School Promotion Foundation. M.U. is grateful to the Research Foundation for Pharmaceutical Sciences for a research grant.

## Notes and references

- (a) J. Fossey, D. Lefort and J. Sorba, Free Radicals in Organic Chemistry. translated by J. Lomas, John Wiley & Sons, New York, 1995; (b) I. Ryu, N. Sonoda and D. P. Curran, Chem. Rev., 1996, 96, 177; (c) B. B. Snider, Chem. Rev., 1996, 96, 339; (d) B. Giese, B. Kopping, T. Göbel, J. Dickhaut, G. Thoma, K. J. Kulicke and F. Trach, Org. React. (N.Y.), 1996, 48, 301; (e) P. Renaud and M. Gerster, Angew. Chem., Int. Ed., 1998, 37, 2562; (f) M. P. Sibi and N. A. Porter, Acc. Chem. Res., 1999, 32, 163; (g) W. R. Bowman, C. F. Bridge and P. Brookes, J. Chem. Soc., Perkin Trans. 1, 2000, 1; (h) Radicals in Organic Synthesis, ed. P. Renaud, and M. P. Sibi, Wiley-VCH, Weinheim, 2001, Vol. 1 and 2.
- 2 For recent reviews, see: (a) G. J. Rowlands, *Tetrahedron*, 2010, 66, 1593;
   (b) G. J. Rowlands, *Tetrahedron*, 2009, 65, 8603.
- 3 (a) C. Ollivier and P. Renaud, *Chem. Rev.*, 2001, **101**, 3415; (b) V. Darmency and P. Renaud, *Top. Curr. Chem.*, 2006, **263**, 71; (c) H. Yorimitsu and K. Oshima in *Radicals in Organic Synthesis*, ed. P. Renaud and M. P. Sibi, Wiley-VCH, Weinheim, 2001, Vol. 1, p. 11; (d) H. Miyabe, M. Ueda and T. Naito, *Synlett*, 2004, 1140.
- 4 S. Bazin, L. Feray, D. Siri, J. V. Naubron and M. P. Bertrand, *Chem. Commun.*, 2002, 2506.
- 5 For the generation and reaction of α-aminoalkyl radicals, see: (a) T. Yoshimitsu, Y. Arano and H. Nagaoka, J. Am. Chem. Soc., 2005, 127, 11610; (b) T. Yoshimitsu, C. Atsumi, E. Iimori, H. Nagaoka and T. Tanaka, *Tetrahedron Lett.*, 2008, 49, 4473; (c) T. Yashimitsu, K. Matsuda, H. Nagaoka, K. Tsukamoto and T. Tanaka, Org. Lett., 2007, 9, 5115; (d) C. Brulé and N. Hoffmann, *Tetrahedron Lett.*, 2002, 43, 69; (e) S. Kim, N. Kim, W. Chung and C. H. Cho, Synlett, 2001, 937.
- 6 For the generation and reaction of α-oxyalkyl radicals, see: (a) T. Yoshimitsu, M. Tsunoda and H. Nagaoka, Chem. Commun., 1999, 1745; (b) T. Yoshimitsu, Y Arano and H. Nagaoka, J. Org. Chem., 2003, 68, 625; (c) T. Yoshimitsu, T. Makino and H. Nagaoka, J. Org. Chem., 2003, 68, 7548; (d) T. Yoshimitsu, Y. Arano and H. Nagaoka, J. Org. Chem., 2005, 70, 2342; (e) A. J. Clark, S. Rooke, T. J. Sparey and P. C. Taylor, Tetrahedron Lett., 1996, 37, 909; (f) K. Yamada, H. Fujihara, Y. Yamamoto, Y. Miwa, T. Taga and K. Tomioka, Org. Lett., 2002, 4,

3509; (g) T. Akindele, Y. Yamamoto, M. Maekawa, H. Umeki, K. Yamada and K. Tomioka, *Org. Lett.*, 2006, **8**, 5729; (h) K. Yamada, H. Umeki, M. Maekawa, Y. Yamamoto, T. Akindele, M. Nakano and K. Tomioka, *Tetrahedron*, 2008, **64**, 7258; (i) Y. Yamamoto, M. Mekawa, T. Akindele, K. Yamada and K. Tomioka, *Tetrahedron*, 2005, **61**, 379; (j) Y. Yamamoto, K. Yamada and K. Tomioka, *Tetrahedron*, 2005, **61**, 379; (j) Y. Yamamoto, K. Yamada, Y. Yamamoto, M. Maekawa and K. Tomioka, *J. Org. Chem.*, 2004, **69**, 1531; (l) K. Hirano, S. Sakaguchi and Y. Ishii, *Tetrahedron Lett.*, 2002, **43**, 3617; (m) see ref. 5e.

- 7 For the generation and reaction of diphenylmethyl radicals, see: (a) Z. Li, L. Cao and C.-J. Li, Angew. Chem., Int. Ed., 2007, 46, 6505; (b) C. A. Correia and C.-J. Li, Tetrahedron Lett., 2010, 51, 1172; (c) N. Borduas and D. A. Powell, J. Org. Chem., 2008, 73, 7822; (d) Á. Pintér, A. Sud, D. Sureshkumar and M. Klussmann, Angew. Chem., Int. Ed., 2010, 49, 1; (e) J. Jin, Y. Li, Z. Wang, W. Qian and W. Bao, Eur. J. Org. Chem., 2010, 1235.
- 8 For the generation and reaction of unstabilized alkyl radicals, see: (a) K. Yamada, Y. Yamamoto, M. Maekawa, J. Chen and K. Tomioka, *Tetrahedron Lett.*, 2004, **45**, 6595; (b) S. Kato, T. Iwahama, S. Sakaguchi and Y. Ishii, J. Org. Chem., 1998, **63**, 222; (c) T. Hara, T. Iwahama, S. Sakaguchi and Y. Ishii, J. Org. Chem., 2001, **66**, 6425; (d) see ref. 5e.
- 9 For the generation and reaction of acyl radicals, see: (a) V. Chudasama, R. J. Fitzmaurice and S. Caddick, *Nature. Chem.*, 2010, **2**, 592; (b) C. Chatgilialoglu, D. Crich, M. Komatsu and I. Ryu, *Chem. Rev.*, 1999, **99**, 1991; (c) K. Yoshikai, T. Hayama, K. Nishimura, K. Yamada and K. Tomioka, *J. Org. Chem.*, 2005, **70**, 681; (d) K. Yoshikai, T. Hayama, K. Nishimura, K. Yamada and K. Tomioka, *Chem. Pharm. Bull.*, 2005, **53**, 586.
- 10 M. Mella, M. Fagnoni and A. Albini, Eur. J. Org. Chem., 1999, 2137.
- (a) M. Ueda, H. Miyabe, M. Torii, T. Kimura, O. Miyata and T. Naito, Synlett, 2010, 1341; (b) M. Ueda, E. Iwasada, H. Miyabe, O. Miyata and T. Naito, Synthesis, 2010, 1999; (c) M. Ueda, H. Miyabe, T. Kimura, E. Kondoh, T. Naito and O. Miyata, Org. Lett., 2009, 11, 4632; (d) H. Rahaman, M. Ueda, O. Miyata and T. Naito, Org. Lett., 2009, 11, 2651; (e) M. Ueda, H. Miyabe, H. Shimizu, H. Sugino, O. Miyata and T. Naito, Angew. Chem., Int. Ed., 2008, 47, 5600.
- (a) J. P. Devlin, W. D. Ollis, J. E. Thorpe, R. S. Wood, B. J. Broughton, P. J. Warren, K. R. H. Wooldbridge and D. E. Wright, J. Chem. Soc., Perkin Trans. 1, 1975, 830; (b) D. E. Levy, F. Lapierre, W. Liang, W. Ye, C. W. Lange, X. Li, D. Grobelny, M. Casabonne, D. Tyrrell, K. Holme, A. Nadzan and R. E. Galardy, J. Med. Chem., 1998, 41, 199; (c) W. C. Groutas, M. J. Brubaker, M. A. Stanga, J. C. Castrisos, J. P. Crowley and E. J. Schatz, J. Med. Chem., 1989, 32, 1607; (d) R. B. Bates, R. S. Cuther and R. M. Freeman, J. Org. Chem., 1977, 42, 4162; (e) G. Sabitha, R. Srividya and J. S. Yadav, Tetrahedron, 1999, 55, 4015.
- 13 (a) Y. Liu, C. A. Sandoval, Y. Yamaguchi, X. Zhang, Z. Wang, K. Kato and K. Ding, J. Am. Chem. Soc., 2006, **128**, 14212; (b) R. J. Moss, K. J. Wadsworth, C. J. Chapman and C. G. Frost, Chem. Commun., 2004, 1984; (c) M. Dai, C. Wang, G. Dong, J. Xiang, T. Luo, B. Liang, J. Chen and Z. Yang, Eur. J. Org. Chem., 2003, 4346; (d) S. Takeuchi, Y. Ukaji and K. Inomata, Bull. Chem. Soc. Jpn., 2001, **74**, 955; (e) N. A. Porter, D. M. Scott, I. J. Rosenstein, B. Giese, A. Veit and H. G. Zeitz, J. Am. Chem. Soc., 1991, **113**, 1791; (f) S. C. Bergmeier and K. A. Ismail, Synthesis, 2000, 1369; (g) R. Ballini, G. Bosica, D. Fiorini and P. Righi, Synthesis, 2002, 681.
- 14 Direct introduction of the benzyl group to fumarate with benzyl bromide<sup>14a</sup> or benzyltrimethylsilane<sup>14b</sup> has been reported, see: (a) P. He, P. Watts, F. Marken and S. J. Haswell, *Angew. Chem., Int. Ed.*, 2006, **45**, 4146; (b) L. Cermenati, M. Mella and A. Albini, *Tetrahedron*, 1998, **54**, 2575.