View Article Online View Journal

Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. Costil, F. Fernandez-Nieto, R. Atkinson and J. Clayden, *Org. Biomol. Chem.*, 2018, DOI: 10.1039/C8OB00551F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the **author guidelines**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the ethical guidelines, outlined in our <u>author and reviewer resource centre</u>, still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/obc



COMMUNICATION

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

 α -Methyl phenylglycines by asymmetric α -arylation of alanine and their effect on the conformational preference of helical Aib foldamers

Romain Costil,^a Fernando Fernández Nieto,^b Rachel C. Atkinson,^b and Jonathan Clayden^a*

 α -Arylated alanine derivatives were made enantioselectively by migratory rearrangement of a urea derivative using (*R*,*R*)pseudoephedrine as a chiral auxiliary. Incorporation of a single residue of the product α -methyl phenylglycine into an otherwise achiral oligomer of aminoisobutyric acid oligomer induced a preferred screw sense, detectable by a NMR reporter located at the remote terminus of the oligomer. The magnitude of the screw sense induction was greater when the chiral residue was located at the N-terminus of the foldamer, and in some cases the sense of induction was opposite to that of related α -methylated amino acids with α -substituents other than aryl.

Incorporation of an additional substituent at the α position of a typical proteinogenic amino acid has important consequences for the properties of the resulting quaternary amino acid. Several families of fungal non-ribosomal peptides incorporate quaternary amino acids into their type sequence. The most common of these is the achiral residue α -aminoisobutyric acid (Aib),¹ but also prevalent are the dialkylated chiral residues as isovaline, α -methylvaline, and α -methyl phenylalanine. These quaternary amino acid residues typically induce a strong preference for helical conformations, and extensive conformational studies have been carried out on peptides containing such residues.²

 α -Methyl phenylglycine ((α Me)Phg) and its derivatives have been studied over the last two decades in programs involving peptide conformation and catalysis,³ as resolving agents in resolutions through selective crystallization⁴ or as alternative sweeteners.⁵ Its derivatives are also found as a structural feature in many targets of medicinal chemistry (Figure 1),⁶ making effective methods for its synthesis particularly valuable.



Figure 1: Examples of bioactive α -methyl phenylglycine derivatives.

In general, quaternary amino acids have been synthesised by alkylation of their naturally occurring tertiary parent, with the configuration of the product being controlled either by some form of chiral memory or by an asymmetric alkylation step.⁷ The corresponding introduction of an aryl substituent is a much more challenging prospect. We have shown that N'-aryl urea derivatives of amino acids undergo rearrangements that yield hydantoin derivatives of quaternary amino acids bearing an lpha aryl substituent, 8 and Kawabata *et al.* concurrently reported a related reaction that allows incorporation of aryl substituents with chiral memory.⁹ More recently, we reported a general approach to the stereoselective arylation of α -amino acids using pseudoephedrine as a chiral auxiliary.¹⁰ This rearrangement was facilitated by temporary in situ protection of the urea as its N- or O-silyl derivative by treatment with trimethylsilyl chloride.

We now report that it is possible to synthesise enantioenriched α -methyl phenylglycine derivatives by a practical modification of this rearrangement method, in which a 2,4dimethoxybenzyl (DMB) group is used for protection of the urea *N* atom. We show that the resulting quaternary amino acid may be incorporated into a helical Aib-containing oligomer at either the C- or the N-terminus, and that the stereogenic centre of this single quaternary residue may

^{a.} School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK. E-mail: j.clayden@bristol.ac.uk

^{b.} School of Chemistry, University of Manchester, Oxford Road, Manchester M13 9PL, UK.

Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x

induces a screw sense preference in the whole of the resulting helical oligomer.

Results and Discussion

Enantioselective α -arylation of alanine

Starting materials **3a-d** for stereoselective rearrangement were made straightforwardly from alanine derivatives in four steps (Scheme 1). *N*-protection of alanine esters **1** with the 2,4-dimethoxybenzyl (DMB) group was achieved by reductive amination with 2,4-dimethoxybenzaldehyde. Coupling of the amine with a series of carbamoyl chlorides, followed by hydrolysis of the ester moiety with lithium hydroxide in a biphasic system, afforded carboxyureas **2a-d** in excellent yields. These carboxylic acids were coupled to (*R*,*R*)pseudoephedrine without epimerisation using standard coupling reagents EDC.HCl and HOBt.H₂O to give **3a-d** in excellent overall yield.

The reaction conditions for stereoselective arylation were optimised by exploring base-promoted phenyl migration within compound 3a. Optimal conditions were found to require an excess of lithium diisopropylamide (LDA) in THF in the presence of ten equivalents of lithium chloride. Full conversion was attained after deprotonation at -78 °C followed by a period of three hours at room temperature. As observed previously by IR,⁸ enolate formation is followed by migration of the ring to the α position of the alanine residue, followed by cyclisation to the hydantoin with concurrent expulsion of the pseudoephedrine auxiliary. N-to-C migration of a selection of aryl rings was performed in moderate to good yield (Table 1), with enantiomeric excesses ranging from nearracemic (compound 4c) to complete enantiopurity (compound 4a). The reason for the strong variation in stereoselectivity with migrating ring is unclear, and shows that an alternative more general method is required for rings other than phenyl.¹¹



Scheme 1: Reagents and conditions: (a) 2,6-dimethoxybenzaldehyde, NaOAc, NaHB(OAc)₃, EtOH, rt; (b) ArN(R')COCl, triethylamine, MeCN, reflux; (c) LiOH, THF/H₂O, 45°C; (d) EDC.HCl, HOBt.H₂O, (*R*,*R*)-pseudoephedrine, CH_2Cl_2 , rt. DMB = 2,4-dimethoxybenzyl.

Table 1: Stereoelective rearrangement of the pseudoephedrine derived compounds New Article Online







Scheme 2: Hydrolysis of the hydantoin. Reagents and conditions: (a) TfOH, TFA, 0°C; (b) NaOH, H_2O , dioxane, 100°C; (c) SOCl₂, MeOH, rt. DMB = 2,4-dimethoxybenzyl.

hvdantoin **4a** was converted into the Enantiopure corresponding amino acid methyl ester (Scheme 2). Acidic removal of the dimethoxybenzyl group afforded free hydantoin 5a, which was hydrolysed under basic conditions to α-methyl phenylglycine, isolated as its methyl ester. Comparison of the optical rotation with reported values allowed us to assign the stereochemistry at the quaternary carbon to be $R.^{5}$ Interestingly, the stereocenter formed in this rearrangement is the opposite of the major enantiomer formed in our previous, protecting group-free, rearrangement using the same enantiomer of pseudoephedrine.¹⁰ Deprotection of hydantoin **4b** followed by hydrolysis gave only the corresponding amino N-methyl amide 6b despite forcing conditions (See the Supporting Information for details), suggesting that additional substitution not only decreases stereoselectivity but also hinders significantly the hydrolysis of the products.

Synthesis and conformation of foldamers containing an α -aryl quaternary amino acid residue

Homo-oligomers of the achiral quaternary amino acid Aib form 3_{10} helices as an interconverting mixture of M and P enantiomeric conformers, but even a single chiral amino acid can powerfully bias the M:P ratio, with the absolute screw sense induction depending on the nature and location of the chiral residue.¹² No information is available on the influence of α -arylated residues on screw sense preference, so the chiral quaternary amino acid derivatives previously made were coupled to achiral Aib oligomers at either the N- or the C-

Journal Name

Published on 26 March 2018. Downloaded by University of New England on 26/03/2018 21:25:25

COMMUNICATION

Journal Name

terminus and their conformational influence was explored by NMR and by Circular Dichroism (CD).

(aMe)PhgOMe was converted to its N-methyl amide derivative 6a (See SI for details) and, together with its orthomethoxylated analogue 6b, coupled to the C-terminus of an Aib tetramer by nucleophilic attack on the foldamer's azlactone derivative 7 (Scheme 3). In order to monitor the global screw sense preference of the resulting foldamer, the tetramer was coupled to a N-terminal azepine probe.^{12b,13} Hydrogenation of the N-terminal azide allowed introduction of the probe in good yield by acylation with its succinimidyl ester derivative

Addition of DMSO-d6 to a solution of 9a in CD₂Cl₂/CD₃OH 9:1 led to a significant deshielding of the two highest field N-H protons (Figure 2). This downfield shift suggests that the two NH protons furthest from the controller in 9a are not part of a hydrogen-bonding network, consistent with the adoption of a conformation. Further 3₁₀-helical support for this interpretation was gained from the CH-NH regions of the ROESY spectrum in CD_2Cl_2/CD_3OH . A diagnostic CH_{ℓ}^{β} -NH_{i+3} cross peak is visible (Figure 3, circled in red), confirming the presence of a 3_{10} -helix along the structure of **9a**. The absence of $CH_{i-}NH_{i+4}$ cross peak suggests that there is no α -helical contribution to the structure.



dibenz[c,e]azepine hydrochloride, DIPEA, MeCN, rt. DSC = N,N'-Disuccinimidyl carbonate.



Figure 2: DMSO-d6 titration of 9a (approx. 0.01 M in CD₂Cl₂/CD₃OH 9:1)



Figure 3: Aromatic region of the ROESY spectrum of 9a (approx. 0.01 M in CD₂Cl₂/CD₃OH 9:1).



Figure 4: CD spectrum of 9a (0.68 mM in MeOH).

¹H NMR in CD₃OD revealed diastereotopic signals for the methylene groups of the azepine probe, indicating an excess of one screw-sense conformer over the other.¹⁴ By comparison with the values obtained for other conformational controllers with known equilibrium ratios, 13a we deduced that (*R*)-(αMe)Phg exhibits rather modest levels of control, with a helical excess (h.e.) of just 10% in a mixture of CD₃OH and CDCl₃ (9:1). The ortho-methoxy substituent of compound 9b increases the control slightly to a h.e. of 21% in deuterated methanol.

Published on 26 March 2018. Downloaded by University of New England on 26/03/2018 21:25:25

The CD spectrum of peptide **9a** in methanol showed a local maximum centered around 250 nm for the biaryl probe (Figure 4). The molar ellipticity at this wavelength indicates a helical excess of about 7% induced at the probe (in line with the low value deduced by ¹H NMR spectroscopy), and its positive value indicates a right-handed (or *P*) screw-sense preference in solution. C-terminal amides of D-amino acids typically induce *M* helices whether the chiral residue is tertiary or quaternary,^{12c} so this weak preference is opposite to that induced by related amino acids, presumably because of the unusual α -arylated structure of the controlling residue.

The effect of introducing **6a** at the *N*-terminus of the oligomer was explored using an Aib tetramer bearing a C-terminal thionoglycine probe (Scheme 4).¹⁵ The methyl ester of **6a** was first capped with a carboxybenzyl (Cbz) protecting group and hydrolysed to the corresponding carboxylic acid. Tetramethylfluoroformamidinium hexafluorophosphate was used to form its acyl fluoride derivative, which was coupled to a hexamer already containing the embedded thioamide probe to afford foldamer **10**.

Variable Temperature ¹H NMR of peptide **10** in CD_3OH , between 5 and 35°C showed a significant downfield drift of the chemical shift of two of the amide protons. This suggests that only these two amide protons lie outside of the hydrogenbonded network, consistent with the folding of **10** into a 3₁₀ helix (Figure 5). In CD₃OD at room temperature, the foldamer shows a characteristic pair of signals for the diastereotopic methylene protons of the thionoglycine residue, centred around 4.20 ppm. Comparison of their anisochronicity with reported values¹⁵ indicates that this N-terminal controller induces a helical excess of 42%, similar to that of L-CbzPhe.



Scheme 4: Reagents and conditions: (a) CbzCl, DIPEA, CH₂Cl₂, rt; (b) NaOH, MeOH, H₂O, rt; (c) TFFH, Pyridine, CH₂Cl₂, rt; (d) H-Aib₄Gly-ψ[CSNH]AibOMe, DIPEA, CH₂Cl₂, rt.





Figure 6: CD spectrum of 10 (0.62 mM in MeOH).

The CD spectrum of the foldamer shows the typical absorption band for the π - π^* transition of the thionoglycine probe centered at 268 nm (Figure 6). The negative value of the ellipticity in this region indicates a left-handed screw-sense preference (or *M* helix) for compound **10**, indicating that, at this N terminus of the foldamer, (*R*)-(α Me)Phg induces the same screw-sense preference as the related quaternary amino acid (*R*)-(α Me)Val, presumably by the mechanism of induction previously proposed.^{12,16,17}

Conclusions

Aryl substituents may be introduced enantioselectively to the α -carbon of alanine derivatives using the base-promoted rearrangement of a urea derivative carrying using pseudo-ephedrine as a chiral auxiliary. While migration of a phenyl ring occurs with excellent enantiomeric excess, other migrating rings showed low selectivity or low yield. Protection of the urea by 2,4-dimethoxybenzyl permits simple operational conditions, leading to quaternary arylated amino acids after deprotection and hydrolysis of the resulting hydantoin. Isolation of the opposite enantiomer to that obtained by a previously reported procedure¹⁰ allows access to both enantiomers of the quaternary amino acid derivative by using the same chiral auxiliary.

 $(\alpha Me)Phg$ **6a** (and its *ortho*-methoxylated derivative **6b**) participates in a 3₁₀ helix when incorporated at either the N- or the C- terminus of an Aib oligomer. While these two unnatural amino acid derivatives induce moderate control over the overall screw-sense from the C-terminus, $(\alpha Me)Phg$ displays more promising levels of conformational control from the Nterminus.

Acknowledgements

This work was supported by the European Research Council (Advanced Grant ROCOCO).

Conflicts of interest

Page 4 of 6

Journal Name

DOI: 10.1039/C8OB00551F

Figure 5: Variable Temperature ¹H NMR of **10** (approx. 0.01 M in CD₃OH). View Article Online Published on 26 March 2018. Downloaded by University of New England on 26/03/2018 21:25:25

Journal Name

There are no conflicts to declare.

16 M. De Poli, M. De Zotti, J. Raftery, J. A. Aguilar, G. A. Morris and J. Clayden, J. Org. Chem., 2013, 78 22:48.1039/C8OB00551F

Notes and references

- (a) C. de la Fuente-Núñez, L. Whitmore and B.A. Wallace, in Handbook of Biologically Active Peptides, ed. A. J. Kastin, Elsevier, Amsterdam, 2nd edn, 2013, vol. 2, ch. 22, pp 150-156; (b) S. J. Pike, J. E. Jones, J. Raftery, J. Clayden and S. J. Webb, Org. Biomol. Chem., 2015, 13, 9580.
- (a) J. E. Jones, V. Diemer, C. Adam, J. Raftery, R. E. Ruscoe, J. Sengel, M. I. Wallace, A. Bader, S. L. Cockroft, J. Clayden and S. J. Webb, J. Am. Chem. Soc., 2016, 132, 688; (b) I. Maffucci, S. Pellegrino, J. Clayden and A. Contini, J. Phys. Chem. B, 2015, **119**, 1350.
- 3 (a) F. Formaggio, A. Barazza, A. Bertocco, C. Toniolo, Q. B. Broxterman, B. Kaptein, E. Brasola, P. Pengo, L. Pasquato and P. Scrimin, J. Org. Chem., 2004, 69, 3849; (b) H. Gregory, D. S. Jones and J. S. Morley, J. Chem. Soc. C, 1968, 531.
- 4 S. Mueller, G. J. A. Ariaans, B. Kaptein, Q. B. Broxterman, F. Formaggio, E. Battan, M. Crisma, C. Toniolo and A. Bruggink, Tetrahedron: Asymmetry, 2004, 15, 1919.
- 5 E. Mossel, F. Formaggio, M. Crisma, C. Toniolo, Q. B. Broxterman, W. H. J. Boesten, J. Kamphuis, P. J. L. M. Quaedflieg and P. Temussi, Tetrahedron: Asymmetry, 1997, 8, 1305.
- (a) A. Mete, K. Bowers, E. Chevalier, D. K. Donald, H. 6 Edwards, K. J. Escott, R. Ford, K. Grime, I. Millichip, B. Teobalda and V. Russell, Bioorg. Med. Chem. Lett., 2011, 21, 7440; (b) J. A. Sternberg, D. Geffken, J. B Adams Jr, R. Pöstages, C. G. Sternberg, C. L. Campbell and W. K Moberg, Pest. Manag. Sci., 2001, 57, 143.
- 7 (a) Y. Ohfune and T. Shinada, Eur. J. Org. Chem., 2005, 5127; (b) C. Cativiela and M. D. Díaz-de-Villegas, Tetrahedron: Asymmetry, 2007, 18, 56; (c) H. Vogt and S. Bräse, Org. Biomol. Chem. 2007, 5, 406; (d) M. I. Calaza and C. Cativiela, Eur. J. Org. Chem., 2008, 3427.
- R. C. Atkinson, D. J. Leonard, J. Maury, D. Castagnolo, N. Volz and J. Clayden, Chem. Commun., 2013, 49, 9734.
- K. Tomohara, T. Yoshimura, R. Hyakutake, P. Yang and T. Kawabata, J. Am. Chem. Soc., 2013, 135, 13294.
- R. C. Atkinson, F. Fernández-Nieto, J. Mas Roselló and J. 10 Clayden, Angew. Chem. Int. Ed., 2015, 54, 8961.
- D. Leonard, J. W. Ward and J. Clayden, Manuscript in 11 preparation.
- 12 (a) M. De Poli, L. Byrne, R. A. Brown, J. Solà, A. Castellanos, T. Boddaert, R. Wechsel, J. D. Beadle and J. Clayden, J. Org. Chem., 2014, 79, 4659; (b) J. Clayden, A. Castellanos, J. Sola and G. A. Morris, Angew. Chem. Int. Ed., 2009, 48, 5962; (c) B. A. F. Le Bailly and J. Clayden, Chem. Commun. 2014, 50, 7949; (d) M. de Zotti, F. Formaggio, M. Crisma, C. Peggion, A. Moretto, C. Toniolo J. Pept. Sci. 2014, 20, 148; (e) M. Crisma, M. de Zotti, F. Formaggio, C. Peggion, A. Moretto, C. Toniolo J. Pept. Sci. 2015, 21, 147; (f) I. Maffucci, S. Pellegrino, J. Clayden, A. Contini J. Phys. Chem. B 2015, 119, 1350; (g) I. Maffucci, J. Clayden, A. Contini J. Phys. Chem. B 2015, 119, 14003; (h) J. Solà, M. Helliwell, J. Clayden J. Am. Chem. Soc. 2010. 132. 4548.
- 13 (a) V. Diemer, J. Maury, B. A. F. Le Bailly, S. J. Webb and J. Clayden, Chem. Commun. 2017, 53, 10768; (b) B. A. F. Le Bailly, L. Byrne, V. Diemer, M. Foroozandeh, G. A. Morris and J. Clayden, Chem. Sci., 2015, 6, 2313.
- 14 Chemical shift separation quantifies conformational preference in a system such as this, which is in fast conformational exchange on the NMR timescale: J. Solà, G. A. Morris and J. Clayden, J. Am. Chem. Soc., 2011, 133, 3712.
- 15 M. De Poli and J. Clayden, Org. Biomol. Chem., 2014, 12, 836.

- 17 The molar ellipticity at this wavelength is consistent with a foldamer with a helical excess of only 16%. This discrepancy is presumably due to the Cotton effect arising from the aromatic ring in the chiral residue. Related 'local' effects have been shown to make CD an unreliable reporter of screw sense preference in some cases, see: R. A. Brown, T. Marcelli, M. De Poli, J. Solà and J. Clayden Angew. Chem. Int. Ed., 2012, 51, 1395.

Graphical abstract:



Asymmetric arylation of an alanine derivative provides (R)- α -methyl phenylglycine, which induces a right handed screw sense in an Aib oligomer.

Organic & Biomolecular Chemistry Accepted Manuscript