## Reductive Cleavage of the Se–Se Bond by the Sm–Me<sub>3</sub>SiCl–H<sub>2</sub>O System: Preparation of Unsymmetrical Phenyl Selenides<sup>†</sup>

Lei Wang and Yongmin Zhang\*

Department of Chemistry, Hangzhou University, Hangzhou, 310028, PR China

The reduction of diphenyl diselenide by the Sm–Me<sub>3</sub>SiCl–H<sub>2</sub>O system led to a selenide anion. This 'living' species reacted with organic halides, epoxides,  $\alpha$ , $\beta$ -unsaturated esters and  $\alpha$ , $\beta$ -unsaturated nitriles to afford unsymmetrical phenylselenides in good yields under mild and neutral conditions.

Organoselenium compounds have received considerable attention as useful synthetic reagents and intermediates in organic synthesis.<sup>1</sup> While there are many methods for the introduction of a seleno-substituent into organic molecules, the use of phenyl selenide anions is especially convenient and common. Several methods for the synthesis of selenide anions have been recommended, the more important of which include the use of diphenyl diselenide with sodium borohydride,<sup>2</sup> reduction of the diselenide with sodium,<sup>3</sup> samarium diiodide<sup>4</sup> and lithium aluminium hydride.<sup>5</sup> Alternatively, sodium phenyl selenide may be obtained from the selenol using sodium hydride or even by treatment with aqueous sodium hydroxide under certain conditions.<sup>6</sup> Moreover, Grignard reagents also react with selenium to give selenide anions.<sup>7</sup>

As a powerful, versatile and either-soluble one-electron transfer reducing agent,  $SmI_2$  has been widely applied in organic synthesis.<sup>8</sup> Though  $SmI_2$  is a useful reagent, storage is difficult because it is very sensitive to air oxidation. On the other hand, metallic samarium is stable in air and its strong reducing power ( $Sm^{3+}/Sm - 2.41$  V) is similar to that of magnesium ( $Mg^{2+}/Mg - 2.37$  V) and superior to that of zinc ( $Zn^{2+}/Zn - 0.71$  V). These properties prompted us to use the more convenient and cheaper samarium directly as a reductant instead of samarium(II) iodide. Recently, there have been reports on the direct use of Sm in organic synthesis.<sup>9</sup> Ishii and co-workers have shown that the Sm-Me<sub>3</sub>SiCl-NaI and Sm-Me<sub>3</sub>SiBr systems can be used for the intermolecular carbon–carbon bond formation reaction of carbonyl compounds.<sup>10</sup> Herein, we report that the Sm-Me<sub>3</sub>SiCl-H<sub>2</sub>O system promotes cleavage of the Se-Se



bond to form a phenyl selenide anion. This species reacted with organic halides, epoxides,  $\alpha,\beta$ -unsaturated esters and  $\alpha,\beta$ -unsaturated nitriles to afford unsymmetrical phenylselenides in good yields under mild and neutral conditions. The results are summarized in Table 1.

We found that under mild conditions, phenyl selenide anion, readily prepared *in situ* from the cleavage of the Se–Se bond of diphenyl diselenide with the Sm–Me<sub>3</sub>SiCl– H<sub>2</sub>O reduction system, reacted easily with epoxides,  $\alpha,\beta$ unsaturated esters and  $\alpha,\beta$ -unsaturated nitrile respectively to afford the desired unsymmetrical alkyl phenyl selenides in good yields. From Table 1, we also found that the phenyl selenide anion, a 'living' species, reacted smoothly with active organic halides such as 2-bromoacetophenone and ethyl bromoacetate to give products in good yields at 45 °C.

 Table 1
 Preparation of unsymmetrical phenyl selenides via phenyl selenide anion intermediate

Entry	R <sup>1</sup> X	R <sup>2</sup>	R <sup>3</sup>	Z	Coupling reaction conditions		
					<i>7</i> /°C	<i>t</i> /h	Yield (%) <sup>a</sup>
ab cd <sup>o</sup> ef gh i j k	PhCH <sub>2</sub> Cl PhCOCH <sub>2</sub> Br BrCH <sub>2</sub> CO <sub>2</sub> Et n-C <sub>4</sub> H <sub>9</sub> Br n-C <sub>4</sub> H <sub>9</sub> Br n-C <sub>4</sub> H <sub>9</sub> Cl n-C <sub>6</sub> H <sub>13</sub> Br n-C <sub>8</sub> H <sub>17</sub> Br	H CH₂CI	H H Me	CN CO₂Me CO₂Me	65 45 65 65 65 65 65 45 45 45 45 45 45	4 3 4 4 10 4 2 2 2 2 2 2 2	75 72 66 68 32  70 72 80 82 79 84 83

<sup>a</sup>Isolated yields. Reaction conditions: diphenyl diselenide (0.5 mmol), Sm (1.0 mmol), Me<sub>3</sub>SiCl (1.0 ml), THF (5 ml), H<sub>2</sub>O (18  $\mu$ l), cleavage reaction temperature, 45 °C, 3 h. <sup>b</sup>In the absence of water.

\*To receive any correspondence. †This is a **Short Paper** as defined in the Instructions for Authors, Section 5.0 [see *J. Chem. Research* (*S*), 1998, Issue 1]; there is therefore no corresponding material in *J. Chem. Research* (*M*).

With the less active halides  $R^2X$  such as alkyl bromides and benzyl chloride, the coupling reaction should be carried out at a higher temperature (65 °C). Unfortunately, alkyl

J. Chem. Research (S), 1998, 598–599<sup>†</sup>

The result of our experiment indicated that as a small amount of water was added to the  $Sm-Me_3SiCl$  system, it not only accelerated the cleavage reaction, but also increased the yield of product. However, the mode of action is not fully understood and a more detailed study is in progress in our laboratory.

In conclusion, it has been found that the  $Sm-Me_3SiCl-H_2O$  reductive system is an effective one for cleaving the Se–Se bond in diphenyl diselenide at mild temperatures. The notable advantages of this reaction are neutral reaction conditions, simple operation and good yields.

## Experimental

<sup>1</sup>H NMR spectra were recorded on a JEOL PMX 60 SI instrument. All NMR samples were measured in CCl<sub>4</sub> using Me<sub>4</sub>Si as internal standard. IR spectra were obtained on a Perkin-Elmer 683 spectrophotometer as liquid films.

Metallic samarium and other chemicals were purchased from commercial sources and used without purification. Chlorotrimethylsilane was redistilled prior to use and kept under an inert atmosphere with molecular sieves. THF was freshly distilled from sodium-benzophenone ketyl prior to use.

General Procedure for the Preparation of Unsymmetrical Phenyl Selenides.—Under a nitrogen atmosphere, metallic samarium powder (0.15 g, 1.0 mmol) and diephenyl diselenide (0.16 g, 0.5 mmol) were placed in a three-necked reaction flask and Me<sub>3</sub>SiCl (1.0 ml) and THF (5 ml) were added in one portion. Then H<sub>2</sub>O (18  $\mu$ l) was added to the mixture and the resulting mixture was magnetically stirred for 3 h at 45 °C until the powdered samarium was almost consumed and the yellow solution had become almost colorless. To the mixture was added organic halide (1.2 mmol), (or epoxides, *etc.*) in THF (5 ml). When the reaction was complete, water (4 ml) was added to quench the reaction and the mixture was extracted with diethyl ether (2 × 20 ml). The extracts were washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed under reduced pressure; the residue was then purified by preparative TLC on silica gel to give pure product.

preparative TLC on silica gel to give pure product. Benzyl phenyl selenide.<sup>11</sup> Oil.  $\delta_{\rm H}$  3.95 (2 H, s, CH<sub>2</sub>Se), 7.05–7.45 (10 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3090, 2980, 1580, 1490, 1440, 1150, 940, 740, 690.

2-(Phenylseleno)acetophenone.<sup>4</sup> Oil.  $\delta_{\rm H}$  4.05 (2 H, s, CH<sub>2</sub>Se), 7.00–7.95 (10 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3085, 2960, 1682, 1600, 1590, 1480, 1410, 1275, 1180, 925, 735, 700, 680.

Ethyl 2-(phenylseleno)acetate.<sup>12</sup> Oil.  $\delta_{\rm H}$  1.12 (3 H, t, CH<sub>3</sub>), 3.29 (2 H, s, CH<sub>2</sub>Se), 3.96 (2 H, q, OCH<sub>2</sub>), 7.01–7.65 (5 H, m, ArH).  $\nu_{\rm max/cm^{-1}}$  3070, 2960, 1750, 1590, 1485, 1460, 1350, 1220, 1170, 935, 740, 690.

Butyl phenyl selenide.<sup>13</sup> Oil.  $\delta_{\rm H}$  0.80 (3 H, t, CH<sub>3</sub>), 1.00–1.85 [3 H, m, (CH<sub>2</sub>)<sub>2</sub>], 2.70 (2 H, t, CH<sub>2</sub>Se), 6.95–7.55 (5 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3050, 2950, 1590, 1470, 1420, 1370, 1220, 1115, 920, 735, 700.

Hexyl phenyl selenide.<sup>14</sup> Oil.  $\delta_{\rm H}$  0.80 (3 H, t, CH<sub>3</sub>), 1.00–1.95 [8 H, m, (CH<sub>2</sub>)<sub>4</sub>], 2.95 (2 H, t, CH<sub>2</sub>Se), 7.00–7.60 (5 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3040, 2940, 1580, 1480, 1420, 1380, 1230, 1100, 930, 740, 690.

Octyl phenyl selenide.<sup>15</sup> Oil.  $\delta_{\rm H}$  0.80 (3 H, t, CH<sub>3</sub>), 1.00–2.10 [12 H, m, (CH<sub>2</sub>)<sub>6</sub>], 3.20 (2 H, t, CH<sub>2</sub>Se), 7.02–7.70 (5 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3020, 2945, 2860, 1589, 1460, 1400, 1220, 1110, 1041, 928, 790, 740, 690.

2-(Phenylseleno)ethanol.<sup>16</sup> Oil.  $\delta_{\rm H}$  2.34 (1 H, s, OH, disappeared on adding D<sub>2</sub>O), 2.90 (2 H, t, CH<sub>2</sub>Se), 3.59 (2 H, t, CH<sub>2</sub>O), 7.00–7.54 (5 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3530, 3040, 2955, 2900, 1600, 1490, 1462, 925, 740, 700.

1-Chloro-3-(phenylseleno)propan-2-ol.<sup>17</sup> Oil.  $\delta_{\rm H}$  2.73 (1 H, s, OH, disappeared on adding D<sub>2</sub>O), 3.00 (2 H, d, CH<sub>2</sub>Se), 3.52 (2 H,

d, CH<sub>2</sub>Cl), 3.64–4.00 (1 H, m, CH), 7.03–7.49 (5 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3450, 2089, 3050, 2945, 2900, 1590, 1485, 1460, 1050, 1020, 940, 735, 700.

β-(Phenylseleno)propionitrile.<sup>18</sup> Oil.  $\delta_{\rm H}$  2.30–2.70 (2 H, m, CH<sub>2</sub>CN), 2.74–3.13 (2 H, m, CH<sub>2</sub>Se), 7.00–7.52 (5 H, m, ArH).  $\nu_{\rm max/cm^{-1}}$  3100, 3080, 2950, 2254, 1588, 1485, 1445, 1420, 1070, 1020, 932, 735, 685.

Methyl 3-(phenylseleno)propanate.<sup>15</sup> Oil.  $\delta_{\rm H}$  2.33–2.77 (2 H, m, CH<sub>2</sub>CO<sub>2</sub>R), 2.80–3.18 (2 H, m, CH<sub>2</sub>Se), 3.50 (3 H, s, OCH<sub>3</sub>), 6.97–7.53 (5 H, m, ArH).  $\nu_{\rm max}/{\rm cm}^{-1}$  3060, 2965, 1750, 1590, 1485, 1460, 1355, 1225, 1165, 1021, 935, 735, 690.

Methyl 2-methyl-3-(phenylseleno)propanate.<sup>19</sup> Oil.  $\delta_{\rm H}$  1.1–1.2 (3 H, d, CH<sub>3</sub>), 2.40–2.60 (2 H, m, CH), 2.65–3.20 (2 H, m, CH<sub>2</sub>Se), 3.54 (3 H, s, OCH<sub>3</sub>), 7.00–7.65 (5 H, m, ArH).  $\nu_{\rm max}$  /cm<sup>-1</sup> 3065, 2950, 1745, 1590, 1480, 1450, 1360, 1230, 1170, 930, 730, 690.

We thank the National Natural Science Foundation of China (Grant no. 29672007) and the Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, and the Chinese Academy of Sciences for financial supports.

Received, 1st May 1998; Accepted, 28th May 1998 Paper E/8/03305F

## References

- 1 The Chemistry of Organic Selenium and Tellurium Compounds, ed. S. Patai and Z. Rappoport, John Wiley, Chichester, 1986, vol. 1.
- 2 K. B. Sharpless and R. F. Laura, J. Am. Chem. Soc., 1973, 95, 2697.
- 3 E. E. Aynsley, N. N. Greenwood and J. B. Leach, *Chem. Ind.*, 1966, 379.
- 4 Y. Zhang, Y. Yu and R. Lin, Synth Commun., 1993, 23, 189.
- 5 H. Suzuki, M. Yoshinaga, K. Takaoka and Y. Hiroi, *Synthesis*, 1985, 497.
- 6 D. Liotta, U. Sunay, H. Santiesteban and W. Markiewicz, J. Org. Chem., 1981, 46, 2605; D. L. Klayman and W. H. H. Gunther, Organic Selenium Compounds: Their Chemistry and Biology, John Wiley, New York, 1973, p. 191.
- 7 F. Taboury, Bull. Soc. Chim. Fr., 1903, 29, 761.
- 8 H. B. Kagan and J. L. Namy, *Tetrahedron*, 1986, **42**, 6573; G. A. Molander, *Chem. Rev.*, 1992, **92**, 29; G. A. Molander, *Org. React.*, 1994, **46**, 211; G. A. Molander and C. R. Harris, *Chem. Rev.*, 1996, **96**, 307.
- 9 R. Yanada, N. Negoro, K. Yanada and T. Fujita, *Tetrahedron Lett.*, 1997, 38, 3271 and references cited therein.
- 10 N. Akane, Y. Kanagawa, Y. Nishiyama and Y. Ishii, *Chem. Lett.*, 1992, 2431; N. Akane, T. Hatano, H. Kusui, Y. Nishiyama and Y. Ishii, *J. Org. Chem.*, 1994, **59**, 7902.
- 11 V. F. Patel, G. Pattenden and D. M. Thompson, J. Chem. Soc., Perkin Trans. 1, 1990, 2729.
- 12 K. Haraguchi, H. Tanaka and T. Miyasaka, *Tetrahedron Lett.*, 1990, **31**, 227.
- 13 R. Raucher and G. A. Koolpe, J. Org. Chem., 1978, 43, 4252.
- 14 G. A. Russell, P. Ngoviwatchai, H. I. Tashtoush, A. Pla-Dalmau and R. K. Khanna, J. Am. Chem. Soc., 1988, 110, 3530.
- 15 S. V. Ley, I. A. O'Neil and C. M. R. Low, *Tetrahedron*, 1986, 42, 5363.
- 16 P. Rollin, V. V. Bencomo and P. Sinay, Synthesis, 1984, 134.
- 17 L. A. Khazemova and V. M. Al'bitskaya, Zh. Org. Khim., 1969, 5, 1926.
- 18 A. Ogawa, Y. Nishiyama, N. Kambe, S. Murai and N. Sonoda, *Tetrahedron Lett.*, 1987, 28, 3271.
- 19 E. G. Kataev and F. G. Gabdrakhmanav, Zh. Obshch. Khim., 1967, 37, 772.