View Article Online

ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: J. Bell, A. Harkiss, D. Nobis, E. Malcolm, A. Knuhtsen, C. Wellaway, A. G. Jamieson, S. Magennis and A. Sutherland, *Chem. Commun.*, 2020, DOI: 10.1039/C9CC09064A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the Information for Authors.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/chemcomm

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Dpen Access Article. Published on 13 January 2020. Downloaded on 1/13/2020 6:40:46 PM.

COMMUNICATION

Conformationally rigid pyrazoloquinazoline α -amino acids: oneand two-photon induced fluorescence

Received 00th January 20xx, Accepted 00th January 20xx Jonathan D. Bell,^a Alexander H. Harkiss,^a David Nobis,^a Eilidh Malcolm,^a Astrid Knuhtsen,^a Christopher R. Wellaway,^b Andrew G. Jamieson,^a Steven W. Magennis^a and Andrew Sutherland^{*a}

DOI: 10.1039/x0xx00000x

The synthesis and photophysical properties of a new class of α amino acid bearing a rigid pyrazoloquinazoline chromophore are described. Confromational constraint of the amino acid side-chains resulted in high emission quantum yields, while the demonstration of two-photon-induced fluorescence via near-IR excitation signifies their potential for sensitive bioimaging applications.

Fluorescence spectroscopy is a powerful technique for studying biological structure and function and for visualizing molecular interactions and intracellular processes.¹ This is due to its high sensitivity and information content, its multiplexing capability, the ability to probe processes over a wide time range, and the availability of an array of versatile fluorescent probes. The investigation of cellular functions such as enzyme mechanisms and protein-protein interactions have often utilized extrinsic fluorescent labels such as green fluorescent protein (GFP) and its derivatives² or intrinsic labels such as the proteinogenic fluorescent α -amino acids, tyrosine or tryptophan.³ However, the attachment of a fluorescent protein can alter the stability and functionality of the fusion partner, while tyrosine and tryptophan have poor optical properties and the occurrence of multiple residues in different environments can complicate their analysis.³

In recent years these limitations have been overcome by the development of unnatural fluorescent α -amino acids.⁴ The photophysical properties of such amino acids can be tuned for a particular application and they can be incorporated into proteins and peptides by solid phase peptide synthesis (SPPS) or by unnatural amino acid mutagenesis.⁵ Of particular interest are amino acids with side-chain chromophores that can be easily incorporated into peptides and proteins with minimal disruption to structure and function.^{4,6} While some of these fluorescent α -amino acids have found application in biological and medicinal imaging, an inherent issue of small chromophores is that ultraviolet (UV) light is required

for excitation. Prolonged exposure to UV light causes cell and tissue damage through photo-bleaching. In recent years, this limitation has been overcome by the use of two-photon fluorescence microscopy, which permits excitation of the chromophore at longer wavelength, allowing long-term imaging of biological specimens, deeper tissue penetration, with minimization of photo-bleaching and phototoxicity.⁷ While this technique has been utilized with a wide range of fluorophores and applied to the imaging of various biological processes and diseases,⁸ there are relatively few reports with unnatural α-amino acids. Examples include the seminal work of Mély and co-workers who used two-photon excitation of peptides containing 3-hydroxyflavone derived α -amino acids (e.g. 1, Fig. 1) for monitoring interactions with cell membranes and intracellular RNA,9 while the Vendrell group used two-photon excitation of a cyclic peptide bearing a tryptophan-BODIPY conjugate (2) for the visualization of fungal infections.¹⁰



Fig. 1 Selected Unnatural Fluorescent $\alpha\text{-Amino}$ Acids.

We previously reported the synthesis of 5-arylpyrazole-derived α -amino acids such as **3** (Fig. 1).¹¹ These compounds possessed some interesting photophysical properties however, the rotational flexibility between the aryl and pyrazole rings resulted in low quantum yields and brightness. As conformationally rigid, molecular structures permit more efficient π -overlap and subsequently brighter

^{a.} WestCHEM, School of Chemistry, The Joseph Black Building, University of

Glasgow, Glasgow, G12 8QQ, UK. Email: Andrew.Sutherland@glasgow.ac.uk ^{b.} Medicinal Chemistry, GlaxoSmithKline Medicines Research Centre, Gunnels Wood Road, Stevenage, SG1 2NY, UK.

⁺Electronic Supplementary Information (ESI) available: Experimental procedures, characterisation data for compounds, photophysical data for amino acids and, NMR spectra for all novel compounds. See DOI: 10.1039/x0xx00000x

COMMUNICATION

Open Access Article. Published on 13 January 2020. Downloaded on 1/13/2020 6:40:46 PM.

chromophores,¹² we proposed that conformational restriction of these 5-arylpyrazole motifs would lead to α -amino acids with enhanced photophysical properties. Here we describe a new class of α -amino acid (4), bearing planar pyrazologuinazoline chromophores. We also report the photophysical properties of these compounds, demonstrating how conformational restraint leads to fluorophores with high quantum yields and spectroscopic properties that are sensitive to environment polarity. We also show the application of these α -amino acids to two-photon excitation, for potential use in imaging UV sensitive, biological systems.

A synthetic route for the preparation of pyrazologuinazolinecontaining α -amino acids **4** was developed from L-aspartic acid. Initially, L-aspartic acid was converted to phosphononorvaline derivative 5 in five steps and 84% overall yield using previously described methods.^{13,14} A range of 2-nitrobenzaldehydes was reacted with phosphononorvaline 5 via a Horner-Wadsworth-Emmons reaction, which gave the E-isomer of enones 6a-e in 42-82% yields (Scheme 1).^{14,15} The enone moiety was used to construct the pyrazole ring by reaction of **6a-e** with hydrazine via a one-pot condensation/aza-Michael process. Low temperature oxidation of the resulting 2-pyrazolines with DDQ gave 5-(2-nitroaryl)pyrazoles 7a-e.11 Carbonylation was then achieved by reduction of the nitroaryl moiety, followed by reaction with triphosgene under basic conditions.¹⁶ This gave pyrazologuinazolines 8a-e in 30-75% yields over the two steps. Deprotection to the parent amino acids was either conducted in one step under acidic conditions (4b, 37%; 4c, 40%) or more efficiently by a two-step strategy involving ester hydrolysis, followed by removal of the Boc-protecting group under milder acidic conditions (4a, 80%; 4d, 82%; 4e, 78%).



č Scheme 1 Synthesis of pyrazologuinazoline α -amino acid 4a-e

н'n

4d

The UV/Vis absorption and emission spectra of the amino acids were recorded in methanol at a concentration of @15 30109END(Fig42 and Table 1).¹⁷ As expected for chromophores that are planar in the ground state, the amino acids, particularly 4a-d, showed structured absorption spectra with well-resolved vibrational bands (Fig. 2a).18 The emission spectra of amino acids 4a-e, following one-photon excitation at the longest wavelength absorption band, showed strong fluorescence ranging from 342-414 nm, with a vibrational progression which mirrored that of the absorption spectra (Fig. 2b).¹⁹ Amino acid 4e bearing the electron rich dimethyl-amino substituent gave a red-shifted, structureless emission maximum at 414 nm, which is indicative of a combination of locally excited and internal

charge transfer states.²⁰ More importantly, the quantum yields of 4a-e were characteristic of conformational restriction and showed a ten-fold increase in comparison to the non-rigid pyrazole-derived amino acid **3**. As a result, these α -amino acids are substantially brighter than the corresponding non-rigid systems.



Fig. 2 (a) Absorption spectra of amino acids 4a–4e recorded at 0.5×10^{-5} M in MeOH (b) Emission spectra of 4a-4e at 0.5 × 10⁻⁵ M in MeOH.

Table 1: Photophysical data for 5'-(naphthylen-2"-yl)-1'-phenylpyrazole amino acid 3 and
pyrazologuinazoline-derived α -amino acids 4a–e . ^{<i>a</i>}

Amino	λ_{Abs}	ε (cm ⁻¹ M ⁻¹)	λ_{Em}	$arPsi_{F}{}^{b}$	Brightness
acid	(nm)		(nm)		(cm ^{−1} M ^{−1}) ^c
3	249	23700	356	0.04	950
4a	321	8300	342	0.42	3500
4b	350	7700	372	0.47	3600
	(279)	(24,800)	(373)	(0.27)	(6700)
4c	328	10000	353	0.47	4700
4d	337	11200	363	0.56	6300
4e	349	12200	414	0.48	5900

^a All spectra were recorded at 0.5×10^{-5} M in MeOH. ^b Quantum yields ($\Phi_{\rm F}$) were determined in MeOH using anthracene and L-tryptophan as standards. ^c Brightness was calculated as the product of the QY and the molar absorptivity at the wavelength maximum specified.

Page 2 of 4

'n

4e

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Dpen Access Article. Published on 13 January 2020. Downloaded on 1/13/2020 6:40:46 PM.

COMMUNICATION

Journal Name

As amino acid **4e** showed the most interesting photophysical properties, this compound was further investigated. The fluorescence decay was measured at an emission wavelength of 430 nm in methanol using time-correlated single-photon counting, following excitation at 390 nm (see SI). Three decay components were fitted with lifetimes of 0.79, 2.17 and 7.90 ns, and A-factors of 0.60, 0.36 and 0.04, respectively. This suggests the molecule exists predominantly in two states, as has been observed previously for dimethylamino-substituted fluorophores.²¹ The amplitude-weighted average lifetime is 1.57 ns, which is comparable to fluorophores typically used in fluorescence lifetime imaging microscopy (FLIM).²²

To assess the environmental sensitivity of amino acid **4e**, its photophysical properties were examined in a range of solvents. The absorption maxima (342–348 nm) were found to be independent of polarity (see SI), indicating negligible intramolecular interaction between the electron-rich dimethylamino and electron-deficient *N*-acylpyrazole moieties in the ground state. In contrast, the emission spectra were found to be highly sensitive to the solvent used, with increasing polarity leading to concomitantly broadened, structureless emission spectra at longer wavelengths (Fig. 3a).^{23,24} For example, in THF, an emission maximum at 374 nm was observed, while in phosphate-buffered saline (PBS), the emission maximum was found at 424 nm. These spectral features and strong solvatochromism are likely due to increasing solvent stabilization of both the locally excited and intramolecular charge transfer states.

Application of amino acid **4e** for two-photon spectroscopy was next investigated. Two- and three-photon excitation has recently been shown to be a promising approach for the ultrasensitive analysis of fluorescent nucleobases, suggesting a similar approach may be applicable to α -amino acids.²⁵ Two-photon absorption was confirmed with a quadratic dependence of the emitted intensity on the excitation intensity (see SI). Furthermore, the molecule has no one-photon absorption in the near IR, precluding linear excitation. The emission spectra of amino acid **4e** in methanol after one- and two-photon excitation have a similar profile (Fig. 3b). The two-photon cross section was measured to be 0.38±0.06 GM using rhodamine B as a reference compound (see SI). This value is comparable to other molecules of similar size.^{8d,e} Thus, **4e** can undergo two-photon absorption using near-IR excitation, thereby avoiding the use of UV excitation and the issues of photobleaching.

The potential of these amino acids for incorporation into peptides via SPPS methodology was also investigated. A model pentapeptide (Val-Pro-Thr-Leu-Lys), based on the Bax-binding domain of Ku70, a multifunctional protein involved in DNA repair and cell-death regulation was chosen.²⁶ This pentapeptide and similar analogues have been shown to have low cytotoxicity and are effective at penetrating living cells.^{26b} The pentapeptide was prepared in an automated, peptide synthesizer using TentaGel[™] S Rink Amide (RAM) resin as the polymer support²⁷ and routine SPPS methodology (Scheme 2). Coupling of each Fmoc-protected amino acid was performed by HCTU²⁸ activation, followed by piperidinemediated N-deprotection, which gave the N-terminal unprotected pentapeptide. Boc-Protected pyrazoloquinazoline-derived amino acid 9 (see SI for preparation) was then coupled manually onto the polymer-supported pentapeptide. A TFA cleavage cocktail was used to remove the protecting groups and release the hexapeptide from the polymer support. Purification by reverse phase-HPLC gave hexapeptide 10 in 72% yield with >95% purity. Analysis of the photophysical properties of hexapeptide 10 showed that apart from a small hypsochromic shift in the emission spectra (Fig. 4), the key

photophysical properties of α -amino acid **4e** were retained including a high quantum yield (0.45) and brightness value (6900/cmc2vP4)4A



Fig. 3 (a) Solvatochromic study of **4e**. Spectra were recorded using a concentration of 0.5×10^{-5} M. (b) Emission of **4e** after two-photon absorption (red) and one-photon (blue) absorption in MeOH. For the two-photon measurement, a solution of 200 μ M in MeOH and an excitation centred around 700 nm with FWHM of 10 nm was used.



Scheme 2 SPPS and photophysical data of hexapeptide 10.



Fig. 4 Absorption (dotted lines) and emission (solid lines) spectra of amino acid 4e (blue) and hexapeptide 10 (red), recorded at 0.5×10^{-5} M in MeOH.

Journal Name

In summary, a new class of conformationally rigid, unnatural α amino acid has been synthesized using a highly regioselective olefination reaction to introduce side-chain diversity, followed by a one-pot condensation/aza-Michael reaction and carbonylation process to form the key pyrazologuinazoline ring system. Although containing relatively small side-chains, these compounds were found to be strongly fluorescent with high quantum yields (42-56%). The potential of amino acid 4e as a sensitive bio-imaging tool was demonstrated with successful excitation of this compound at 700 nm using two-photon spectroscopy. Furthermore, amino acid 4e was efficiently incorporated into a biologically relevant peptide as part of a SPPS process. Based on the exciting optical properties of this new class of α -amino acid, current work is investigating the introduction of these minimally invasive compounds into other biologically relevant peptides for in vitro and in vivo spectroscopic and microscopic applications.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Article. Published on 13 January 2020.

oen Access

Downloaded on 1/13/2020 6:40:46 PM.

Financial support from EPSRC (studentships to J.D.B., EP/N509176/1 and A.H.H., EP/K503058/1), University of Glasgow (studentship to D.N.) and GlaxoSmithKline is gratefully acknowledged.

Notes and references

- For some recent reviews, see: (a) L.-Y. Niu, Y.-Z. Chen, H.-R. Zheng, L.-Z. Wu, C.-H. Tung and Q.-Z. Yang, *Chem. Soc. Rev.*, 2015, **44**, 6143. (b) A. B. Chinen, C. M. Guan, J. R. Ferrer, S. N. Barnaby, T. J. Merkel and C. A. Mirkin, *Chem. Rev.*, 2015, **115**, 10530. (c) X. Chen, F. Wang, J. Y. Hyun, T. Wei, J. Qiang, X. Ren, I. Shin and J. Yoon, *Chem. Soc. Rev.*, 2016, **45**, 2976.
- (a) R. Y. Tsien, Annu. Rev. Biochem., 1998, 67, 509. (b) R. Heim,
 A. B. Cubitt and R. Y. Tsien, Nature, 1995, 373, 663.
- 3 R. W. Sinkeldam, N. J. Greco and Y. Tor, *Chem. Rev.*, 2010, **110**, 2579 and references therein.
- 4 For recent reviews, see: (a) A. T. Krueger and B. Imperiali, ChemBioChem., 2013, 14, 788. (b) R. Kubota and I. Hamachi, Chem. Soc. Rev., 2015, 44, 4454. (c) A. H. Harkiss and A. Sutherland, A. Org. Biomol. Chem., 2016, 14, 8911.
- For examples, see: (a) A. Chatterjee, H. Xiao, P.-Y. Yang, G. Soundararajan and P. G. Shultz, Angew. Chem. Int. Ed., 2013, 52, 5106. (b) S. Chen, N. E. Fahmi, L. Wang, C. Bhattacharya, S. J. Benkovic and S. M. Hecht, J. Am. Chem. Soc., 2013, 135, 12924. (c) L. C. Speight, A. K. Muthusamy, J. M. Goldberg, J. B. Warner, R. F. Wissner, T. S. Willi, B. F. Woodman, R. A. Mehl and E. J. Petersson, J. Am. Chem. Soc., 2013, 135, 18806.
- For recent examples, see: (a) P. Cheruku, J.-H. Huang, H.-J. Yen, R. S. Iyer, K. D. Rector, J. S. Martinez and H.-L. Wang, *Chem. Sci.*, 2015, **6**, 1150. (b) M. R. Hilaire, I. A. Ahmed, C.-W. Lin, H. Jo, W. F. DeGrado and F. Gai, *PNAS*, 2017, **114**, 6005. (c) M. Arribat, E. Rémond, S. Clément, A. Van Der Lee and F. Cavelier, *J. Am. Chem. Soc.*, 2018, **140**, 1028. (d) A. H. Harkiss, J. D. Bell, A. Knuhtsen, A. G. Jamieson and A. Sutherland, *J. Org. Chem.*, 2019, **84**, 2879.
- 7 A. Diaspro, G. Chirico and M. Collini, *Q. Rev. Biophys.*, 2005, 38, 97.
- 8 (a) R. Yuste and W. Denk, *Nature*, 1995, **375**, 682. (b) C. Xu, W.
 Zipfel, J. B. Shear, R. M. Williams and W. W. Webb, *PNAS*, 1996, **93**, 10763. (c) W. R. Zipfel, R. M. Williams, R. Christie, A.

Y. Nikitin, B. T. Hyman and W. W. Webb, *PNAS* 2003, **100**, 7075. (*d*) R. S. K. Lane and S. W. Magennis, *RSGADC* 2013, **4**, 11397. (*e*) R. S. K. Lane, R. Jones, R. W. Sinkeldam, Y. Tor and S. W. Magennis, *ChemPhysChem*, 2014, **15**, 867.

- 9 (a) V. Y. Postupalenko, O. M. Zamotaiev, V. V. Shvadchak, A. V. Strizhak, V. G. Pivovarenko, A. S. Klymchenko and Y. Mély, *Bioconjugate Chem.*, 2013, 24, 1998. (b) M. Sholokh, O. M. Zamotaiev, R. Das, V. Y. Postupalenko, L. Richert, D. Dujardin, O. A. Zaporozhets, V. G. Pivarenko, A. S. Klymenchenko and Y. Mély, *J. Phys. Chem. B*, 2015, 119, 2585.
- L. Mendive-Tapia, C. Zhao, A. R. Akram, S. Preciado, F. Alberico, M. Lee, A. Serrels, N. Kielland, N. D. Read, R. Lavilla and M. Vendrell, *Nat. Commun.*, 2016, 7, 10940.
- 11 L. Gilfillan, R. Artschwager, A. H. Harkiss, R. M. J. Liskamp and A. Sutherland, *Org. Biomol. Chem.*, 2015, **13**, 4514.
- 12 (a) M. Nepraš, N. Almonasy, F. Bureš, J. Kulhánek, M. Dvorák and M. Michl, *Dyes Pigm.*, 2011, **91**, 466. (b) M. S. Michie, R. Götz, C. Franke, M. Bowler, N. Kumari, V. Magidson, M. Levitus, J. Loncarek, M. Sauer and M. J. Schnermann, *J. Am. Chem. Soc.*, 2017, **139**, 12406. (c) J. C. Anderson, C.-H. Chang, A. P. Jathoul and A. J. Syed, *Tetrahedron*, 2019, **75**, 347.
- 13 D. E. Rudisill and J. P. Whitten, *Synthesis*, 1994, 851.
- (a) L. S. Fowler, D. Ellis and A. Sutherland, *Org. Biomol. Chem.*, 2009, **7**, 4309. (b) L. S. Fowler, L. H. Thomas, D. Ellis and A. Sutherland, *Chem. Commun.*, 2011, **47**, 6569.
- 15 F. Gosselin and W. D. Lubell, J. Org. Chem., 1998, 63, 7463.
- 16 F. Varano, D. Catarzi, V. Colotta, G. Filacchioni, A. Galli, C. Costagli and V. Carlà, J. Med. Chem., 2002, 45, 1035.
- 17 The absorption and emission spectra for each amino acid are presented in the Supporting Information.
- 18 (a) I. B. Berlman, J. Phys. Chem., 1970, 74, 3085. (b) N. I. Nijegorodov and W. S. Downey, J. Phys. Chem., 1994, 98, 5639.
- 19 It should be noted that while excitation of the absorption band at longest wavelength gave the most intense emission spectra for most of the amino acids, this was not the case for **4b**. Excitation of the peak at 279 nm gave the most intense fluorescence and highest brightness value (Table 1).
- 20 V. A. Galievsky, S. I. Druzhinin, A. Demeter, S. A. Kovalenko, T. Senyushkina, P. Mayer and K. A. Zachariasse, J. Phys. Chem. A, 2011, 115, 10823.
- 21 S. R. Meech and D. Phillips, J. Chem. Soc., Faraday Trans. 2, 1987, 83, 1941.
- 22 P. Sarder, D. Maji and S. Achilefu, *Bioconjugate Chem.*, 2015, 26, 963.
- 23 S. Achelle, I. Nouira, B. Pfaffinger, Y. Ramondenc, N. Plé and J. Rodríguez-López, J. Org. Chem., 2009, 74, 3711.
- 24 J. R. Lakowicz, Principles of Fluorescence Spectroscopy, Springer, New York, 2006.
- (a) D. Nobis, R. S. Fisher, M. Simmermacher, P. A. Hopkins, Y. Tor, A. C. Jones and S. W. Magennis, *J. Phys. Chem. Lett.*, 2019, 10, 5008. (b) R. S. Fisher, D. Nobis, A. F. Füchtbauer, M. Bood, M. Grøtli, L. M. Wilhelmsson, A. C. Jones and S. W. Magennis, *Phys. Chem. Chem. Phys.*, 2018, 20, 28487.
- 26 (a) J. A. Downs and S. P. Jackson, *Nat. Rev. Mol. Cell. Biol.*, 2004, **5**, 367. (b) J. A. Gomez, J. Chen, J. Ngo, D. Hajkova, I.-J. Yeh, V. Gama, M. Miyagi and S. Matsuyama, *Pharmaceuticals*, 2010, **3**, 3594.
- 27 W. Li and B. Yan, J. Org. Chem., 1998, 63, 4092.
- 28 HCTU: *O*-(1-*H*-6-Chlorobenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate.