

Baeyer–Villiger Reaction in the Solid State

Fumio Toda,* Minoru Yagi, and Koji Kiyoshige










Department of Industrial Chemistry, Faculty of Engineering, Ehime University, Matsuyama 790, Japan

Some Baeyer–Villiger oxidations of ketones with *m*-chloroperbenzoic acid proceed much faster in the solid state than in solution.

Relatively few organic reactions have been carried out in the solid state.¹ We have found that some Baeyer–Villiger oxidations of ketones with *m*-chloroperbenzoic acid (1) proceed much faster in the solid state than in solution.

The oxidations were carried out at room temperature with a mixture of powdered ketone and 2 mol. equiv. of powdered peroxy acid (1). When the reaction time was longer than 1 day, the reaction mixture was ground once a day with agate pestle

Table 1. Yields of Baeyer–Villiger oxidation products in the solid state and in CHCl₃^a.

		$\text{R}^1\text{COR}^2 \xrightarrow[m\text{-ClC}_6\text{H}_4\text{CO}_3\text{H}]{(1)} \text{R}^1\text{CO}_2\text{R}^2$		
Ketone	Reaction time	Product	Yield (%)	
			Solid state	CHCl ₃ ^b
 (2)	30 min	 (3)	95	94
 (4)	5 days	 (5)	64	50
PhCOCH ₂ Ph (6)	24 h	PhCO ₂ CH ₂ Ph (7)	97	46
PhCOPh (8)	24 h	PhCO ₂ Ph (9)	85	13
 (10)	24 h	 (11)	50	12
 (12)	4 days	 (13)	(1:1) 39	6
		 (14)		

^a Molar ratio of ketone and (1) is 1:2. ^b The reaction was carried out with 1 g of ketone in 50 ml of CHCl₃.

and mortar. The excess of peroxy acid (**1**) was decomposed with aqueous 20% NaHSO_3 , and the product was taken up in ether. The solution was washed with aqueous 20% NaHCO_3 and water, dried (Na_2SO_4), and evaporated. The crude product was chromatographed on silica gel (benzene- CHCl_3); yields are shown in Table 1. The ratio of (**13**) to (**14**) was determined by ^1H n.m.r.

For comparison, the oxidation was also carried out in CHCl_3 . Yields obtained from a solution of the ketone and 2 molar equiv. of peroxy acid (**1**) in CHCl_3 at room temperature are also summarized in Table 1.

In the case of ketones (**2**) and (**4**), there is no marked difference between reactions in the solid state and in CHCl_3 . However, there is a large difference with the ketones (**6**), (**8**), (**10**), and (**12**) (Table 1). It is surprising that the reaction occurs more easily in the solid state than in solution. Since Baeyer-Villiger reaction in solution usually takes a long time,²

a reaction which proceeds faster and does not need solvent has many advantages.

We have shown that the movement of molecules in host-guest complex formation in the solid state is easy. Some host-guest complexes can be formed by mixing host and guest compounds in the solid state or by keeping a mixture of powdered host and guest at room temperature.³ Therefore, the ease of oxidation in the solid state is not unexpected.

Received, 25th January 1988; Com. 8/00271A

References

- 1 'Organic Solid State Chemistry,' ed. G. R. Desiraju, Elsevier, Amsterdam, 1987.
- 2 S. L. Friess and A. H. Soloway, *J. Am. Chem. Soc.*, 1951, **73**, 3968.
- 3 F. Toda, K. Tanaka, and A. Sekikawa, *J. Chem. Soc., Chem. Commun.*, 1987, 279.