

View Article Online View Journal

ChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: X. zhu, M. xiong, H. Liu, G. Mao, L. Zhou, J. Zhang, X. hu, X. Zhang and W. Tan, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC08695G.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

Journal Name



A FRET-based ratiometric two-photon fluorescent probe for dual channel imaging of nitroxyl in living cells and tissues

Received 00th January 20xx, Accepted 00th January 20xx

Xiaoyan Zhu[†]a, Mengyi Xiong^{†a}, Hong-wen Liu^a, Guo-jiang Mao^b, Liyi Zhou^a, Jing Zhang^a, Xiaoxiao Hu^a, Xiao-Bing Zhang^{a*} and Weihong Tan^{a*}

www.rsc.org/

A FRET-based two-photon fluorescent probe, P-Np-Rhod, which exhibited a fast and high selective ratiometric response to nitroxyl, was first proposed. P-Np-Rhod was successfully applied to twophoton dual-channel imaging of nitroxyl in living cells and tissues with less cross-talk between channels and satisfactory deep-tissue imaging depth.

Nitroxyl (HNO), the one-electron reduced form of NO, has recently been found to possess unique biological activity and pharmacology applications.¹ For example, HNO can inhibit the activity of aldehyde dehydrogenase by interacting with its protein thiols.² And it has been demonstrated to exhibit the potential for treating heart failure through upregulating the calcitonin gene-related peptide which can induce vasorelaxation.³ Moreover, HNO could also be used to treat alcoholism, mitigate ischemia-reperfusion injury and functionalize as a potential anti-cancer drug.⁴ In this light, the development of a reliable assay method that can monitor HNO in biosystems is highly desired.

Until recently, several analytical methods including colorimetric approach,⁵ electron paramagnetic resonance spectroscopy,⁶ high performance liquid chromatography,⁷ mass spectrometry⁸ and electrochemical analysis⁹ have been developed for HNO detection. In despite of appropriate sensitivity, these methods are unsuitable for detection of HNO in living systems, due to the complicated pretreatment and potential destruction of biological samples. In recent years, fluorescent probes have been widely applied for bioimaging of various analytes in living cells, tissues and organisms by virtue of their high sensitivity, fast response time, and little biodamage.¹⁰ Recently, based on the reaction of HNO with triarylphosphine, various fluorescent probes were rationally designed to discriminate HNO from other biological reductants.¹¹

Nevertheless, the biological applications of these fluorescent probeare limited by one-photon excitation with short wavelengths, when could lead to photobleaching of probes, restricted tissue-penetration depth, and high cellular autofluorescence. In sharp contrast, ..., photon microscopy (TPM) approach, which excites a two-photo. (TP) fluorophore with near-infrared (NIR) laser pulses, car overcome these shortages and provide improved three-dimension. spatial localization with reduced photodamage to biosamples and increased tissue penetration depth.¹² Despite all these advantages only few TP fluorescent turn-on probe has been reported for the detection of HNO.^{11k} Moreover, this turn-on probe was limited for practical applications, due to the fact that the fluorescence intensi measurement within a single detection window may vary with the experimental conditions, such as the incident laser power and t e distribution of probes. Ratiometric probes, by measuring the changes of the intensity ratio at two different emission peaks, can provide a built-in correction to minimize these interferences, resulting in a more effective system for imaging living cells and tissues.¹³ It worth pointing out that, although several one-photon excut ratiometric fluorescent probes have been developed for the detection of HNO^{11d, e, g, j}, to our best knowledge, no ratiometric TP fluorescent probes for detecting HNO has been reported to date.

In this work, by combining the advantages of TPM technique win ratiometric probe, we designed and synthesized, for the first time, a ratiometric TP fluorescent probe (P-Np-Rhod, Scheme 1) for HN y based on a FRET strategy. In this FRET system, a TP naphthale derivative served as the energy donor and a rhodol fluorophore w.... chosen as the energy acceptor which was modified with (diphenylphosphino)-benzoate moiety as a recognition unit for HNC In the absence of HNO, the rhodol existed in a non-fluorescent lactone form and the FRET is off. As reported by previous literal res, HNO can react with 2-(diphenylphosphino)-benzoate moiety to give the corresponding aza-ylide, which can attack the adjacent est . linker in an intramolecular manner to release hydroxyl. And then the closed spirolactone form was converted to a conjugated fluoresce xanthene form to induce the occurrence of FRET (Scheme S2, ESI*) The sensing process was confirmed by both NMR spectroscopy and mass spectrometry (Fig. S9 and Fig. S10, ESI[†]). The detaile synthesis procedures and characterization of all the new compound

^aMolecular Science and Biomedicine Laboratory, State Key Laboratory of Chemo/Biosensing and Chemometrics, College of Chemistry and Chemical Engineering, Hunan University, Changsha 410082, China. E-mail: <u>xbzhanq@hnu.edu.cn</u>: <u>tan@chem.ufl.edu</u>.

^bSchool of Chemistry and Chemical Engineering, Henan ; Normal University, Xinxia ng, Henan 453007, P. R. China.

⁺ Co-first authors

 $[\]label{eq:electronic Supplementary Information (ESI) available: Synthesis, experimental details and characterization of new compounds. See DOI: 10.1039/x0xx00000x$

are described in the ESI[†].



Scheme 1 Chemical structure and the response mechanism of probe **P-Np-Rhodol** to HNO.

The spectroscopic properties of the probe P-Np-Rhod were firstly investigated in buffered (10 mM PBS, pH = 7.4) aqueous CH₃CH₂OH solution (H₂O/CH₃CH₂OH = 1:1, v/v). In the absence of HNO, the free probe P-Np-Rhod displayed a cyan-colored fluorescence with an emission peak at 448 nm (Fig. 1). After treatment with enhanced concentrations of AS (from 0 to 100 µM, Angeli's salt (AS) acted as the source of HNO in our experiments), a new fluorescence emission peak at 541 nm appeared and increased gradually, while the donor's characteristic emission peak (448 nm) showed a decreased tendency (Fig. 1), allowing a ratiometric fluorescence response for the detection of HNO. It is noteworthy that the large wavelength difference (93 nm) between the two emissions leads to two well-resolved emission bands for the probe, which is favourable for dual-channel imaging of HNO in biological samples with less cross-talk observed. We proceeded to investigate the UVabsorption properties of the probe. Upon addition of AS, no obvious absorption changes of the donor occurred, while a new absorption peak (at 511 nm) belonging to the conjugated xanthene form of the acceptor increased with an AS dose-dependent pattern. Meanwhile, the solution changed from colorless to orange, allowing colorimetric detection of HNO by naked eyes (Fig. S1, ESI⁺). The remarkable changes in absorption and fluorescence spectra might be attributed to the HNO-induced occurrence of FRET between the donor naphthalene and acceptor rhodol. The Förster energy transfer efficiency between the donor and acceptor rhodol was calculated to be 88.6% (Fig. S2, ESI⁺). The fluorescence intensity ratio value between the two wavelengths (I541 nm/I448 nm) varied from 0.13 to 4.67 with the concentration of HNO changed from 0 to 100 µM (Fig. 1). The detection limit for HNO was calculated to be $5.9{\times}10^{-7}~M$ (3o/slope, Fig. S3, ESI⁺), indicating that the probe P-Np-Rhod exhibited a high sensitivity for HNO.



Fig. 1 Fluorescence emission spectra of probe **P-Np-Rhod** (15 μ M) in the presence of various concentrations of AS (0-100 μ M) in PBS buffered (10 mM, pH = 7.4) aqueous EtOH solution (1:1, v/v). Inset:

The fluorescence intensity ratio (I_{541 nm}/I_{448 nm}) of prob₆ P₂Np₂R)_{Mne} (15 μM) versus AS concentration after incubation 43346 probe WFG

AS for 30 min. $\lambda_{ex} = 370$ nm. The time-dependent fluorescence response of probe **P-Np-R** (C) to different concentrations of AS (5, 35, 55 or 100 μ M) we examined by monitoring the change of fluorescent intensity ratio between the two wavelengths (I_{541 nm}/I_{448 nm}) after the addition of A S in PBS buffer. As shown in Fig. S4 (ESI[†]), the fluorescent intensity ratio (I_{541 nm}/I_{448 nm}) obviously increased and reached a plateau t around 25 min after the addition of AS, demonstrated that the probe **P-Np-Rhod** can provide a rapid response for HNO.

A highly selective response to the target molecule over other potentially competing molecules is of great significance for bioimaging probe with potential application in the complebiological system. Therefore, the probe **P-Np-Rhod** was treated with various biologically relevant analytes to evaluate its selectivit, including reactive oxygen species, reactive nitrogen specreducing agents, small molecule thiols, and biologically related metal ions. As shown in Fig. 2, no obvious changes of fluorescence intensity ratio (I_{541 nm}/I_{448 nm}) were observed in the presence of these interference species. In contrast, the introduction of AS could trigger remarkable change of the fluorescence intensity ratio at I_{541 nm}/I_{448 nm}. This result indicated that the probe **P-Np-Rhod** shows a high selectivity for HNO and possesses the ability f **r** detecting HNO in complex biological samples.



Fig. 2 Fluorescence response of the probe **P-Np–Rhod** (15 μÅ) toward various analytes in PBS buffered (10 mM, pH = 7.4) aqueous EtOH solution (1:1, v/v) at room temperature for 30 min. The fluorescence intensity ratio (I_{541 nm/I448 nm}) was plotted vers s substances: 1. Probe only (15 μM); 2-12. Ca²⁺, Ascorbic Acid, NO₂, N₃⁻, S²⁻, K⁺, Zn²⁺, NO₃⁻, Na⁺, Fe²⁺, Mg²⁺ (1 mM); 13. Fe³⁺ (100 μM, 14-15. Cys, GSH (3 mM); 16-23. H₂O₂, ClO⁻, NO, O₂⁻, •OH, TBH₁ •'OBu, ONOO⁻ (500 μM); 24. AS (100 μM). λ_{ex} = 370 nm.

The influence of pH on the fluorescence response of **P**- [**p**-**Rhod** to HNO was further investigated. As shown in Fig. So (ESI[†]), in the absence of HNO, the probe **P-Np-Rhod** was p **1** insensitive over a pH range of 5.0-10.0, indicating that the est **1** group was stable in this pH range. Upon addition of AS, the maximal fluorescence intensity ratio ($I_{541 \text{ nm}}/I_{448 \text{ nm}}$) appeared **i** the pH range of 6.0-7.8, which demonstrated that probe **P-Np**-**Rhod** can be employed to detect HNO in physiological system . With the pH value larger than 8.0, fluorescence intensity **1**

Journal Name

Journal Name

 $(I_{541\ nm}/I_{448\ nm})$ showed a decreased tendency due to the decreased decomposition rate of AS into nitroxyl under the strong basic conditions. 14

The TP active cross-section ($\delta\Phi$) value of **P-Np-Rhod** and **Np-Rhod** were respectively estimated to be 48 GM and 27 GM at 740 nm (Fig. S6, ESI[†]), indicating high capability of the probe **P-Np-Rhod** for detecting HNO with TP mode. In the absence of HNO, **P-Np-Rhod** showed a fluorescence emission peak at 448 nm under excitation with 740 nm femtosecond pulses. However, a new fluorescence emission peak at 541 nm appeared while the donor's emission peak at 448 nm vanished in the presence of HNO (Fig. 3). The probe **P-Np-Rhod** was demonstrated to possess potential ability for ratiometric TP fluorescent imaging of HNO.



Fig. 3 TP excited emission spectra of **P-Np-Rhod** (black line) and **Np-Rhod** (red line). TP excitation at 740 nm in PBS/EtOH (1:1, v/v, 10 mM, pH=7.4).

Encouraged by the remarkable features of the probe P-Np-Rhod including high selectivity and sensitivity, rapid response and good TP fluorescence properties, we next investigated whether this probe can be applied in TP excited dual-channel imaging of HNO in living cells. MTT assays were first performed to evaluate the potential cytotoxicity of P-Np-Rhod and Np-Rhod at various concentrations against HeLa cells. The results indicate that both P-Np-Rhod and Np-Rhod have no obvious cytotoxicity to live cells under the experimental conditions. (Fig. S7, ESI[†]). We then examined the applicability of the probe for detecting HNO in living cells, with results shown in Fig. 4. The HeLa cells incubated with only the probe P-Np-Rhod (15 µM) displayed a strong TP fluorescence signal in the cyan channel but weak fluorescence signal in the yellow channel, corresponding to strong fluorescence of the donor and weak fluorescence of the acceptor due to its spirolactone form. In contrast, HeLa cells incubated with P-Np-Rhod (15 µM) for 30 min, and then treated with AS (100 µM or 200 µM) exhibited a significant fluorescence enhancement in the yellow channel and a concomitant decrease in cyan channel, corresponding to strong fluorescence from the open-ring acceptor and the weak fluorescence from the donor due to the effective FRET process triggered by HNO. All above results showed that P-Np-Rhod was cell membrane permeable and could be successfully applied to TP-excited ratiometric dualchannel imaging of HNO in living cells.

To further investigate the applications of P-Np-Rhod fluorescence imaging of living samples, 19P0 32xe5xe60 3485G channel imaging of P-Np-Rhod for HNO in rat liver tissu slices was then performed (Fig. 5). The changes of fluorescent signal with scan depth were recorded by TPM in the Z-sca. mode. In the absence of AS, the tissue slices incubated with only P-Np-Rhod exhibited obvious fluorescence at the cyt channel with an imaging depths of 40-150 µm. In contrast, when the tissue slices were incubated with P-Np-Rhod (5) μ M) for 60 min, and then treated with 1 mM AS for another 60 min, relatively significant fluorescence was observed at the yellow channel with an imaging depths of 20-130 µn, followed with a remarkable decrease at the cyan channel. The results indicated that P-Np-Rhod possesses excellent tissu penetration and staining ability as well as ratiometric dua' colour (cyan and yellow) imaging performance with less cros. talk between dual emission channels.



Fig. 4 TP fluorescence images of HNO in live HeLa cells with **P**-**Np-Rhod.** (a) TP fluorescence images HeLa cells incubated with only probe **P-Np-Rhod** (15 μ M) for 30 min in the cyan channel; (b, images of (a) in the yellow channel; (c) Merged images of (a) and (b); (d, g) TP fluorescence images of HeLa cells incubated with. **Np-Rhod** (15 μ M) for 30 min, then with AS (100 μ M or 200 μ M) for 30 min in the cyan channel; (e, h) TP fluorescence images of (d, g) in the yellow channel; (f) Merged images of (d) and (e); () Merged images of (g) and (h). $\lambda_{ex} = 740$ nm, cyan channel: $\lambda_{en} = 470-530$ nm; yellow channel 550-650 nm. Scale bar: 50 μ m.

8	b
2 90 00 μm	Z 90.00 µm
c	d
Z 90.00 um	2 90.00 um

Fig. 5 TP fluorescence imaging of a rat liver frozen slice stained with **P-Np-Rhod** (50 μ M) at ~90 μ m for 60 min (a, b), followed 1 treatment with AS (1 mM) and incubated for another 60 min (c

Journal Name

The images were collected at 450-530 nm (cyan channel, a, c) and 540-650 nm (yellow channel, b, d) upon excitation at 740 nm with femtosecond pulses. Scale bars: $200 \ \mu m$

In summary, we have developed a ratiometric TP fluorescent probe **P-Np-Rhod** for detecting HNO for the first time. The diphenylphosphinobenzoyl group was employed as a recognition unit and a FRET cassette, termed **Np-Rhod**, acted as the fluorescent reporter. This ratiometric probe **P-Np-Rhod** can rapidly measure HNO with large emission shift (93 nm), high sensitivity and high selectivity. More importantly, **P-Np-Rhod** could be succesfully applied for TP fluorescence imaging of HNO in living cells and tissues with less cross-talk between dual emission channels. Far more than a promising tool to investigate the biological functions of HNO, this FRET-based TP strategy might also be used to design dual-channel imaging probes for other biological researches.

This work was supported by the National Key Scientific Program of China (2011CB911000), the National Key Basic Research Program of China (2013CB932702), NSFC (Grants 21325520, 21327009, J1210040, 21177036), the Foundation for Innovative Research Groups of NSFC (Grant 21221003), National Instrumentation Program (2011YQ030124). National Natural Science Foundation of China (21505032). Key scientific research project of higher education of the Henan province (15A150016)

notes and references

COMMUNICATION

- (a) F. Doctorovich, D. Bikiel, J. Pellegrino, S. A. Suárez, A. Larsen and M. A. Martí, *Coord. Chem. Rev.*, 2011, **255**, 2764; (b) J. C. Irvine, R. H. Ritchie, J. L. Favaloro, K. L. Andrews, R. E. Widdop and B. K. Kemp-Harper, *Trends Pharmacol. Sci.*, 2008, **29**, 601; (c) K. M. Miranda, *Coord. Chem. Rev.*, 2005, **249**, 433.
- E. G. DeMaster, B. Redfern and H. T. Nagasawa, Biochem. pharmacol., 1998, 55, 2007.
- N. Paolocci, T. Katori, H. C. Champion, M. E. St John, K. M. Miranda, J. M. Fukuto, D. A. Wink and D. A. Kass, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 5537.
- (a) A. J. Norris, M. R. Sartippour, M. Lu, T. Park, J. Y. Rao, M. I. Jackson, J. M. Fukuto and M. N. Brooks, *Int. J. Cancer*, 2008, **122**, 1905; (b) P. Pagliaro, D. Mancardi, R. Rastaldo, C. Penna, D. Gattullo, K. M. Miranda, M. Feelisch, D. A. Wink, D. A. Kass and N. Paolocci, *Free Radical Biol. Med.*, 2003, **34**, 33; (c) N. Paolocci, M. I. Jackson, B. E. Lopez, K. Miranda, C. G. Tocchetti, D. A. Wink, A. J. Hobbs and J. M. Fukuto, *Pharmacol. Ther.*, 2007, **113**, 442.
- (a) S. E. Bari, M. A. Marti, V. T. Amorebieta, D. A. Estrin and F. Doctorovich, *J. Am. Chem. Soc.*, 2003, **125**, 15272; (b) K. P. Dobmeier, D. A. Riccio and M. H. Schoenfisch, *Anal. Chem.*, 2008, **80**, 1247; (c) M. A. Marti, S. E. Bari, D. A. Estrin and F. Doctorovich, *J. Am. Chem. Soc.*, 2005, **127**, 4680.
- (a) U. Samuni, Y. Samuni and S. Goldstein, J. Am. Chem. Soc., 2010, 132, 8428; (b) Y. Xia, A. Cardounel, A. F. Vanin and J. L. Zweier, Free Radical Biol. Med., 2000, 29, 793.
- S. Donzelli, M. G. Espey, D. D. Thomas, D. Mancardi, C. G. Tocchetti, L. A. Ridnour, N. Paolocci, S. B. King, K. M. Miranda, G. Lazzarino, J. M. Fukuto and D. A. Wink, *Free Radical Biol. Med.*, 2006, 40, 1056.
- M. R. Cline, C. Tu, D. N. Silverman and J. P. Toscano, *Free Radical Biol. Med.*, 2011, **50**, 1274.

- 9. (a) S. A. Suarez, D. E. Bikiel, D. E. Wetzler, M. Avie Martician Doctorovich, Anal. Chem., 2013, 85, 10262;1(b) (Searchine Doctorovich, Anal. Chem., 2013, 85, 10262;1(b) (Searchine Doctorovich, Chem., 2010, A. Salvarezza, M. A. Marti and F. Doctorovich, Inorg. Chem., 2010, 49, 6955.
- (a) J. Fan, M. Hu, P. Zhan and X. Peng, *Chem. Soc. Rev.*, 2013, 4
 29; (b) J.-T. Hou, M.-Y. Wu, K. Li, J. Yang, K.-K. Yu, Y.-M. Xie an
 X.-Q. Yu, *Chem. Commun.*, 2014, **50**, 8640; (c) X. Li, X. Gao, V
 Shi and H. Ma, *Chem. Rev.*, 2014, **114**, 590; (d) L. Yuan, W. Lin, K.
 Zheng, L. He and W. Huang, *Chem. Soc. Rev.*, 2013, **42**, 622.
- 11. (a) K. N. Bobba, Y. Zhou, L. E. Guo, T. N. Zang, J. F. Zhang and S. Bhuniya, RSC Adv., 2015, 5, 84543; (b) X. Jing, F. Yu and L. Chen, Chem. Commun., 2014, 50, 14253; (c) K. Kawai, N. Ieda, ... Aizawa, T. Suzuki, N. Miyata and H. Nakagawa, J. Am. Chem. So 2013, 135, 12690; (d) C. Liu, Z. Cao, Z. Wang, P. Jia, J. Liu, 🧏 Wang, B. Han, X. Huang, X. Li, B. Zhu and X. Zhang, Sens. Actuators, B, 2015, 220, 727; (e) C. Liu, H. Wu, Z. Wang, C. Sha B. Zhu and X. Zhang, Chem. Commun., 2014, 50, 6013; (f) P. Liu, X. Jing, F. Yu, C. Lv and L. Chen, Analyst, 2015, 140, 4576; (g M. Lv, Y. Chen, J. Lei, C.-T. Au and S.-F. Yin, Anal. Methods, 2015 7, 3883; (h) G.-J. Mao, X.-B. Zhang, X.-L. Shi, H.-W. Liu, Y.-X. L.-Y. Zhou, W. Tan and R.-Q. Yu, Chem. Commun., 2014, 50, 5790; (i) Z. Miao, J. A. Reisz, S. M. Mitroka, J. Pan, M. Xian ar B. King, Bioorg. Med. Chem. Lett., 2015, 25, 16; (j) H. Zhang, Liu, Y. Tan, W. H. Xie, H. Lei, H. Y. Cheung and H. Sun, ACS Appl. Mater. Interfaces, 2015, 7, 5438; (k) K. Zheng, W. Lin, D. Chen, H. Chen, Y. Liu and K. Liu, Chem. Commun., 2015, 51, 5754.
- (a) D. Kim, H. G. Ryu and K. H. Ahn, Org. Biomol. Chem., 20
 12, 4550; (b) H. M. Kim and B. R. Cho, Acc. Chem. Res., 2009, 4
 863; (c) H. M. Kim and B. R. Cho, Chem. Rev., 2015, 115, 5014.
- M. H. Lee, J. S. Kim and J. L. Sessler, Chem. Soc. Rev., 2015, 44, 4185.
- A. S. Dutton, J. M. Fukuto and K. N. Houk, J. Am. Chem. Soc., 2004, 126, 3795.

4 | J. Name., 2012, 00, 1-3