

Tetrahedron Letters 40 (1999) 1941-1944

TETRAHEDRON LETTERS

## Synthesis of longithorone B, a sixteen-membered farnesylated *p*-benzoquinone<sup>1</sup>

Tadahiro Kato,\* Kentaro Nagae, and Masahiro Hoshikawa

Department of Chemistry, Faculty of Science, Science University of Tokyo, Kagurazaka 1-3, Shinjuku ku, Tokyo 162-8601. Japan

Received 16 November 1998; accepted 28 December 1998

**Abstract**: The intramolecular coupling reactions at the C-1 position of the 10Z-farnesyl moiety appended at the 2'-position of 4'-methoxy phenol derivatives were examined with the aid of  $Hf(OTf)_4$  and  $LiClO_4$ . When the phenol was protected with a TBDMS group, meta coupling predominated while para coupling occurred efficiently when protected with a pivaloyl group. The para-coupled product was transformed to longithorone B, recently isolated from a tunicate. © 1999 Elsevier Science Ltd. All rights reserved.

Recently, a group of farnesylated quinones featuring new macrocyclic skeletons has been isolated from a tunicate, *Aplidium longithorax* by Schmitz and his coworkers.<sup>2</sup> Among the quinones characterized, longithorone **B**(1) is a representative compound containing a 16-membered carbocycle comprised of benzoquinone bridged across the para position by a farnesyl unit having two trans and one cis double bonds. The unique structural features of 1, which is shown to have an atropisomerism, prompted us to explore an efficient synthetic route to longithorone B. Here we report the preparation of the compounds corresponding to 1 and its position isomer, 2 in *dl*-forms.



An 8:1 mixture of ethyl farnesate (3) was selected as our starting material, which was prepared by Horner-Emmons reaction of *trans* geranyl acetone with diethyl phosphonoacetate and NaH (80% yield). By treatment of 3 with  $SO_2Cl_2$  and  $Na_2CO_{3^3}$  the terminal double bond was selectively chlorinated to give an 8:1 mixture of 4a and 4b<sup>4</sup> in 95% yield, which was separated easily by usual silica gel column chromatography. The formation of phenyl ether proceeded smoothly when the allyl chloride (4a) was treated with *p*-MeOC<sub>6</sub>H<sub>4</sub>OK in the presence of 18-crown-6, affording 5 in 62% yield (81% conversion yield).

In the Claisen rearrangement of 5, the formation ratio of 6a and 6b depended largely on the reaction

1942

conditions as summarized in Table 1. Reaction of 5 in refluxing xylene for 23 h without any additive (run 1) resulted in a 1:4 ratio of **6a** and **6b** while exclusive formation of **6b** was observed when the Claisen rearrangement was carried out in the presence of Na, CO, in refluxing xylene (run 2). These results suggested that the phenolic proton (M = HOPh) of the initially formed phenol derivative interacts with the oxygen atom of the phenyl ether (5) through the hydrogen bonding as depicted in figure 1. In the transition states, the energy difference between Ts-E and Ts-Z may become smaller to permit the formation of the Z isomer (6a) in addition to the predominant E isomer (6b). In the presence of Na, CO, the Claisen rearrangement proceeded exclusively through the energetically more stable Ts-E, in which no interaction between the oxygen atom of the phenyl ether and the phenoxide ion (M = OPh) may occur. The slower reaction in the presence of Na<sub>2</sub>CO<sub>3</sub> as compared with that in run 1, in which the Claisen rearrangement is accelerated by the hydrogen bonding, may support the present suggestion.<sup>5</sup> Based on these considerations, the effects of other additives capable of interacting with the oxygen atom in the transition state were explored and we found that the Claisen rearrangement of the phenyl ether (5) proceeded in the presence of silica gel (run 3) or activated Al<sub>2</sub>O<sub>2</sub> (run 4) in refluxing xylene. The latter conditions furnished a 4.3:1 mixture of 6a and 6b in 71% yield after heating for 36 h. In the transition state, the oxygen atom of the ether linkage may interact with the aluminum atom of Al<sub>2</sub>O<sub>4</sub> to make the complexes, in which the amount of Ts-E decreases due to the repulsion between M (Al<sub>2</sub>O<sub>3</sub>) and the equatorial R group. The lack of such repulsion in Ts-Z may bring about the predominant formation of 6a.<sup>6</sup> Both isomers (6a and 6b) were separated by HPLC and the geometry of the double bonds in each compound was deduced from <sup>13</sup>C chemical shifts of the olefinic methyl groups.7



Run	Additive	Solv.	Temp. (°C)	Time (h)	5 : (6a+6b) (%) 6a : 6b	
1	-	xylene	140	23	28 : 72	1:4
2	Na <sub>2</sub> CO <sub>3</sub>	xylene	140	24	51 : <b>49</b>	0:1
3	SiO <sub>2</sub>	-	100	24	43 : 57	3:1
4	Al <sub>2</sub> O <sub>3</sub>	xylene	140	36	29:71	4.3 : 1

 Table.1
 Effect of reaction conditions on Claisen rearrangement of 5



Figure 1 Possible transition states of Claisen rearrangement

The intramolecular coupling reaction was examined using three types of protected allyl alcohols (7a, 7b and 7c), which were prepared as follows. Protection of 6a with TBDMS (TBDMSCl, imidazole, DMAP in DMF) or Bn (BnBr, NaH, TBAI in THF) groups followed by DIBAL-H reduction afforded the allyl alcohols 7a (73 %) and 7b (81 %), respectively. The pivalate (7c) was directly prepared from 7a in 51% yield by treatment with PivCl and TBAF in THF.<sup>8</sup> When 7a was treated with Hf(OTf), in the presence of LiClO, in CH<sub>2</sub>Cl<sub>2</sub>,<sup>9</sup> a mixture of 8a, 8b, and unidentified products (R = TBDMS) were obtained. Due to the difficulty of separation, the mixture was converted to the corresponding pivalates (8a and 8b, R = Piv), from which the pivalate (8b) was isolated in 64% yield by HPLC separation.<sup>12</sup> When the pivalate (7c) was subjected to the coupling reaction under the same conditions, 8a (R = Piv) was isolated in 78 % yield.<sup>13</sup> The pivalate (8a, R = Piv) was transformed to 1 by the sequential reactions of hydrolysis with KOH/MeOH followed by CAN oxidation in the presence of  $C_{s}H_{3}NO(CO_{2}H)_{2}$  in 58% yield.<sup>14</sup> The meta-coupled product (8b) was similarly transformed to the quinone (2). The predominant meta coupling of the TBDMS ether (7a) may be caused by the electron-donating nature of the TBDMS group overcoming its steric hindrance. In the meantime, steric hindrance as well as electron-withdrawing nature of pivaloyl group of 7c controlled the coupling reaction, leading to the para-coupled 8a as the major product. The ortho coupling product corresponding to 8c was not characterized from either 7a or 7c.<sup>15</sup> When the sequential reactions were applied to benzyl ether (7b), a complex mixture was detected without formation of any predominant product, suggesting that the benzyl group, in addition to the *p*-hydroquinone skeleton, underwent the coupling reaction.

The <sup>1</sup>H and <sup>13</sup>C chemical shifts and the coupling modes of the synthesized product (1) agree approximately with the reported values.<sup>16</sup> Thus, we have developed an efficient route for the construction of the macrocyclic farnesylated quinone derivatives (1 and 2).



## **Footnotes and References**

- \* E-mail: tkato@ch.kagu.sut.ac.jp (t.kato).
- 1. This constitutes part 58 of the series of cyclization of polyenes. Part 57, T. Kato, M. Tanaka, M. Hoshikawa and M. Yagi, *Tetrahedron Lett.*, **1998**, *39*, 7553-7556.
- 2. X. Fu, M. B. Hossain, F. J. Schmitz, and D. van der Helm, J. Org. Chem., 1997, 62, 3810-3819.
- 3. a) Y. Masaki, K. Hashimoto, K. Sakura, and K. Kaji, *Chem Pharm. Bull. Jpn.*, 1984, 32, 3952-3958.
  b) M. Bulliard, G. Balme, and J. Gore, *Tetrahedron Lett.*, 1989, 30, 5767-5770.
- 4. The <sup>13</sup>C chemical shifts of C<sub>3</sub>-methyls (CDCl<sub>2</sub>); 4a 16.7 and 4b 26.3 ppm, respectively.
- 5. It needed 90 h for the completion of the reaction of run 2.
- 6. The effect of other Lewis acids is being examined.

- 7. The <sup>13</sup>C chemical shifts of olefinic methyls (CDCl<sub>3</sub>); **6a** 16.7, 16.8, and 26.3, and **6b** 16.5, 16.8, and 16.9 ppm, respectively.
- 8. D. G. Saunders, Synthesis, 1988, 377-379. The formation conditions of 7c was not optimized.
- 9. Kobayashi et al succeeded in the intermolecular coupling reaction of allyl chlorides with various types of aromatics with  $Hf(OTf)_4$  and  $LiClO_4$ .<sup>10</sup> In our compound, however, the allyl chloride (**7b**, Cl instead of OH), prepared from **7b** in 90 % yield by the action of MsCl, Et<sub>3</sub>N, LiCl, 12-crown-4 in CH<sub>2</sub>Cl<sub>2</sub>, afforded no coupling product under these conditions. Fukuzawa et al<sup>11</sup> carried out the intermolecular coupling reactions with Sc(OTf)<sub>3</sub> using allyl alcohols and aromatics. In our case, however, application of Sc(OTf)<sub>3</sub> to **7b** gave no satisfactory fesults.

The authors are indebted to Professor S. Kobayashi for his kind gift of qualified Hf(OTf)<sub>4</sub>.

- 10. a) I. Hachiya, M. Moriwaki, and S. Kobayashi, *Tetrahedron Lett.*, 1995, *36*, 409-412.
  b) I. Hachiya, M. Moriwaki, and S. Kobayashi, *Bull. Chem. Soc. Jpn.*, 1995, *68*, 2053-2060.
- 11. T. Tsuchimoto, K. Tobita, T. Hiyama, and S. Fukuzawa, J. Org. Chem., 1997, 62, 6997-7005.
- 12. After deprotection of TBDMS group with TBAF in THF (90% yield), the resultant phenolic mixture was treated with PivCl, pyr., DMAP in  $CH_2Cl_2$  to give the pivalates. The HPLC analysis showed an 8 : 1 ratio of **8b** and **8a** (R = Piv).
- 13. The HPLC analysis indicated a 6 : 1 ratio of 8a and 8b (R = Piv).
- 14. As reported by Schmitz et al,<sup>2</sup> the quinones (1 and 2) were fairly unstable, particularly in oxygen atmosphere.
- 15. The preliminary MM2 calculation shows the molecular energies of 1, 2 and quinone derived from 8c are 15.34, 17.06, and 23.12 kcal/mol, respectively, reflecting the sterically crowded nature of 8c.
- 16. Some efforts to resolve the *dl*-form into the optically active compounds were unsuccessful using chiral columns. The synthetic compound (1) has the following physical data. <sup>1</sup>H-NMR(500MHz, C<sub>6</sub>D<sub>6</sub>); 6.37(t, *J*=1.5Hz, Δ-0.08; H-21), 6.23(d, *J*=1.0Hz, Δ0; H-18), 5.45(t, *J*=7.5Hz, Δ+0.05; H-2), 5.18(dm, *J*=9.5Hz, Δ+0.08; H-10), 4.71(br m, Δ-0.02; H-6), 3.38(dd, *J*=18.0, 1.5Hz, Δ-0.08; H-12), 3.34(dd, *J*=10.5, 7.5Hz, Δ+0.01; H-1), 2.44(dd, *J*=11.5, 9.5Hz, Δ-0.01; H-1), 2.42(dd, *J*=17.9, 1.2Hz, Δ-0.02; H-12), 1.97(m, Δ+0.03; H-4, H-5), 1.87(m, Δ-0.02; H-9a), 1.86(m, Δ-0.01; H-8a), 1.71(m, Δ-0.02; H-8b), 1.64(s, Δ+0.01; H-15), 1.54(s, Δ+0.16; H-14), 1.45(m, Δ0; H-9b), 1.38(s, Δ+0.05; H-13) <sup>13</sup>C-NMR(125MHz, C<sub>6</sub>D<sub>6</sub>); 187.2(s, Δ-0.9; C-20), 187.1(s, Δ-0.7; C-17), 147.3(s, Δ-1.0; C-16), 147.3(s, Δ-0.5; C-19), 140.1(s, Δ-1.6; C-3), 135.0(s, Δ-0.2; C-7), 132.5(d, Δ+0.1; C-21), 130.3(s, Δ-0.6; C-11), 129.7(d, Δ-0.5; C-18), 128.3(d, Δ+0.9; C-10), 123.2(d, Δ+0.6; C-6), 121.3(d, Δ-0.8; C-2), 39.3(t, Δ-0.2; C-8), 38.4(t, Δ-0.6; C-4), 31.8(t, Δ-0.3; C-12), 29.5(t, Δ+0.4; C-11), 28.3(t, Δ-0.9; C-9), 26.2(q, Δ-0.3; C-15), 24.7(t, Δ+0.6; C-5), 16.1(q, Δ+0.5; C-14), 14.4(q, Δ-0.2; C-13).

The  $\Delta$  values in the above chemical shifts are the deviation of the synthetic compound (1) from the reported values.<sup>2</sup> Some of the chemical shifts in both <sup>1</sup>H- and <sup>13</sup>C-NMR spectra are obviously inconsistent beyond experimental deviation. The chemical shifts of <sup>1</sup>H and <sup>13</sup>C NMR spectra of the synthetic compound (1) were almost same when measured under two different concentrations in 0.5 ml C<sub>6</sub>D<sub>6</sub> solutions, the sample being 7 mg and 2 mg, respectively. We have independently carried out the NMR analysis of our synthetic compound (1) using H-H COSY, C-H COSY and NOESY spectra, leading the synthetic compound to the several structures including the formula (1).

At present, the reason of the inconsistency remains obscure.