

RSC Advances



This article can be cited before page numbers have been issued, to do this please use: J. Doroszuk, M. Musiejuk, S. Demkowicz, J. Rachon and D. Witt, *RSC Adv.*, 2016, DOI: 10.1039/C6RA19440K.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

COMMUNICATION

Convenient and efficient synthesis of functionalized unsymmetrical alkynyl sulfides

J. Doroszk, ^a M. Musiejuk, ^a S. Demkowicz, ^a J. Rachon, ^a and D. Witt ^{a*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

We developed a simple and efficient method for the synthesis of functionalized unsymmetrical alkynyl sulfides under mild conditions in good yields. The designed method is based on the reaction of 5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinan-2-disulfanyl derivatives with lithium acetylides. The developed method allows the preparation of unsymmetrical alkynyl sulfides bearing additional hydroxyl, carboxyl, or amino functionalities.

Introduction

The development of practical and versatile methods for introduction of functional groups into molecules to modify their properties is one of the most important task of organic chemistry. The easy to perform reactions that do not require highly specialized synthetic skills have a particularly broad impact to other fields such as medicine, biology, or materials science. From this point of view, among all functional groups in organic chemistry, alkynes occupy a privileged position.¹ Although alkynes contain one of the simplest functional group, two carbon atoms connected with triple bond, the reactivity of that bond makes alkynes exceptionally useful in organic synthesis. They were applied in the stereoselective construction of the carbon backbone of complex natural products,² bulk chemical synthesis based on acetylene gas³ and in the variety of complexity-enhancing metal-catalyzed cyclization reactions to provide carbo- and heterocycles.⁴ The electronic properties of alkynes have found also wide applications in preparation of organic materials and dyes.⁵ Moreover, their [3+2] cycloaddition with azides is recognized as one of the best biorthogonal conjugation method to modify

biomolecules and polymers.⁶ In this context, the development of new synthetic methods to access alkynes and their derivatives efficiently under user-friendly conditions is highly desirable.

Driven by the exceptional reactivity of sulfur and its importance in biology, medicine, and materials science,⁷ recent research efforts targeted a series of thiol-based transformations including thiol alkylations and the thiol addition to alkenes and alkynes respectively.⁸ Unlike the well-established S-Csp³ bond forming reactions, existing methods to construct S-Csp bonds are rare in number and often lack generality or require harsh conditions.

Alkynyl sulfides can be obtained by methods utilize transition-metal catalysts, such as the copper-catalyzed carbon sulfur coupling between terminal alkynes and disulfides,⁹ or as in the elegant study by Yamaguchi, the use of catalytic rhodium to achieve C-S bond formation by C-H and S-S bond metathesis (Scheme 1, A).¹⁰ Alternatively, a range of processes utilize alkenyl¹¹ or alkynyl¹² halides bearing leaving groups that undergo elimination under strongly basic conditions to furnish the desired alkynyl sulfides (Scheme 1, B). Consequently, the limited functional group tolerance exhibited by these methods is not surprising as they require harsh conditions, proceed via highly reactive intermediates, or involve the use of sensitive catalytic systems. Recently, Waser has developed a thiol-alkynylation procedure utilizing the hypervalent iodine alkyne transfer reagent TIPS-ethynylbenziodoxolone (Scheme 1, C).¹³ Although the method is highly chemoselective as a vast array of functional groups are tolerated, the problems associated with preparation and stability hypervalent iodine alkyne transfer reagent are the major disadvantages of that transformation. Currently, the most common methods to form alkynyl sulfides require a prefunctionalization of the thiol (Scheme 1, D). These methods are generally based on nucleophilic substitutions between highly reactive lithium acetylide intermediates with pre-activated thiols or disulfide species.¹⁴ The major drawback of that approached emerges from availability and long term stability of pre-activated thiols.


^a Department of Organic Chemistry, Faculty of Chemistry, Gdansk University of Technology, Narutowicza 11/12, 80-233 Gdansk, Poland. Fax: +48 58 3472694. E-mail: chemwitt@pg.gda.pl

† Footnotes relating to the title and/or authors should appear here.
Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x


COMMUNICATION

Journal Name


A. S-Csp bonds from transitions metals promoted coupling of disulfides and alkynes



B. S-Csp bonds directly from thiols




C. S-Csp bonds from thiols and hypervalent iodine alkynes



D. S-Csp bonds from pre-activated thiols



E. This work




Scheme 1. Previously reported methods for the synthesis of alkynyl sulfides (A-D) and our new approach (E)

We have previously demonstrated the preparation of functionalized unsymmetrical molecules, such as dialkyldisulfanes,¹⁵ alkyl-aryl disulfanes,¹⁶ 'bioresistant' disulfanes,¹⁷ the unsymmetrical disulfanes of L-cysteine and L-cystine,¹⁸ and diaryldisulfanes,¹⁹ based on the readily available electrophilic 5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinane-2-disulfanyl derivatives **2**.²⁰ These disulfanyl derivatives of phosphorodithioic acid were also convenient for the preparation of α -sulfenylated carbonyl compounds,²¹ functionalized phosphorothioates,²² as well as symmetrical²³⁻²⁴ and unsymmetrical²⁵⁻²⁶ trisulfanes. All these transformations are based on the electrophilic properties of disulfanyl derivatives of phosphorodithioic acid **2**, so we expected that also reaction with nucleophilic acetylides should provide corresponding alkynyl sulfides (Scheme 1, E). Furthermore, the development of a robust, efficient, and orthogonal alkynyl sulfides synthesis is particularly attractive to the field of

material and polymer science as an alternative tool to the current array of thiol-functionalization reactions. In this context, we set out to investigate the feasibility of more convenient and experimentally practical method to gain access to alkynyl sulfides.


Results and discussion

The terminal alkynes can be easily converted into the 1-Cu(I) acetylides.²⁷ The presence of additional groups (as carbonyl, carboxyl or hydroxyl) did not disturb the formation of corresponding salts. Moreover, these compounds were stored in a bottle and no notable decomposition was observed within one year. We began our study with copper (I) phenylacetylide **1a** as starting substrate while screening different disulfanyl derivatives reagents capable to produce corresponding phenylethynyl sulfides **3** (Scheme 2).




Scheme 2. The reaction of copper (I) phenylacetylide **1a** with disulfanyl derivatives **2**.

Unfortunately, in the most cases the reaction did not occur or provided complex mixture without formation of compound **3**. The only successful transformation was the reaction of copper (I) phenylacetylide with phosphorodithioic acid disulfane **2a** when the expected product **3a** was isolated in 67% yield. Additionally, we have accomplished the synthesis of **3a** by independent method based on the phenylethynylphenyliodonium tosylate²⁸ to confirm the structure of the final product (Scheme 3).

Scheme 3. The synthesis of 1-[(5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinan-2-yl)sulfanyl]-2-phenylacetylene **3a**

We examined the reactivity of functionalized copper acetylides **1** with disulfane **2a**. The results are summarized in the Table 1.


Table 1. Reaction of 1-Copper (I) acetylides **1** with disulfane **2a**^a.

Entry	R	Yield ^b (%)	
		3	4
1	Ph- 1a	67	30
2	PhCH ₂ OCH ₂ - 1b	40	55
3	BocNH-CH ₂ - 1c	30	40

^a Conditions: copper alkyne **1** (1.0 mmol), disulfane **2a** (1.0 mmol), 10.0 mL of chloroform, reflux 15 min. under N₂. ^b Isolated yields.

As the data in Table 1 demonstrate, the yield of product **3** is moderate (30-67%) and the reaction is hampered by the formation of dimer **4** (30-55%). Although acetylides **1** are stable and readily available their application in the alkynyl sulfides synthesis, under developed conditions, seems to be limited by an oxidative coupling that produces bis-acetylenes **4**.

The reaction of electrophilic disulfanyl derivatives **2** and lithium acetylides **5** was expected to produce the corresponding alkynyl sulfides **3** with higher yield due to the limited formation of bis-acetylenes **4**. The results are summarized in Table 2.

Table 2. Reaction of Lithium Acetylides **5** with disulfanes **2**

Entry	R ¹	R ²	Yield ^b of 3
1	Ph- 5a		(78) 3a
		2a	
2	PhCH ₂ OCH ₂ - 5b		(75) 3b
		2a	
3	BocNHCH ₂ - 5c		(60) 3c
		2a	
4	Ph- 5a	CH ₃ (CH ₂) ₁₁ - 2b	(99) 3d
5	PhCH ₂ OCH ₂ - 5b	CH ₃ (CH ₂) ₁₁ - 2b	(83) 3e
6	BocNHCH ₂ - 5c	CH ₃ (CH ₂) ₁₁ - 2b	(57) 3f
7	Ph- 5a	HO(CH ₂) ₁₁ - 2i	(63) 3g
8	PhCH ₂ OCH ₂ - 5b	HO(CH ₂) ₁₁ - 2i	(55) 3h
9	BocNHCH ₂ - 5c	THPO(CH ₂) ₁₁ - 2l	(50) 3i
10	Ph- 5a	HO ₂ C(CH ₂) ₁₁ - 2j	(72) 3j
11	PhCH ₂ OCH ₂ - 5b	HO ₂ C(CH ₂) ₁₁ - 2j	(73) 3k
12	BocNHCH ₂ - 5c	MeO ₂ C(CH ₂) ₁₁ - 2m	(51) 3l
13	Ph- 5a	4-CH ₃ -C ₆ H ₄ - 2k	(98) 3m
14	PhCH ₂ OCH ₂ - 5b	4-CH ₃ -C ₆ H ₄ - 2k	(89) 3n
15	BocNHCH ₂ - 5c	4-CH ₃ -C ₆ H ₄ - 2k	(78) 3o
16	4-MeO-2-MeC ₆ H ₃ - 5d	CH ₃ (CH ₂) ₁₁ - 2b	(91) 3p
17	(CH ₃) ₂ CHCH ₂ CH ₂ - 5e	CH ₃ (CH ₂) ₁₁ - 2b	(100) 3r
18	EtO ₂ C- 5f	CH ₃ (CH ₂) ₁₁ - 2b	(54) 3s
19	(CH ₃) ₂ CHCH ₂ CH ₂ - 5e	2-naphthyl- 2n	(86) 3t
20	(CH ₃) ₂ CHCH ₂ CH ₂ - 5e		(100) 3u
		2o	
21	4-NO ₂ -C ₆ H ₄ - 5g	CH ₃ (CH ₂) ₁₁ - 2b	(0) 3w

^a Conditions: Lithium acetylide **5** (1.0 mmol), TMEDA (1.0 mmol), disulfane **2** (1.1 mmol), 12 mL of THF, rt 30 min. under N₂. ^b Isolated yields in parenthesis.

The lithium acetylides **5** were generated from the corresponding terminal alkynes and BuLi in the presence of *N,N,N,N'*-tetramethylethylenediamine (TMEDA) in THF to avoid the potential aggregation of lithium salts. The lithium phenylacetylide **5a** was obtained at -78 °C, -35 °C and 0 °C. The subsequent reaction with **2a** provided product **3a** in 76, 81 and 78% yields respectively. We did not observed substantial influence of temperature on the generation of acetylide **5a** and the yield of final product **3a**. That is why, acetylides **5b-g** were generated at 0 °C. In the case of disulfanes **2i** and **2j** with acidic protons (OH and CO₂H respectively, Table 2 entries 7-8 and 10-11) the two fold excess of **5a** and **5b** was used. It was also possible to treat **2i** and **2j** with NaH before addition to

solution of **5a** or **5b** in THF. Both methods provided appropriate alkynyl sulfides **3g**, **3h** and **3j**, **3k** with comparable yield respectively. However, the generation of **5c** from *N*-Boc propargylamine required the using of two equivalents of BuLi (Table 2 entries 3, 6, 9, 12, 15). Moreover, the corresponding alkynyl sulfides **3i** and **3l** were obtained when hydroxyl and carboxyl groups were protected in disulfanes **2l** and **2m** (THP or methyl ester respectively, Table 2 entries 9 and 12). The corresponding alkynyl sulfides **3** can be obtained from alkyl, aryl acetylides **5** and disulfanyl derivatives **2** produced from alkyl, aryl and heteroarylthiols. When electron-withdrawing group was present in the acetylide **5e**, the corresponding product **3s** was obtained in moderate yield (Table 2 entry 18). However, reaction of lithium 4-nitrophenylacetylide **5f** with disulfane **2b** did not produce expected product **3w** (Table 2 entry 21). Both starting materials were consumed and formation of complex reaction mixture was observed. The same result was obtained when the reaction was performed at -78 °C. The influence of other electron-withdrawing groups on the course of the reaction is under investigation. On other hand, the presence of electron-donating groups on the phenylacetylide **5d** did not disturb the formation of alkynyl sulfide **3p** and the product was isolated in 91% yield (Table 2 entry 16). The preparation of alkynyl sulfides **3** from disulfanyl derivatives **2** and lithium acetylides **5** is very convenient because both starting materials are stable and readily available. The transformation tolerates the presence of additional hydroxyl, carboxyl and amino groups. However, the presence of acidic protons in the starting materials may require the using of larger quantity of base to avoid the protonation of acetylides anions.

Conclusions

In summary, we developed an efficient and convenient method for the preparation of unsymmetrical alkynyl sulfides **3** directly from the readily available disulfanyl derivatives of phosphorodithioic acid **2** and functionalized alkynes in the presence of TMEDA and BuLi. A variety of functional groups is tolerated, including the hydroxyl, carboxyl, and protected amino, hydroxyl and carboxyl group. Reactions of lithium acetylides **5** with a variety of disulfanes **2** in the presence of TMEDA in THF at rt were generally complete within 30 minutes and gave unsymmetrical alkynyl sulfides **3** exclusively in good or very good yield after isolation. The simplicity and good yields render this method one of the most attractive approaches to the preparation of functionalized unsymmetrical alkynyl sulfides.

Experimental

A typical procedure for the preparation of alkynyl sulfide **3d** and representative analytical data

BuLi (2.5M, 0.44 mL, 1.1 mmol) was added to a solution of phenylacetylene (0.11 mL, 1.0 mmol) and *N,N,N',N',*-tertmetylethylenediamine (TMEDA) (0.15 mL, 1.0 mmol) in

anhydrous THF (10 mL) at 0 °C under N₂. After 5 min, a solution of 1-[(5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinan-2-yl)disulfanyl]dodecane **2b** (0.438 g, 1.1 mmol) in anhydrous THF (2 mL) was added. The mixture was stirred at r.t. for 30 min. The reaction was quenched by addition of MeOH (1 mL) and evaporated under vacuum. The residue was purified by column chromatography (SiO₂, petroleum ether, R_f = 0.6) to give **3d** as a colorless oil; yield: 0.300 g (99%).

IR (ATR): 2900 (s), 2850 (s) (C-H), 2170 (w) (C≡C), 1600 (w), 1450 (w), 1370 (w), 750 (m), 700 (m) cm⁻¹

¹H NMR (500 MHz, CDCl₃): δ = 0.89 (t, *J* = 7.0 Hz, 3H, CH₃), 1.20-1.40 (m, 16H, CH₂), 1.46 (qu, *J* = 7.0 Hz, 2H, CH₂), 1.8 (qu, *J* = 7.4 Hz, 2H, CH₂), 2.81 (t, *J* = 7.3 Hz, 2H, SCH₂), 7.26-7.35 (m, 3H, Ar), 7.40-7.45 (m, 2H, Ar).

¹³C NMR (125 MHz, CDCl₃): δ = 131.4, 128.2, 127.9, 123.6, 92.8, 79.7, 35.8, 31.9, 29.6, 29.6, 29.6, 29.5, 29.3, 29.3, 29.1, 28.2, 22.7, 14.1; signals: 19 expected, 18 observed.

HRMS (ESI): *m/z* [M + H]⁺ calcd for C₂₀H₃₁S: 303.2146; found: 303.2152.

Acknowledgments

We gratefully acknowledge the National Science Centre (NCN) for financial support (grant no. 2015/19/B/ST5/03359)


Notes and references

- 1 *Acetylene Chemistry: Chemistry, Biology and Material Science*; F. Diederich, P. J. Stang, R. R. Tykwinski, Eds.; Wiley-VCH: Weinheim, 2005.
- 2 (a) D. E. Frantz, R. Fassler, E. M. Carreira, *J. Am. Chem. Soc.*, 2000, **122**, 1806. (b) B. M. Trost, A. H. Weiss, *Adv. Synth. Catal.*, 2009, **351**, 963.
- 3 (a) H. Schobert, *Chem. Rev.*, 2014, **114**, 1743. (b) I. T. Trotus, T.; Zimmermann, F. Schuth, *Chem. Rev.*, 2014, **114**, 1761.
- 4 (a) G. Zeni, R. C. Larock, *Chem. Rev.*, 2004, **104**, 2285. (b) V. Michelet, P. Y. Toullec, J. P. Genet, *Angew. Chem., Int. Ed.* 2008, **47**, 4268. (c) E. Jimenez-Nunez, A. M. Echavarren, *Chem. Rev.*, 2008, **108**, 3326. (d) B. Godoi, R. F. Schumacher, G. Zeni, *Chem. Rev.*, 2011, **111**, 2937.
- 5 (a) U. H. F. Bunz, *Chem. Rev.*, 2000, **100**, 1605. (b) J. Z. Liu, J. W. Y. Lam, B. Z. Tang, *Chem. Rev.*, 2009, **109**, 5799. (c) A. Yella, H. W. Lee, H. N. Tsao, C. Y. Yi, A. K. Chandiran, M. K. Nazeeruddin, E. W. G. Diau, C. Y. Yeh, S. M. Zakeeruddin, M. Gratzel, *Science* 2011, **334**, 629.
- 6 (a) H. C. Kolb, M. G. Finn, K. B. Sharpless, *Angew. Chem., Int. Ed.* 2001, **40**, 2004. (b) V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, *Angew. Chem., Int. Ed.* 2002, **41**, 2596. (c) C. W. Tornoe, C. Christensen, M. Meldal, *J. Org. Chem.*, 2002, **67**, 3057. (d) M. Meldal, C. W. Tornoe, *Chem. Rev.*, 2008, **108**, 2952. (e) N. J. Agard, J. A. Prescher, C. R. Bertozzi, *J. Am. Chem. Soc.*, 2004, **126**, 15046. (f) E. M. Sletten, C. R. Bertozzi, *Angew. Chem., Int. Ed.* 2009, **48**, 6974. (g) P. Thirumurugan, D. Matosiuk, K. Jozwiak, *Chem. Rev.*, 2013, **113**, 4905.
- 7 (a) B. P. Zambrowicz, A. T. Sands, *Nat. Rev. Drug Discovery* 2003, **2**, 38. (b) L. A. Paquette, *Sulfur-Containing Reagents*; Wiley: Chichester, 2009. (c) R. Masella, *Glutathione and Sulfur Amino Acids in Human Health and Disease*; Wiley: Hoboken, 2009. (d) C. E. Hoyle, A. B. Lowe, C. N. Bowman, *Chem. Soc. Rev.*, 2010, **39**, 1355.

- 8 (a) R. L. Lundblad, *Chemical reagents for protein modification*, 3rd ed.; CRC Press: Boca Raton, 2005. (b) G. T. Hermanson, *Bioconjugate techniques*, 2nd ed.; Academic Press: San Diego, 2008. (c) C. E. Hoyle, C. N. Bowman, *Angew. Chem., Int. Ed.* 2010, **49**, 1540. (d) R. Hoogenboom, *Angew. Chem., Int. Ed.* 2010, **49**, 3415. (e) A. B. Lowe, C. E. Hoyle, C. N. Bowman, *J. Mater. Chem.*, 2010, **20**, 4745. (f) N. Stephanopoulos, M. B. Francis, *Nat. Chem. Biol.*, 2011, **7**, 876. (g) M. H. Stenzel, *ACS Macro Lett.*, 2012, **2**, 14.
- 9 (a) A. L. Braga, A. Reckziegel, P. H. Menezes, H. A. Stefani, *Tetrahedron Lett.*, 1993, **34**, 393. (b) L. W. Bieber, M. F. da Silva, P. H. Menezes, *Tetrahedron Lett.*, 2004, **45**, 2735.
- 10 M. Arisawa, K. Fujimoto, S. Morinaka, M. Yamaguchi, *J. Am. Chem. Soc.*, 2005, **127**, 12226.
- 11 (a) I. Marchueta, E. Montenegro, D. Panov, M. Poch, X. Verdaguer, A. Moyano, M. A. Pericas, A. Riera, *J. Org. Chem.*, 2001, **66**, 6400. (b) Z. Ni, S. Wang, H. Mao, Y. Pan, *Tetrahedron Lett.*, 2012, **53**, 3907.
- 12 G. R. Ziegler, C. A. Welch, C. E. Orzech, S. Kikkawa, S. I. Miller, *J. Am. Chem. Soc.*, 1963, **85**, 1648.
- 13 (a) R. Frei, J. Waser, *J. Am. Chem. Soc.*, 2013, **135**, 9620. (b) R. Frei, M. D. Wodrich, D. P. Hari, P.-A. Borin, C. Chauvier, J. Waser, *J. Am. Chem. Soc.*, 2014, **136**, 16563.
- 14 (a) B. Godoi, A. Speranca, D. F. Back, R. Brandao, C. W. Nogueira, G. Zeni, *J. Org. Chem.*, 2009, **74**, 3469. (b) J. Z. Chandanshive, B. F. Bonini, D. Gentili, M. Fochi, L. Bernardi, M. C. Franchini, *Eur. J. Org. Chem.*, 2010, 6440.
- 15 S. Lach, S. Demkowicz, D. Witt, *Tetrahedron Lett.*, 2013, **54**, 7021.
- 16 S. Antoniow, D. Witt, *Synthesis* **2007**, 363.
- 17 J. Kowalczyk, P. Barski, D. Witt, B. A. Grzybowski, *Langmuir* 2007, **23**, 2318.
- 18 M. Szymelfejnik, S. Demkowicz, J. Rachon, D. Witt, *Synthesis* **2007**, 3528.
- 19 S. Demkowicz, J. Rachon, D. Witt, *Synthesis* **2008**, 2033.
- 20 M. Musiejuk, T. Klucznik, J. Rachon, D. Witt, *RSC Adv.*, 2015, **5**, 31347.
- 21 E. Okragla, S. Demkowicz, J. Rachon, D. Witt, *Synthesis* **2009**, 1720.
- 22 S. Lach, D. Witt, *Synthesis* **2011**, 3975.
- 23 A. Kertmen, S. Lach, J. Rachon, D. Witt, *Synthesis* **2009**, 1459.
- 24 S. Lach, D. Witt, *Heteroatom Chem.*, 2014, **25**, 10.
- 25 S. Lach, M. Sliwka-Kaszynska, D. Witt, *Synlett* **2010**, 2857.
- 26 S. Lach, D. Witt, *Synlett* 2013, **24**, 1927.
- 27 W. Chen, B. Wang, N. Liu, D. Huang, X. Wang, Y. Hu, *Org. Lett.*, 2014, **16**, 6140.
- 28 Z.-D. Liu, Z.-C. Chen, *J. Org. Chem.*, 1993, **58**, 1924.

RSC Author Templates - ChemDraw (CHM) - Graphical Abstracts

All text and images must be placed within the frame.



81x71mm (300 x 300 DPI)