

An Efficient Improve for the Kröhnke Reaction: One-pot Synthesis of 2,4,6-Triarylpyridines Using Raw Materials under Microwave Irradiation

Shujiang Tu,* Tuanjie Li, Feng Shi, Fang Fang, Songlei Zhu, Xianyong Wei,[†] and Zhimin Zong[†]
*Department of Chemistry, Xuzhou Normal University, Key Laboratory of Biotechnology for Medicinal Plant,
 Xuzhou, Jiangsu, 221116, P. R. China*

[†]*School of Chemical Engineering, China University of Mining and Technology, Xuzhou, Jiangsu, 221008, P. R. China*

(Received February 21, 2005; CL-050227)

A series of 2,4,6-triarylpyridines have been prepared by the one-pot reaction of aldehydes with aromatic ketones in the presence of ammonium acetate under microwave irradiation without catalyst. This method has the advantage of easier workup, shorter reaction time, higher yield, and environment-friendly.

Kröhnke type pyridines¹ and substituted phenylpyridines^{2,3} are useful intermediates in the synthesis of drugs, agrochemicals, herbicides, insecticides, desiccants, surfactant agents, and antiinflammatory agents.^{4,5} They are widely used as ligands in coordination complex preparation. Triarylpyridines are prominent building blockers in supramolecular chemistry with their π -stacking ability, directional H-bonding and coordination. Hence, their synthesis has received much attention.


Traditionally, triarylpyridines are synthesized using the Kröhnke approach¹ by the reaction of pyridium salts with an unsaturated ketone, which started from the preparation of pyridium salts and involved the release of pyridium halide in the last step. And many articles adopting the method to synthesize 2,4,6-triarylpyridines have been described.^{6–12} However, these methods coupled with the fact that the pyridinium halide and α,β -unsaturated ketone often have to be synthesized first. So it was an expensive, time consuming protocol and did harm to the environment. Since then, some new methods for the synthesis of 2,4,6-triarylpyridines have also been reported. One employed the reaction of β -enaminophosphonates with chalcones in tetrahydrofuran catalyzed by butyllithium.¹³ Gareth reported the synthesis of triarylpyridines in the presence of sodium hydroxide using a multistep solid-phase reaction.¹⁴

Green chemistry is an environmental, health, and safe strategy that emphasizes pollution prevention and application of chemical process to reduce or eliminate the use and generation of hazardous substance.

We investigated that microwave irradiation could assist this reaction, and streamlined the usual multistep procedure to a single step, cut it use of poisonous catalyst and eliminated the generation of poisonous pyridium halide. In our previous paper, we have reported the synthesis of heterocyclic compounds under microwave irradiation.¹⁵ In this paper, we would like to report the one-pot synthesis of 2,4,6-triarylpyridines using raw materials under microwave irradiation without catalyst.

Under microwave irradiation, the reaction of an aldehyde with an aromatic ketone in the presence of ammonium acetate completed within 6 min and yielded **1a–1x** in 80–96% yields (Scheme 1).¹⁶ While in traditional heating mode (110 °C), the reaction time is 2 h and the yields are 70–78%, which indicates this reaction can be promoted by microwave irradiation.

The results (Table 1) show that a wide range of aromatic al-




Scheme 1.

dehydes and aromatic or heteroaromatic ketones can take part in this reaction. This method is simple and easy work up.

All these products were characterized by IR and ¹H NMR analysis, and their melting points were identical to those of the known compounds reported in the literature. Meanwhile, the structure of the compound **1s** was established on the basis of spectroscopic data and conformed by X-ray diffraction study

Table 1. The synthesis of triarylpyridine **1**


Entry	Ar	R	Yield/%	Mp (°C).Lit.
1a	4-ClC ₆ H ₄	3-O ₂ NC ₆ H ₄	92	285.6–286.6
1b	4-ClC ₆ H ₄	4-CH ₃ C ₆ H ₄	90	200.6–202
1c	4-ClC ₆ H ₄	2-ClC ₆ H ₄	91	155.8–156
1d	4-ClC ₆ H ₄	4-FC ₆ H ₄	92	209.4–210.1
1e	4-ClC ₆ H ₄	4-CH ₃ OC ₆ H ₄	91	113.8–115.0
1f	4-ClC ₆ H ₄	2,4-Cl ₂ C ₆ H ₃	95	176.4–177.0
1g	4-O ₂ NC ₆ H ₄	4-CH ₃ OC ₆ H ₄	90	143.1–144.7
1h	3-O ₂ NC ₆ H ₄	3-O ₂ NC ₆ H ₄	91	>300
1i	4-CH ₃ OC ₆ H ₄	2,4-Cl ₂ C ₆ H ₃	89	161.0–161.8
1j	4-CH ₃ OC ₆ H ₄	4-CH ₃ C ₆ H ₄	92	156.8–157.9 (156–157.5) ¹⁸
1k	4-CH ₃ OC ₆ H ₄	4-CH ₃ OC ₆ H ₄	92	136–137 (135) ¹⁹
1l	2-ClC ₆ H ₄	2-ClC ₆ H ₄	89	146.5–147.3
1m	2-ClC ₆ H ₄	2,4-Cl ₂ C ₆ H ₃	89	201.2–201.6
1n	4-BrC ₆ H ₄	2,4-Cl ₂ C ₆ H ₃	92	177.9–178.2
1o	4-BrC ₆ H ₄	4-CH ₃ OC ₆ H ₄	92	163.9–165
1p	C ₆ H ₅	2,4-Cl ₂ C ₆ H ₃	88	160.1–161
1q	2,4-Cl ₂ C ₆ H ₃	2,4-Cl ₂ C ₆ H ₃	91	203.9–204.8
1r	3,4-Cl ₂ C ₆ H ₃	2,4-Cl ₂ C ₆ H ₃	90	152.9–154.8
1s	3-indole	4-CH ₃ OC ₆ H ₄	89	232.0–233.0
1t	2,4-Cl ₂ C ₆ H ₃	2-pyridyl	84	158.4–159.3
1u	4-CH ₃ OC ₆ H ₄	2-pyridyl	81	167–168 (171–172) ²⁰
1v	4-BrC ₆ H ₄	2-pyridyl	89	154–156 (158–160) ²⁰
1w	4-CH ₃ C ₆ H ₄	2-pyridyl	80	166–167 (166–167) ²⁰
1x	C ₆ H ₅	2-pyridyl	80	210.8–211.6 (206–207) ²⁰

Figure 1. X-ray structures of **1s**.(Figure 1).¹⁷

However, when using aldehyde and two different ketones as the starting material, we obtained mixture including two different chalcone, which are primary products, and a little amount of unsymmetrical triarylpyridine.

This method is applied to aromatic and heteroaromatic ketones. Furthermore, this versatile method is also suitable for the synthesis of 1-indenone and α -tetralone. 11-aryl-10*H*,12*H*-diindeno[1,2-*b*:2',1'-*e*]pyridine **2** and 7-aryl-5,6,8,9-tetrahydro-dibenzo[*c,h*]acridine **3** were obtained with good yields (Scheme 2). The results are listed in Table 2.

Here, we disclose a facile method to synthesize a wide range of symmetrical triarylpyridines. This methodology is superior to the reported methods from the view of green chemistry. The significance of our approach relates to the elimination of toxic starting materials, as well as its simplicity and avoiding the release of



Scheme 2.

Table 2. The synthesis of **2**, **3**

Entry	Ar	Yield/%	Mp (°C).Lit.
2a	C ₆ H ₅	93	>300 (298) ¹
2b	4-OH-3-CH ₃ OC ₆ H ₃	92	>300
2c	4-ClC ₆ H ₄	95	>300
2d	3,4-OCH ₂ OC ₆ H ₃	93	>300
2e	4-BrC ₆ H ₄	96	>300
2f	4-CH ₃ OC ₆ H ₄	93	251.8–252.2
2g	3,4-Cl ₂ C ₆ H ₃	94	>300
2h	3-indole	90	>300
3a	4-OH-3-CH ₃ OC ₆ H ₃	91	292.0–292.8
3b	3,4-Cl ₂ C ₆ H ₃	90	147.6–148.9
3c	C ₆ H ₅	90	167.5–168.9
3d	4-CH ₃ OC ₆ H ₄	92	183.6–184.4
3e	3,4-OCH ₂ OC ₆ H ₃	93	193.0–195.6
3f	4-ClC ₆ H ₄	95	260.0–262.0
3g	4-BrC ₆ H ₄	95	279.6–280.0

any hazardous products. In summary, this procedure is simple, environment-friendly.

We thank the Natural Science Foundation of the China (No. 20372057), the Key Laboratory of Biotechnology for Medicinal Plants of Jiangsu Province (No. 01AXL 14), the Open-end Fund of the Key Experiments of Organic Synthesis, Jiangsu Province (S8109111) for financial support.

References and Notes

- 1 F. Kröhnke, *Synthesis*, **1976**, 1.
- 2 F. Neve, A. Grispi, and S. Campagna, *Inorg. Chem.*, **36**, 6150 (1997).
- 3 B. Olenyuk, J. A. Whiteford, A. Frechtenkötter, and P. J. Stang, *Nature*, **398**, 796 (1999).
- 4 S. Shimizu, N. Abe, N. Goto, T. Niwa, and A. Iguchi, Jpn. Patent, JP 01261367A (1998).
- 5 S. Peter, H. Gerhard, H. Elisabeth, K. Ralf, K. Hartmann, H. Albercht, G. Norbert, W. Helmut, W. Karl-Otto, and M. Uif, U. S. Patent, US 5 733850 (1998).
- 6 R. S. Tewari and N. K. Mishra, *Croat. Chem. Acta*, **54**, 241 (1981).
- 7 R. S. Tewari and A. K. Dubey, *J. Chem. Eng. Data*, **25**, 91 (1980).
- 8 N. K. Mishra, K. S. Mishra, and H. C. Srivastara, *Indian J. Agric. Chem.*, **21**, 91 (1988).
- 9 R. K. Singhal and N. K. Mishra, *Indian J. Chem., Sect. B*, **24B**, 1079 (1985).
- 10 T. Kobayashi and M. Nitta, *Chem. Lett.*, **1986**, 1549.
- 11 S. Sukhwai and B. L. Verma, *Indian J. Heterocycl. Chem.*, **31**, 287 (1994).
- 12 A. S. Kiselyor, *Tetrahedron Lett.*, **36**, 9297 (1995).
- 13 P. Francisco, M. Ana, R. Ochoa, and O. Julen, *Tetrahedron Lett.*, **37**, 4577 (1996).
- 14 W. V. C. Gareth and L. R. Collin, *Chem. Commun.*, **2000**, 2199.
- 15 S. Tu, C. Miao, F. Fang, F. Youjian, T. Li, Q. Zhuang, X. Zhang, S. Zhu, and D. Shi, *Bioorg. Med. Chem. Lett.*, **14**, 1533 (2004).
- 16 A Typical procedure for triarylpyridine: A dry flask (25 mL) was charged with a solution of the appropriate aldehyde (2 mmol), aromatic ketone (4 mmol), NH₄OAc (2 mmol) in a microwave oven. The flask was then connected with refluxing equipment. After irradiation for 5–9 min (irradiation sequences were interrupted with a cooling period in between), the reaction mixture was cooled and the crude solid precipitated from the solution was filtered and washed with 95% ethanol to afford the products. Analytical data of compound **1j**: IR (neat): 3038, 1609, 1598, 1580, 1543, 820 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 2.39 (6H, s, 2 CH₃), 3.85 (3H, s, OCH₃), 7.11 (2H, d, *J* = 8.0 Hz, ArH), 7.35 (2H, d, *J* = 8.0 Hz, ArH), 8.01 (2H, d, *J* = 8.0 Hz, ArH), 8.08 (2H, s, Pyridine-CH), 8.21 (2H, d, *J* = 8.0 Hz, ArH). compound **1s**: IR (neat): 3135, 3105, 3061, 3008, 2829, 1599, 1512, 1458, 1421, 1342, 1299, 1244, 1171, 1103, 1033, 880, 801, 753 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.86 (6H, s, 2OCH₃), 7.10–7.12 (4H, m, ArH), 7.22–7.24 (2H, m, ArH), 7.51–7.58 (2H, m, ArH), 8.08 (2H, s, Pyridine-CH), 8.08–8.11 (1H, m, Indole-CH), 8.24–8.27 (4H, m, Indole-CH), 11.75 (1H, s, Indole-NH). Compound **2f**: IR (neat): 3041, 1612, 1560, 1517, 858, 831, 749 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.88 (3H, s, OCH₃), 3.96 (4H, s, 2CH₂), 7.14 (2H, d, *J* = 8.0 Hz, ArH), 7.42–7.52 (4H, m, ArH), 7.63 (2H, d, *J* = 7.2 Hz, ArH), 7.76 (2H, d, *J* = 8.0 Hz, ArH), 8.11 (2H, d, *J* = 7.2 Hz, ArH). Compound **3a**: IR (neat): 3446, 3028, 1594, 1542, 1511, 766, 750, 736 cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ 2.60–2.87 (8H, m, 4CH₂), 3.81 (3H, s, OCH₃), 6.68 (1H, d, *J* = 8.4 Hz, ArH), 6.84 (s, 1H, ArH), 6.92 (1H, d, *J* = 8.4 Hz, ArH), 7.27–7.42 (6H, m, ArH), 8.42 (2H, d, ArH), 9.17 (1H, s, OH).
- 17 Crystal data for **1s**: Empirical formula C₂₇H₂₂N₂O₂, *M*_r = 406.47, *T* = 193(2) K, Triclinic, space group *P*1, *a* = 8.7438(13) Å, *b* = 16.341(2) Å, *c* = 16.460(2) Å, α = 68.442(11)°, β = 76.059(12)°, γ = 75.827(12)°, *V* = 2090.7(5) Å³, *Z* = 4, *D*_{calcd} = 1.291 Mg/cm³, λ (Mo K α) = 0.71073 Å, μ = 0.082 mm⁻¹, *F*(000) = 856, 3.05° < θ < 27.48°, *R* = 0.0741, *wR* = 0.1445, *S* = 1.160, Largest diff. Peak and hole: 0.248 and -0.207 e/Å³.
- 18 R. Lombard and J. P. Stephan, *Bull. Soc. Chim. Fr.*, **1958**, 1458.
- 19 G. N. Dorofenko, E. P. Olekhovich, and L. I. Laukhina, *Zh. Org. Khim.*, **7**, 1296 (1971).
- 20 T. Mutai, J. D. Cheon, S. Arita, and K. Araki, *J. Chem. Soc., Perkin Trans. 2*, **2001**, 1045.