

Influence of the N-Substituents on the Nucleophilicity and Lewis **Basicity of N-Heterocyclic Carbenes**

Alison Levens,[†] Feng An,[‡] Martin Breugst,[§] Herbert Mayr,[‡] and David W. Lupton^{*,†}

[†]School of Chemistry, Monash University, Clayton, Victoria 3800, Australia

[‡]Department Chemie, Ludwig-Maximilians-Universität München, Butenandtstraße 5-13 (Haus F), 81377 München, Germany [§]Department für Chemie, Universität zu Köln, Greinstraße 4, 50939 Köln, Germany

Supporting Information

ABSTRACT: The ability to modulate the nucleophilicity and Lewis basicity of N-heterocyclic carbenes is pivotal to their application as organocatalysts. Herein we examine the impact of the N-substituent on the nucleophilicity and Lewis basicity. Four N-substituents popular in NHC organocatalysis, namely, the N-2,6-(CH₃O)₂C₆H₃, N-Mes, N-4-CH₃OC₆H₄, and N-tertbutyl groups, have been examined and found to strongly affect the nucleophilicity. Thus, the $N-2,6-(CH_3O)_2C_6H_3$ group provides the most nucleophilic imidazolylidene NHC reported and the N-tert-butyl group one of the least. This difference in nucleophilicity is reflected in the catalyst efficiency, as observed with a recently reported trienyl ester rearrangement.

-Heterocyclic carbene (NHC) catalysis provides access to **V** a diverse range of intermediates for reaction discovery.¹ Among others, the acyl anion equivalent,² homoenolates,³ acyl azoliums⁴ and acyl azolium enols/enolates⁵ are now routinely exploited in increasingly sophisticated reaction designs. While other Lewis base catalysts can access some of these classes of intermediates, few access all.⁶ Although the versatility of NHC organocatalysis is striking, practical application requires highly judicious catalyst selection. This can involve variation of the heterocycle, with imidazole, triazole, and thiazole NHC catalysts being common, although increasingly prevalent is N-substituent modification to deliver optimal catalytic activity for a given reaction design.⁷ In the context of acyl anion-mediated reactions, the role of the N-substituent has been investigated.^{7b,c,8} In contrast, an examination of NHC nucleophilicity as a function of the N-substituent, to our knowledge, is yet to be reported. Hence, catalyst design remains largely empirical.

Since 2009 we have reported a series of transformations involving enol esters and enolic anhydrides.⁹ In these studies, and those from others using ester substrates, it is clear that the required NHC catalysts are distinct from those suited to acyl anion-mediated reactions.^{1g,10,11} While Tolman electronic parameters (TEPs), pK_a values, and ¹³C NMR studies give information that is useful for catalyst selection,¹² measures of nucleophilicity are more scarce. Recently, as part of our studies on reactivity scales for nucleophiles and electrophiles, we reported the nucleophilicities of three common NHCs (SIMes, IMes, and TPT; Figure 1).^{13,14} While this study addressed the role of the heterocycle in nucleophilicity, the impact of the Nsubstituent was not addressed. Herein, the dependence of the nucleophilicity on the N-substituent is examined, with the



Figure 1. Overview of the study.

nucleophilicity, as well as the Lewis basicity, of NHCs bearing the $N-2,6-(CH_3O)_2C_6H_3$ (1a), N-Mes (1b), N-4-CH_3OC_6H_4 (1c), and N-tert-butyl groups (1d) determined. These N-substituents have been exploited in recent discoveries in NHC catalysis,^{9h,15–18} particularly using ester substrates. The studies herein demonstrate that the N-substituent plays a significant role in defining the catalyst nucleophilicity, with 3 orders of magnitude rate difference between 1a and 1d. Thus, the impact of the Nsubstituent is at least as significant as the nature of the azolium moiety in determining NHC nucleophilicity.

1b

1a

A series of p-quinone methide electrophiles (i.e., 2, 3, and 4) were previously used to determine the nucleophilicity parameters (N) of SIMes, IMes, and TPT (Figure 1). This was achieved by exploiting the observation that the rate of addition of nucleophiles to Michael acceptors or carbocations can be predicted by eq 1:19

Received: May 29, 2016

ACS Publications © XXXX American Chemical Society

$$\log k_{20^{\circ}\mathrm{C}} = s_{\mathrm{N}}(N+E) \tag{1}$$

In this equation, nucleophiles are characterized by a solventdependent nucleophilicity parameter N and a sensitivity parameter s_N , while electrophiles are characterized by a solvent-independent electrophilicity parameter E. To date, this relationship has been used to examine 1039 nucleophiles and 272 electrophiles.^{20a}

To determine the effect of the N-substituent on the nucleophilicity, studies began with the preparation of the required NHCs by deprotonation of the corresponding imidazolium salts 1a-d·HI with KOt-Bu. Although triazolylidenes are more commonly exploited in NHC catalysis, the impact on the nucleophilicity of changing between a triazolylidene and an imidazolylidene has been previously studied (vide supra), thus allowing trends to be extrapolated. Since the nucleophilicities of the four new NHCs 1a-d are likely to be within the range of those already reported, the reference electrophiles 2a-c, 3a-d, and 4 used previously were employed in this study (Table 1). These references are easily handled, have absorption maxima monitorable in the presence of various NHCs, and cover a suitable range of electrophilicities.

Table 1. Reference *p*-Quinone Methides 2-4

electrophile (E)		R	E^{a}
Ph O Ph Ph	2a	Н	-11.87
	b	OCH ₃	-12.18
	c	$N(CH_3)_2$	-13.39
/Bu	3a	NO_2	-14.36
	b	CH_3	-15.83
	c	OCH_3	-16.11
	d	$N(CH_3)_2$	-17.29
rBu o rBu	4	-	-17.90

^{*a*}Electrophilicity parameters for 2–4 from refs 20b and 20c.

To demonstrate that the outcome of the reaction of NHCs 1a-d with the reference electrophiles are the expected Michael adducts (i.e., 5a-d), preparatory studies were undertaken (Scheme 1). Paralleling earlier reports, the reactions of NHCs 1a-d with *p*-quinone methide 3b gave the adducts 5a-d in high yields as the only isolable products. Structural confirmation was obtained through regular spectroscopic methods as well as with single-crystal X-ray diffraction for 5d. Having confirmed the





formation of the expected adducts, attention was directed toward the rates of these reactions.

The reactions of NHCs 1a-d with the reference *p*-quinone methides 2-4 were studied in THF at 20 °C by photometric monitoring of the disappearance of the colored quinone methides in the presence of a large excess of the NHC to achieve pseudo-first-order conditions. Least-squares fitting of the exponential function $A = A_0 \exp(-k_{obs}t) + C$ to the observed time-dependent absorbances A of the quinone methide provided the first-order rate constants k_{obs} . The slopes of the linear correlations between k_{obs} and the concentrations of 1 correspond to the second-order rate constants k_2 listed in Table 2. The rate

Table 2. Second-Order Rate Constants and Nucleophilicity Parameters for the Reactions of NHCs 1a-d with Reference Electrophiles 2-4 in THF at 20 °C

NHC 1a-d	Е	$k_2 [\mathrm{L} \mathrm{mol}^{-1} \mathrm{s}^{-1}]$	N , $(s_{\rm N})$
/──\ ^H 3CO	2b	1.16×10^{5}	23.00
H ₃ C-N N	2c	2.11×10^{-1}	(0.46)
 1a H₃CO	30 4	1.02×10^{2} 2.27×10^{2}	
H_3C	2a	3.46×10^4	21.50
	2b	1.52×10^4	(0.45)
Пас СНа	2c	3.02×10^{3}	
1b H ₃ C	3a	5.62×10^{3}	
	3b	5.01×10^{2}	
	3c	3.15×10^{2}	
	3d	6.34×10^1	
	4	5.86×10^1	
	2b	6.70×10^{3}	20.41
H ₃ C-N N CH	2c	1.37×10^{3}	(0.46)
10	4	1.41×10^1	
CH ₃	2b	1.46×10^2	16.54
H ₃ C-N, CH ₃	2c	2.42×10^1	(0.47)
1d	3c	3.10	
	4	2.10×10^{-1}	

constants k_2 were then used in association with the electrophilicity parameters of **2**–**4** to construct correlation lines for the four NHCs (Figure 2). As can be seen, in all cases good fits were obtained, which allowed the nucleophilicity parameters to be accurately determined.

The nucleophilicities of these newly characterized carbenes can be compared with those of other NHCs and nucleophilic



Figure 2. Correlations of log k_2 vs electrophilicity parameters for the reactions of NHCs 1a-d with *p*-quinone methides 2-4.

catalysts (Figure 3). From this comparison, it is clear that the nucleophilicity is heavily influenced by the nature of the N-



Figure 3. Nucleophilicities of 1a-d and other catalysts.

substituent. Thus, the most nucleophilic of the NHCs bears an N-2,6-(CH₃O)₂C₆H₃ substituent (i.e., **1a**) and is more nucleophilic than IMes. The least nucleophilic is the *N*-tertbutyl NHC **1d**, which is only moderately more nucleophilic than the Enders carbene (TPT) and DBU. Interestingly, replacing a single mesityl substituent with a methyl group has little impact on the nucleophilicity of the catalyst (IMes, cf. **1b**). Finally, a *para* electron-donating substituent gives an NHC that is less nucleophilic than either the *N*-Mes NHC **1b** or the *N*-2,6-(CH₃O)₂C₆H₃ NHC **1a**.

In previous studies, the Lewis basicities of NHCs have been studied computationally because of an inability to determine equilibrium constants from their interactions with *p*-quinone methides. This is a result of their exceptionally high Lewis basicity. To allow Lewis basicity to be determined in this study, methyl cation affinities (MCAs), as defined in Figure 4, have been calculated for NHCs 1a-d employing MP2/6-31+G-(2d,p)//B86/6-31G(d), a method previously shown to result in reliable MCAs.^{13,21}

Most strikingly, we found that the $N-2,6-(CH_3O)_2C_6H_3$ substituent in **1a** led to the most Lewis basic catalyst, with an MCA some 28 kJ mol⁻¹ higher than that of the variant bearing a single mesityl group (**1b**). Interestingly, the MCA of **1b** was 23 kJ mol⁻¹ lower than that of IMes (MCA = 767 kJ mol⁻¹),¹³ indicating that replacement of a mesityl group with a methyl



Figure 4. Gas-phase methyl cation affinities (MCAs) calculated at the MP2/6-31+G(2d,p)//B98/6-31G(d) level of theory.

group decreases the Lewis basicity while having little bearing on the nucleophilicity. In contrast, switching from a methyl substituent to a *tert*-butyl substituent has very little impact on the Lewis basicity (719 kJ mol⁻¹ vs 718 kJ mol⁻¹ for $R = CH_{3;}$ Figure 4).¹³ The dihedral angle between the imidazole and the aromatic N-substituent indicates significant twisting out of plane in both carbenes 1a and 1b (80.5° and 83.7°) and methyl imidazoles 6a and 6b (69.5° and 88.9°). While some planarization (1a \rightarrow 6a) is observed, it is not sufficient to allow through-space interactions of the *ortho* group and the azolium.

Studies from our group on NHC-catalyzed (4 + 2) annulations show significant sensitivity to the N-substituent.^{9d-f} To further examine this sensitivity, the conversion of triene 7 to cycloadduct 8 was monitored over time using NHCs bearing the four Nsubstituents examined herein (Figure 5). Preliminary studies on



Figure 5. Conversion of ester 7 to diene 8 with catalysts 9a-d.

imidazolium-derived NHCs were hampered by kinetic difficulties due to very high reaction rates. In contrast, monitoring with the triazolium series of NHCs demonstrated that the reaction is successful with the most nucleophilic triazolium-derived catalysts **9a** and **9b** and fails with N-substituents associated with the least nucleophilic NHCs. These results are consistent with our observations that NHC addition to esters is significantly more challenging than that to aldehydes. The success or failure of this reaction is correlated with the catalyst nucleophilicity and not its Lewis basicity, with catalyst **9b** likely to display similar Lewis basicity to **9c** but giving a significantly different reaction outcome.

The capacity of NHCs to enable reactions that are unattainable with other nucleophilic catalysts is associated with their unique Lewis basicity and nucleophilicity profiles. As applications of NHCs in organocatalysis develop, the capacity to modulate these properties takes on increasing importance. In recent years, the N-substituents examined in this report have allowed new discoveries in NHC catalysis. Herein we report that the *N*-2,6- $(CH_3O)_2C_6H_3$ substituent gives one of the most nucleophilic and Lewis basic NHC catalysts, while the impact of remote electron-donating groups on the NHC properties is modest. Finally, hindered alkyl groups give one of the least nucleophilic and Lewis basic NHCs.

The studies reported herein demonstrate that the Nsubstituent plays a significant role in defining catalyst nucleophilicity. Together with the known role of heterocycle selection in nucleophilicity,¹³ these studies demonstrate that significant variation in catalyst properties is possible by manipulation of these two parameters.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b01525.

Procedures and additional data (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: david.lupton@monash.edu.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors acknowledge financial support from the Australian Research Council (DP150101522 and FT110100319), the Deutsche Forschungsgemeinschaft (SFB 749, Project B1), the Fonds der Chemischen Industrie (Liebig Scholarship to M.B.), and the Alexander von Humboldt Foundation (Ludwig Leichardt Memorial Fellowship to D.W.L.). We are grateful to Ms. Nathalie Hampel (LMU) for synthesizing the reference electrophiles and Dr. Peter Mayer (LMU) for the X-ray analysis. This work used the DFG-funded Cologne High Efficiency Operating Platform for Sciences (CHEOPS).

REFERENCES

 (1) For selected recent reviews of NHC catalysis, see: (a) Enders, D.; Niemeier, O.; Henseler, A. Chem. Rev. 2007, 107, 5606. (b) Douglas, J.; Churchill, G.; Smith, A. D. Synthesis 2012, 44, 2295. (c) Bugaut, X.; Glorius, F. Chem. Soc. Rev. 2012, 41, 3511. (d) Izquierdo, J.; Hutson, G. E.; Cohen, D. T.; Scheidt, K. A. Angew. Chem., Int. Ed. 2012, 51, 11686. (e) Ryan, S. J.; Candish, L.; Lupton, D. W. Chem. Soc. Rev. 2013, 42, 4906. (f) Hopkinson, M. N.; Richter, C.; Schedler, M.; Glorius, F. Nature 2014, 510, 485. (g) Chauhan, P.; Enders, D. Angew. Chem., Int. Ed. 2014, 53, 1485. (h) Flanigan, D. M.; Romanov-Michailidis; White, N. A.; Rovis, T. Chem. Rev. 2015, 115, 9307.

(2) For the seminal contributions of Breslow and Ukai, see: Breslow, R. *J. Am. Chem. Soc.* **1958**, *80*, 3719 and references therein.

(3) Early examples of homolenolate chemistry: (a) Sohn, S. S.; Rosen, E. L.; Bode, J. W. *J. Am. Chem. Soc.* **2004**, *126*, 14370. (b) Burstein, C.; Glorius, F. *Angew. Chem., Int. Ed.* **2004**, *43*, 6205.

(4) Early examples of unsaturated acyl azolium chemistry: (a) Zeitler, K. Org. Lett. **2006**, *8*, 637. (b) Maki, B. E.; Chan, A.; Phillips, E. M.; Scheidt, K. A. Org. Lett. **2007**, *9*, 371. (c) Ryan, S. J.; Candish, L.; Lupton, D. W. J. Am. Chem. Soc. **2009**, *131*, 14176. (d) De Sarkar, S.; Studer, A. Angew. Chem., Int. Ed. **2010**, *49*, 9266.

(5) Early acyl azolium enolate/enol chemistry: (a) Chow, K. Y.-K.; Bode, J. W. J. Am. Chem. Soc. **2004**, *126*, 8126. (b) Reynolds, N. T.; Read de Alaniz, J.; Rovis, T. J. Am. Chem. Soc. **2004**, *126*, 9518.

(6) For reaction design with different nucleophilic catalysts, see: Candish, L.; Nakano, Y.; Lupton, D. W. *Synthesis* **2014**, *46*, 1823.

(7) (a) Rovis, T. *Chem. Lett.* **2008**, *37*, 2. (b) Mahatthananchai, J.; Bode, J. W. *Chem. Sci.* **2012**, *3*, 192. (c) Collett, C. J.; Massey, R. S.; Maguire, O. R.; Batsanov, A. S.; O'Donoghue, A. C.; Smith, A. D. *Chem. Sci.* **2013**, *4*, 1514. (d) Collett, C. J.; Massey, R. S.; Taylor, J. E.; Maguire, O. R.; O'Donoghue, A. C.; Smith, A. D. *Angew. Chem., Int. Ed.* **2015**, *54*, 6887.

(8) For selected mechanistic contributions to the field, see: (a) Paul, M.; Breugst, M.; Neudörfl, J.-M.; Sunoj, R. B.; Berkessel, A. J. Am. Chem. Soc. 2016, 138, 5044. (b) Yatham, V. R.; Neudörfl, J.-M.; Schlörer, N. E.; Berkessel, A. Chem. Sci. 2015, 6, 3706. (c) Berkessel, A.; Yatham, V. R.; Elfert, S.; Neudörfl, J.-M. Angew. Chem., Int. Ed. 2013, 52, 11158. (d) Berkessel, A.; Elfert, S.; Etzenbach-Effers, K.; Teles, J. H. Angew. Chem., Int. Ed. 2010, 49, 7120. (e) Zhao, X.; Glover, G. S.; Oberg, K. M.; Dalton, D. M.; Rovis, T. Synlett 2013, 24, 1229. (f) Moore, J. L.; Silvestri, A. P.; Read de Alaniz, J.; DiRocco, D. A.; Rovis, T. Org. Lett. 2011, 13, 1742. (g) Cohen, D. T.; Johnston, R. C.; Rosson, N. T.; Cheong, P. H.-Y.; Scheidt, K. A. Chem. Commun. 2015, 51, 2690. (h) Johnston, R. C.; Cohen, D. T.; Eichman, C. C.; Scheidt, K. A.; Cheong, P. H.-Y. Chem. Sci. 2014, 5, 1974. (i) Samanta, R. C.; Maji, B.; De Sarkar, S.; Bergander, K.; Fröhlich, R.; Mück-Lichtenfeld, C.; Mayr, H.; Studer, A. Angew. Chem., Int. Ed. 2012, 51, 5234. (j) Ryan, S. J.; Stasch, A.; Paddon-Row, M. N.; Lupton, D. W. J. Org. Chem. 2012, 77, 1113.

(9) For selected studies with esters from our group, see: (a) Reference 4c. (b) Kowalczyk, M.; Lupton, D. W. Angew. Chem., Int. Ed. 2014, 53, 5314. (c) Candish, L.; Levens, A.; Lupton, D. W. J. Am. Chem. Soc. 2014, 136, 14397. (d) Candish, L.; Levens, A.; Lupton, D. W. Chem. Sci. 2015, 6, 2366. (e) Levens, A.; Zhang, C.; Candish, L.; Forsyth, C. M.; Lupton, D. W. Org. Lett. 2015, 17, 5332. For selected studies using activated acids, see: (f) Ryan, S. J.; Candish, L.; Lupton, D. W. J. Am. Chem. Soc. 2011, 133, 4694. (g) Candish, L.; Lupton, D. W. J. Am. Chem. Soc. 2013, 135, 58. (h) Candish, L.; Forsyth, C. M.; Lupton, D. W. Angew. Chem., Int. Ed. 2013, 52, 9149.

(10) For examples from Chi's group that demonstrate sensitivity to the N-substituent, see: (a) Hao, L.; Chen, S.; Xu, J.; Tiwari, B.; Fu, Z.; Li, T.; Lim, J.; Chi, Y. R. *Org. Lett.* **2013**, *15*, 4956. (b) Xu, J.; Jin, Z.; Chi, Y. R. *Org. Lett.* **2013**, *15*, 5028. (c) Chen, S.; Hao, L.; Zhang, Y.; Tiwari, B.; Chi, Y. R. *Org. Lett.* **2013**, *15*, 5822.

(11) For recent reports on NHC catalysis via anhydrides, see: (a) Lee, A.; Younai, A.; Price, C. K.; Izquierdo, J.; Mishra, R. K.; Scheidt, K. A. J. Am. Chem. Soc. **2014**, 136, 10589. (b) Chen, X.-Y.; Gao, Z.-H.; Song, C.-Y.; Zhang, C.-L.; Wang, Z.-X.; Ye, S. Angew. Chem., Int. Ed. **2014**, 53, 11611. (c) Jin, Z.; Chen, S.; Wang, Y.; Zheng, P.; Yang, S.; Chi, Y. R. Angew. Chem., Int. Ed. **2014**, 53, 13506.

(12) For a review of NHC properties, see: (a) Dröge, T.; Glorius, F. Angew. Chem., Int. Ed. **2010**, 49, 6940. For computational studies on π -acidity, see: (b) Andrada, D. M.; Holzmann, N.; Hamadi, T.; Frenking, G. Beilstein J. Org. Chem. **2015**, 11, 2727 and references therein.

(13) Maji, B.; Breugst, M.; Mayr, H. Angew. Chem., Int. Ed. 2011, 50, 6915.

(14) For a review of the use of this scale in organocatalysis, see: Mayr, H.; Lakhdar, S.; Maji, B.; Ofial, A. R. *Beilstein J. Org. Chem.* **2012**, *8*, 1458.

(15) For the N-2,6-(CH₃O)₂C₆H₃ group, see: Liu, F.; Bugaut, X.; Schedler, M.; Fröhlich, R.; Glorius, F. Angew. Chem., Int. Ed. **2011**, 50, 12626.

(16) For the N-2,4,6-(CH₃)₃C₆H₂ group in enantioselective NHC organocatalysis, see: He, M.; Struble, J. R.; Bode, J. W. *J. Am. Chem. Soc.* **2006**, *128*, 8418.

(17) For the N-4-CH₃OC₆H₄ group in enantioselective catalysis, see: Nakano, Y.; Lupton, D. W. Angew. Chem., Int. Ed. **2016**, 55, 3135.

(18) The $N-C_6F_5$ group, which is exploited extensively in acyl anion chemistry, was also examined, but difficulties in preparing the carbene in high purity precluded its inclusion in this study.

(19) (a) Mayr, H.; Patz, M. Angew. Chem., Int. Ed. Engl. 1994, 33, 938.
(b) Mayr, H.; Bug, T.; Gotta, M. F.; Hering, N.; Irrgang, B.; Janker, B.; Kempf, B.; Loos, R.; Ofial, A. R.; Remennikov, G.; Schimmel, H. J. Am. Chem. Soc. 2001, 123, 9500. (c) Mayr, H.; Ofial, A. R. J. Phys. Org. Chem. 2008, 21, 584.

(20) (a) Access to all constants is provided at: Mayr, H. Database of Nucleophilicties and Electrophilicities. http://www.cup.uni-muenchen. de/oc/mayr/DBintro.html (accessed July 3, 2016). (b) For quinone methides 2b-c, 3b-d and 4, see: Lucius, R.; Loos, R.; Mayr, H. Angew. Chem., Int. Ed. 2002, 41, 91. (c) For quinone methides 2a and 3a, see: Richter, D.; Hampel, N.; Singer, T.; Ofial, A. R.; Mayr, H. Eur. J. Org. Chem. 2009, 2009, 3203.

(21) (a) Wei, Y.; Sastry, G. N.; Zipse, H. J. Am. Chem. Soc. 2008, 130, 3473. (b) Wei, Y.; Singer, T.; Mayr, H.; Sastry, G. N.; Zipse, H. J. Comput. Chem. 2008, 29, 291. (c) Lindner, C.; Maryasin, B.; Richter, F.; Zipse, H. J. Phys. Org. Chem. 2010, 23, 1036.