COMMUNICATION

Reaction of gold(III) oxo complexes with alkenes. Synthesis of unprecedented gold alkene complexes, $[Au(N,N)(alkene)][PF_6]$. Crystal structure of $[Au(bipy^{ip})(\eta^2-CH_2=CHPh)][PF_6]$ (bipy^{ip} = 6-isopropyl-2,2'-bipyridine)[†]

Maria A. Cinellu,*a Giovanni Minghetti,a Sergio Stoccoro,a Antonio Zuccaa and Mario Manasserob

^a Dipartimento di Chimica, Università di Sassari, via Vienna 2, I-07100 Sassari, Italy. E-mail: cinellu@uniss.it

^b Dipartimento di Chimica Strutturale e Stereochimica Inorganica, Università di Milano, Centro CNR, via Venezian 21, I-20133 Milano, Italy. E-mail: m.manassero@istm.cnr.it

Received (in Cambridge, UK) 2nd April 2004, Accepted 21st May 2004 First published as an Advance Article on the web 17th June 2004

Gold alkene complexes $[Au(bipy^R)(\eta^2\text{-}alkene)][PF_6]$ (bipy^R = 6-alkyl-2,2'-bipyridine) have been obtained by reaction of gold(111) oxo complexes $[Au_2(bipy^R)_2(\mu\text{-}O)_2][PF_6]_2$ with alkenes. The crystal structure of the styrene adduct $[Au(bipy^{ip})(\eta^2\text{-}CH_2=CHPh)][PF_6]$ (bipy^{ip} = 6-isopropyl-2,2'-bipyridine) has been solved by X-ray analysis.

Gold(III) oxo complexes¹ are a new entry into the relatively small family of late transition metal oxo complexes.² The chemistry of this class of compounds has been of interest for many years. Among the various aspects of the reactivity displayed by metal oxo complexes is the capability of transferring oxygen atoms to organic and inorganic substrates.^{2,3}

In the course of our investigations of the reactivity of the oxo complexes $[Au_2(bipy^R)_2(\mu-O)_2][PF_6]_2$ (bipy^R = 6-alkyl-2,2'-bipyridine) we found that PPh₃ is stoichiometrically converted into OPPh₃; the gold(1) complex $[Au(PPh_3)_2][PF_6]$ and the free bipyridine ligand are also quantitatively obtained from the reaction.^{1*a*} In view of these encouraging results we decided to extend this reaction to alkenes. Alkene oxidation by oxygen atom transfer reaction from late transition metal oxo complexes is, at present, an objective of particular interest.

The μ -oxo complexes $[Au_2(bipy^R)_2(\mu-O)_2][PF_6]_2$ **1** (R = Me **a**, CH₂Me **b**, CHMe₂ **c**, CH₂CMe₃ **d**) react with excess alkene in acetone or acetonitrile solution under different experimental conditions to give, albeit in low yield, unexpected alkene adducts, $[Au(bipy^R)(alkene)][PF_6]$ (alkene = styrene **2**, α -methylstyrene, ethylene and norbornene).^{4,5} Few examples of alkene complexes obtained by reaction of a late transition metal oxo complex have been reported.⁶

The reaction with styrene was used as a model and studied with all the complexes in order to assess the possible influence of the alkyl substituent at the 6 position on the reactivity of the oxo complexes.[‡] In no case is the reaction quantitative—average yield is in the range 15–45%. Most of the starting material is recovered when the reaction is carried out in anhydrous solvents, while complete conversion of the oxo complexes is accomplished in the presence of water, after a long reaction time.[‡] Variable amounts of metallic gold and organic products arising from oxidation of alkenes are also formed in all cases.^{4,5} To the best of our knowledge, alkene oxidation by an isolated late transition metal oxo complex is still rare.^{6b,7}

The thermally stable styrene adducts $[Au(bipy^R)(\eta^2-CH_2=CHPh)][PF_6]$ (**2a–d**)† give satisfactory elemental analyses and their molecular ions M^+ have been detected by FAB mass spectrometry. The ¹H NMR spectra at room temperature, recorded in various solvents, show that both the CH and CH₂ resonances are shifted upfield compared to free styrene. For example, in CDCl₃ the methyne proton resonates at δ 5.43 ($\Delta\delta$ – 1.29) and the methylene protons at δ 4.07 ($\Delta\delta$ – 1.68 ppm) and 4.03 ($\Delta\delta$ – 1.21 ppm). The

† Electronic supplementary information (ESI) available: experimental details. See http://www.rsc.org/suppdata/cc/b4/b404890c/

resonances of the corresponding carbon atoms are likewise shifted upfield with coordination-induced shift $\Delta\delta$ –52.4 ppm for the methyne and –56.4 ppm for the methylene carbon. These upfield shifts are similar to those found in the comparable three-coordinate copper(1) olefin complexes supported by chelating nitrogen ligands ($\Delta\delta$ (H) –0.1 to –2.0 ppm; $\Delta\delta$ (C) –40 to –50 ppm).^{8,9} Addition of one equivalent of styrene to a CD₂Cl₂ solution of **2a** causes broadening of the vinylic protons of the coordinated styrene, indicating fast intermolecular exchange at room temperature.

Variable temperature ¹H NMR spectra of complex **2c** indicate at room temperature a dynamic process involving rotation of the coordinated styrene around the metal–olefin bond. The calculated $\Delta G_{rot}^{\ddagger}$ value (50.3 kJ·mol⁻¹ at 243 K)† is in the range previously reported for similar d¹⁰ olefin complexes with α -diimmine ligands.^{8e,9a,b,10}

Olefin gold complexes have been known for many years,¹¹ however, only a small number of isolable complexes have been reported¹² and only a few of them have been structurally characterized.¹³ Furthermore the available X-ray data are relevant to species incomparable to compounds **2a–2d**.

The structure of **2c**§ consists of the packing of $[Au(bipy^{ip})(\eta^2$ styrene)]⁺ cations, PF₆⁻ anions and MeCN molecules in the molar ratio 2:2:1. Fig. 1 shows an ORTEP view of the cation, and principal bond parameters are reported in the figure caption. The orientation of the olefin is coplanar with the bipyridine backbone, which chelates the gold centre with a bite angle of $75.1(2)^{\circ}$. The dihedral angle between the metal coordination plane and the C(14)-C(15)-C(16) plane is 77.2(3)° which means that there is a bending back angle of 12.8(3) ° of the phenyl ring with respect to the ideal orthogonal coordination. Gold-carbon distances, Au-C(14) 2.114(6) and Au-C(15) 2.098(5) Å, are slightly shorter or similar to those found in two gold(I) alkene complexes (2.21(2), 2.15(2) Å^{13c} and 2.14(4), 2.11(4) Å^{13b}) (consider the high standard deviations of the four latter distances). The bond length between Au and the CH carbon atom, 2.114(6) Å, is slightly longer than that of the other Au-C contact, 2.098(5) Å. This is a common feature of metalcoordinated styrene, compare with e.g. Cu-C distances of 2.014(5) and 1.985(6) Å in the cationic Cu(I) styrene complex [Cu(bipy)(η^2 styrene)][ClO₄] **3**.^{8b} The C(14)–C(15) distance, 1.384(8) Å, is slightly longer than in free styrene (1.346(20) Å)¹⁴ and in 3 (1.358(10) Å)^{8b} and comparable to that found in the neutral β diketiminate complex [$\hat{C}u(Me_2NN)(\eta^2-CH_2=CHPh)$] (1.373(6) Å).^{8e} The Au–N distances, 2.150(3) and 2.217(5) Å are very long, probably because of the trans-influence of the styrene molecule. The elongation of the Au–N(2) bond, 2.217(5) Å, compared to the Au–N(1) bond, 2.150(3) Å, is probably due to the steric hindrance of the bulky isopropyl substituent. A similar effect has been found previously, for instance, in 1d.^{1c}

The coordinated styrene is readily displaced from 2d by triphenylphosphine, as shown by a proton spectrum recorded straight after the addition of one equivalent of PPh₃. [Au(PPh₃)₂][PF₆] is quantitatively isolated from the reaction with two equivalents of PPh₃.

1618



Fig. 1 ORTEP view of the [Au(bipy^{ip})(η^2 -styrene)]⁺ cation. Ellipsoids are drawn at the 30% probability level. Selected bond distances (Å) and angles (°): Au–N(1) 2.150(3), Au–N(2) 2.217(5), Au–C(14) 2.114(6), Au–C(15) 2.098(5), C(14)–C(15) 1.384(8), C(14)–C(16) 1.488(6) Å, N(1)–Au–N(2) 75.1(2), N(1)–Au–C(14) 119.6(2), N(1)–Au–C(15) 157.8(2), N(2)–Au–C(14), 165.4(2), N(2)–Au–C(15) 127.0(2), C(14)–Au–C(15) 38.4(2), C(15)–C(14)–L(16) 126.3(5)°.

On the whole the spectroscopic and structural data, taken together, point to a non negligible π contribution in the olefin–gold bond.

We are grateful to the University of Sassari for financial support (60%) and to the Regione Autonoma della Sardegna (RAS) for a grant to M.A.C. (L.R.9/8/1950, n.43).

Notes and references

[‡] Reactions of compounds **1a–1d** with styrene. *Anhydrous solvent*: styrene (2 mmol) was added to a solution of **1** (0.2 mmol) in acetonitrile (40 cm³) (**1a** and **1b**) or acetone (40 cm³) (**1c** and **1d**). The resulting yellow solution was stirred for 3 days at room temperature and then filtered through Celite. The solution was evaporated to dryness and the residue extracted with CHCl₃ (3 × 15 cm³). The combined chloroform extracts were filtered and concentrated to a small volume. Addition of diethyl ether gave a whitish precipitate of compound **2** (**2a**, 15%; **2b**, 20%; **2c** 25%; **2d**, 45%). Unreacted complex **1** was recovered from the insoluble residue (**1a**, 66%; **1b**, 61%; **1c**, 50%; **1d**, 30%).

Aqueous solvent: to a solution of **1a** (0.320 g, 0.3 mmol) in acetonitrile (50 cm³) styrene (6 mmol) and water (5 cm³) were added. The resulting solution was stirred under Ar for 7 days at room temperature. Concentration of the filtered solution gave a white precipitate of **2a** which was recovered by filtration under vacuum. Recrystallization from dichloromethane gave the analytical sample (0.130 g, 35%). Compounds **2c** and **2d** were obtained similarly from **1c** and **1d**, respectively. In contrast to **2a**, **2c** and **2d** were extracted with CHCl₃ (3 × 15 cm³) after evaporation to dryness of the filtered solution. (**2c**, 31 %; **2d**, 46%).

§ Crystal data for **2c**·0.5MeCN: C₂₂H_{23.5}Au₁F₆N_{2.5}P, $M_r = 664.88$, monoclinic, space group $P2_1/c$ (no. 14), a = 13.717(1), b = 12.833(1), c = 15.272(2) Å, $\beta = 114.15(1)^\circ$, U = 2453.1(4) Å³, Z = 4, $d_c = 1.800$ g cm⁻³, T = 223 K, $\mu = 61.1$ cm⁻¹, F(000) = 1284. Reflections measured 33227, independent 5027 with $R_{int} = 0.027$. Final R_2 (F^2 , all reflections) = 0.054, $R_{2w} = 0.090$, conventional $R_1 = 0.032$ for 291 variables. Bruker SMART CCD area-detector, Mo-K α radiation ($\lambda = 0.71073$ Å), ω scan mode, $\theta_{min} = 3^\circ$, $\theta_{max} = 26^\circ$. Structure solved by direct methods and refined by full-matrix least squares. Atom C(13) is disordered and split into two half atoms, C(13a) and C(13b), with occupancy factors = 0.50; only C(13a) has been drawn in Fig. 1. The half MeCN molecule is disordered around an inversion centre. The programme used was Personal SDP on a

Pentium III computer. CCDC 236738. See http://www.rsc.org/suppdata/cc/ b4/b404890c/ for crystallographic data in .cif or other electronic format.

- (a) M. A. Cinellu and G. Minghetti, Gold Bull., 2002, 35, 11; (b) M. A. Cinellu, G. Minghetti, M. V. Pinna, S. Stoccoro, A. Zucca and M. Manassero, J. Chem. Soc., Dalton Trans., 2000, 1261; (c) M. A. Cinellu, G. Minghetti, M. V. Pinna, S. Stoccoro, A. Zucca, M. Manassero and M. Sansoni, J. Chem. Soc., Dalton Trans., 1998, 1735; (d) M. A. Cinellu, G. Minghetti, M. V. Pinna, S. Stoccoro, A. Zucca and M. Manassero, Chem. Commun., 1998, 2397.
- 2 P. R. Sharp, J. Chem. Soc., Dalton Trans., 2000, 2647; P. R. Sharp, Comments Inorg. Chem., 1999, 21, 85.
- 3 Active Oxygen in Chemistry, ed. C. S. Foote, J. S. Valentine, A. Greenberg and J. F. Liebman, Chapman and Hall, London, 1995; Active Oxygen in Biochemistry, ed. J. S. Valentine, C. S. Foote, A. Greenberg and J. F. Liebman, Chapman and Hall, London, 1995; Biomimetic Oxidations Catalyzed by Transition Metal Complexes, ed. B. Meunier, Imperial College Press, London, 2000; Metal-Oxo and Metal-Peroxo Species in Catalytic Oxidations, ed. B. Meunier, Springer, Berlin, 2000; vol. 97; R. H. Holm, Chem. Rev., 1987, 87, 1401; L. Que, Jr. and W. B. Tolman, Angew. Chem., Int. Ed., 2002, 41, 1114; M. Taki, S. Itoh and S. Fukuzumi, J. Am. Chem. Soc., 2002, 124, 998 and references therein.
- 4 Part of this work has been presented in oral form: M. A. Cinellu, G. Minghetti, D. Bacciu, S. Stoccoro, A. Zucca and M. Manassero, *Abstracts of Papers, Gold 2003, New Industrial Applications of Gold*, Vancouver, Canada, September 28–October 1, 2003, No.1113.
- 5 Full characterization of complexes with α-methylstyrene, ethylene and norbornene is on-going and will be reported in a full paper.
- 6 (a) B. Flint, J.-J. Li and P. R. Sharp, Organometallics, 2002, 21, 997; (b) E. Szuromi, H. Shan and P. R. Sharp, J. Am. Chem. Soc., 2003, 125, 10522.
- 7 T. Hosokawa, M. Takano and S.-I. Murahashi, J. Am. Chem Soc., 1996, 118, 3990.
- 8 (a) M. Munakata, S. Kitagawa, S. Kosome and A. Asahara, *Inorg. Chem.*, 1986, **25**, 2622 and references therein (b) H. Masuda, K. Machida, M. Munakata, S. Kitagawa and H. Shimono, *J. Chem. Soc., Dalton Trans.*, 1988, 1907; (c) L. Cavallo, M. E. Cucciolito, A. De Martino, F. Giordano, I. Orabona and A. Vitagliano, *Chem. Eur. J.*, 2000, **6**, 1127; (d) B. F. Straub, F. Eisenträger and P. Hoffmann, *Chem. Commun.*, 1999, 2507; (e) X. Dai and T. H. Warren, *Chem. Commun.*, 2001, 1998.
- 9 Larger values are found for analogous d¹⁰ metal complexes (M = Pd, Pt) with more electron accepting alkenes ($\Delta\delta$ (H) -3.0 to -3.5 ppm; $\Delta\delta$ (C) -90 to -120 ppm). See for example: (a) R. van Asselt, C. J. Elsevier, W. J. J. Smeets and A. L. Spek, *Inorg. Chem.*, 1994, **33**, 1521; (b) K. J. Cavell, D. J. Stufken and K. Vrieze, *Inorg. Chim. Acta*, 1980, **47**, 67; (c) M. W. van Laren, M. A. Duin, C. Klerk, M. Naglia, D. Rogolino, P. Pelagatti, A. Bacchi, C. Pelizzi and C. J. Elsevier, *Organometallics*, 2002, **21**, 1546; (d) M. L. Ferrara, F. Giordano, I. Orabona, A. Panunzi and F. Ruffo, *Eur. J. Inorg. Chem.*, 1999, 1939; (e) L. Canovese, F. Visentin, G. Chessa, C. Santo, P. Uguagliati, L. Maini and M. Polito, *J. Chem. Soc., Dalton Trans.*, 2002, 3696; (f) M. L. Ferrara, I. Orabona, F. Ruffo, M. Funicello and A. Panunzi, *Organometallics*, 1998, **17**, 3832.
- 10 L. Canovese, F. Visentin, P. Uguagliati and B. Crociani, J. Chem. Soc., Dalton Trans., 1996, 1921.
- 11 A. J. Chalk, J. Am. Chem. Soc., 1964, 86, 4733.
- H. Schmidbaur, A. Grohmann and M. E. Olmos in *Gold Progress in Chemistry, Biochemistry and Technology*, ed. H. Schmidbaur, John Wiley and Sons. New York, 1999, pp. 728–730; A. Grohmann and H. Schmidbaur, 'Gold' in *Comprehensive Organometallic Chemistry*, ed. J. L. Wardell, E. W. Abel, F. G. A. Stone and G. Wilkinson, F. Gold', in *Comprehensive Organometallic Chemistry*, ed. J. Puddephatt, 'Gold', in *Comprehensive Organometallic Chemistry*, ed. G. Wilkinson, F. G. A. Stone and E. W. Abel, Pergamon Press, Oxford, 1982, vol. 2, pp. 809–812; R. J. Puddephatt, *The Chemistry of Gold*, Elsevier, The Netherlands, 1978, pp. 148–151.
 (a) M. Håkansson, H. Eriksson and S. Janger, *J. Organomet. Chem.*,
- 13 (a) M. Håkansson, H. Eriksson and S. Janger, J. Organomet. Chem., 2000, 602, 133; (b) R. M. Dávila, R. J. Staples and J. P. Fackler, Jr., Organometallics, 1994, 13, 418; (c) D. Belli Dell'Amico, F. Calderazzo, R. Dantona, J. Strähle and H. Weiss, Organometallics, 1987, 6, 1207.
- 14 J. C. Cochran, K. Hagen, G. Paulen, Q. Shen, S. Tom, M. Traetteberg and C. Wells, *J. Mol. Struct.*, 1997, **413–414**, 313.