View Article Online

Dalton Transactions

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. Arévalo, C. M. Vogels, G. MacNeil, L. Riera, J. Perez and S. Westcott, *Dalton Trans.*, 2017, DOI: 10.1039/C7DT01549F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the **author guidelines**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the ethical guidelines, outlined in our <u>author and reviewer resource centre</u>, still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/dalton

Dalton Transactions

PAPER

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/



Rebeca Arévalo,^a Christopher M. Vogels,^b Gregory A. MacNeil,^b Lucía Riera,^c Julio Pérez,^{ac} and Stephen A. Westcott*^b

The first examples for the rhenium-catalysed hydroboration of aldehydes, ketones and aldimines, including heteroaromatic quinoline, are reported herein. Reactions are remarkably chemoselective and tolerant of several functional groups. A wide array of rhenium complexes were efficient pre-catalysts for these hydroborations, including new low-valent complexes of the formula $[Re(N-N)(CO)_3(L)]X$ (N-N = bipy derivative, L = labile ligand/solvent, and X = $[BAr^{f}_4]^-$ and $[B(3,5-di-tBu-cat)_2]^-$), which have been characterized fully including an X-ray diffraction study for $[Re(bipy)(CO)_3(quin)][BAr^{f}_4]$ (2). A new silver spiroboronate ester Ag[B(3,5-di-tBu-cat)_2](NCCH_3)_3 (3) was prepared and characterized fully, including an X-ray diffraction study, and used to make one of the new rhenium complexes.

Introduction

Published on 30 May 2017. Downloaded by State University of New York at Binghamton on 31/05/2017 11:35:17.

The catalysed hydroboration of unsaturated organic molecules using either catecholborane (HBcat: cat = $1,2-O_2C_6H_4$) or pinacolborane (HBpin: pin = 1,2-O₂C₂Me₄) offers the synthetic chemist a gentle and efficient methodology for the reduction of double and triple bonds. Depending upon the nature of the substrate, fine-tuning of the catalyst system and the borane, these catalysed reactions can generate products with complementary, opposite, chemo-, or regio-. and stereoselectivities compared to those obtained in conventional hydroborations using H₃B•LB (LB = Lewis base), alkyl- or dialkyl boranes.¹ Although catalysed hydroborations have traditionally employed late metal catalysts (Ru, Rh, Ir, etc) and have focused on the reduction of alkenes and alkynes, a considerable amount of research has recently investigated the possibility of using alternate catalyst systems for the hydroboration of aldehydes and ketones as a gentle route for generating the corresponding alcohols (Scheme 1).² Not surprising the boron usually ends up bound to the electronegative oxygen atom of the aldehyde or ketone while the basic hydride adds to the carbon atom.² Indeed, a number of different metal complexes and Lewis acids have been found to be effective pre-catalysts for the reduction of C=O double bonds, including Zr,³ Ti,⁴ Mn,⁵ Fe,⁶ Co,⁷ Ni,⁸ Cu,⁹ Zn,¹⁰ Mo,¹¹ Ru,¹² Rh/Ir,¹³ Ge/Sn,¹⁴ P,¹⁵ Al,¹⁶ Ga/In,¹⁷ Mg,¹⁸ and La.19



Scheme 1 Hydroboration and Oxidative Workup of Ketones and Aldehydes using HBpin.

It is somewhat surprising, therefore, that the use of rhenium complexes in catalysed hydroboration reactions has been largely ignored. Indeed, the only reported examples so far have been by Fernandes and co-workers who have elegantly demonstrated that high-valent rhenium complexes, such as $[(PPh_3)_2Re(O)_2I]$, are effective pre-catalysts for the reduction or sulfoxides using either HBcat or HBpin.²⁰ These reactions are believed to proceed via initial dissociation of the phosphine ligands and coordination of two sulfoxide groups. The next step involves activation of the B-H bond by a formal addition of the borane to the rhenium-oxide bond to give the corresponding rhenium hydrides where the Lewis acid boryl group has added to the oxygen atom to form transient Re-OBcat or Re-OBpin appendages. In support of this mechanism, Fernandes and coworkers have found that stoichiometric addition of HBcat and HBpin to $[(PPh_3)_2Re(O)_2I]$ (1) cleanly affords the metal hydrides (trans-[(Ph₃P)₂(O)(I)Re(H)OBcat] and trans-[(Ph₃P)₂(O)(I)Re(H)OBpin]) in high yields (Scheme 2). Following this remarkable chemistry, and considering our interest in rhenium chemistry²¹ and developing novel borylation strategies,²² we decided to explore the possibility of using various rhenium complexes as pre-catalysts for the hydroboration of aldehydes, ketones and aldimines, the results of which are presented within.

Dalton Trans., 2017, 46, x-y | 1

^{a.} Departamento de Química Orgánica e Inorgánica, Universidad de Oviedo, Julián Clavería, 8, 33006, Oviedo, Spain.

b. Department of Chemistry and Biochemistry, Mount Allison University, Sackville, New Brunswick, E4L 1G8, Canada.

^{c.} Centro de Investigación en Nanomateriales y Nanotecnología (CINN), CSIC-

Universidad de Oviedo-Principado de Asturias, Avenida de la Vega 4–6, 33940, El Entrego, Spain.

⁺ Electronic Supplementary Information (ESI) available: experimental and catalytic data provided. See DOI: 10.1039/x0xx00000x

Published on 30 May 2017. Downloaded by State University of New York at Binghamton on 31/05/2017 11:35:17.

Results and discussion

Synthesis of rhenium complexes

It is well known that rhenium complexes in various oxidation states can effectively be used as pre-catalysts for the hydrosilylation of carbonyl compounds.²³ Although a number of simple neutral rhenium complexes are readily-available, we decided to prepare a series of low-valent rhenium cationic salts of the general formula [Re(bipy)(CO)₃(L)]X (Lewis-base, Scheme 3) in an effort to expand the range of rhenium complexes investigated for the analogous hydroboration reactions. For complete spectroscopic data and catalytic results on these species, please see the Supporting Information section. These complexes have been designed to be strong Lewis acids, combining labile ligands and non-coordinating anions with a cationic metal centre and strongly π -acceptor carbonyl ligands.²¹ Variation of the labile monodentate L, allowed us to fine-tune the physicochemical properties of the rhenium complexes. All new complexes have been characterized fully by a number of physical techniques including multinuclear NMR spectroscopy, FT-IR spectroscopy, elemental analyses and X-ray diffraction in the case of 2. The molecular structure of 2 is shown in Fig. 1 and bond distances and angles are similar to those reported earlier for related complexes.²¹



Page 2 of 9

the reaction of $B_2(3,5-di-tBu-cat)_3^{25}$ with $[Ag(acac)_{ni}]_{ni}$ acetonitrile. The reaction proceeded in the dark for 18 h to give the desired silver salt in moderate yield (61%) along with concomitant formation of (acac)B(3,5-di-tBu-cat). The ¹¹B NMR spectrum for Ag[B(3,5-di-tBu-cat)_2](NCCH_3)_3



Fig. 1 The molecular structure of **2** with thermal ellipsoids shown at the 50% probability level. The anion has been omitted for clarity. Selected bond distances (Å) and angles (°): Re(1)-C(23) 1.924(4), Re(1)-C(21) 1.925(6), Re(1)-C(22) 1.939(5), Re(1)-N(1) 2.171(3), Re(1)-N(2) 2.181(3), Re(1)-N(3) 2.257(4); C(22)-Re(1)-N(1) 173.88(15), C(23)-Re(1)-N(2) 173.74(15), C(21)-Re(1)-N(3) 172.42(16).

shows the expected sharp singlet at 14.1 ppm which is typical for four-coordinate boron bound to two catecholato groups. 25 To confirm the formation of this silver salt we have carried out a single crystal X-ray diffraction study on 3, the molecular structure of which is shown in Fig. 2. As expected, the borate anion lies in a slightly distorted tetrahedral environment. The silver cation has three molecules of acetonitrile arranged in a distorted tetrahedral geometry as there appears to be significant interaction with one of the oxygen atoms of the catecholato groups of the bulky [B(3,5-di-tBu-cat)2]. Indeed, the long silver-oxygen distance of 2.4333(16) Å is close to that found in related structures containing bound THF ligands, demonstrating that the anion is only weakly coordinated to the silver cation in the solid state.²⁶ No evidence was found using multinuclear NMR spectroscopy for any such coordination of the bulky [B(3,5-di-tBu-cat)2] anion to the silver cation in solution.



Fig. 2 The molecular structure of $Ag[B(3,5-di-tBu-cat)_2](NCCH_3)_3$ (3) with thermal ellipsoids shown at the 50% probability level. Selected bond distances (Å) and angles (°): Ag(1)-N(1) 2.240(2), Ag(1)-N(3) 2.250(2), Ag(1)-N(2) 2.274(2), Ag(1)-O(4) 2.4333(16); N(1)-Ag(1)-N(3) 130.83(9), N(1)-Ag(1)-N(2) 109.49(9), N(3)-Ag(1)-N(2) 108.33(8), N(1)-Ag(1)-O(4) 101.86(7), N(3)-Ag(1)-O(4) 97.60(7), N(2)-Ag(1)-O(4) 104.84(7).

Published on 30 May 2017. Downloaded by State University of New York at Binghamton on 31/05/2017 11:35:17.

Dalton Transactions

Furthermore, we have found that addition of **3** in CH₃CN to $[Re(bipy)(CO)_3Br]$ resulted in the formation of $[Re(bipy)(CO)_3(CH_3CN)][B(3,5-di-tBu-cat)_2]$ (**4**, Scheme 3b) in high yields (88%) along with concomitant formation of AgBr. With these new rhenium complexes in hand, along with a variety of known compounds, we then decided to examine their potential to facilitate the hydroboration of compounds containing carbonyl and imine groups.

Catalytic hydroborations

Our initial studies focused on reactions of 4-anisaldehyde using HBpin as the reducing agent as it is considerably less-reactive than its counterpart, HBcat.²⁷ Although no reaction was observed at room temperature over two hours, we have found that a variety of rhenium complexes are effective pre-catalysts for this reduction at elevated temperatures (Table 1 illustrates selected examples, where complexes 2 and 4 were chosen as representative examples of complexes of the type [Re(bipy)(CO)₃(L)]X, additional results are shown in the SI section). It should be noted, however, that under these conditions, depending upon the solvent, about 30-40% conversion to give the saturated boronate ester product is observed even in the absence of a rhenium pre-catalyst. The rhenium dimer [Re2(CO)10] (Entry 2) showed no significant improvements in conversion (as ascertained by ¹H NMR spectroscopy) in C₆D₆ but up to 90% of the starting aldehyde was reduced in reactions carried out in CD₂Cl₂, illustrating that the Re(0) species had some effectiveness in this solvent. More remarkable is that complete conversion was achieved in either solvent using the readily available rhenium(I) complex [Re(CO)₅Br] (Entry 3). Substitution of two of the carbonyl ligands with a more basic bipy group (Entry 4) in the neutral rhenium(I) complex showed reduced conversion in CD 2Cl2 and no improvements in reactions carried out in C₆D₆. We then examined a variety of complexes of the type [Re(N-N)(CO) 3(L)]X (SI) and are pleased to report that several of these low-valent species proceeded with complete conversions in reactions carried out in both solvents. Unfortunately, complexes containing either basic pyridine or quinoline (Entry 5) gave lower conversions as considerable degradation of the borane to B₂pin₃ and H₂ was observed using multinuclear NMR spectroscopy in these cases. The base or transition-metal promoted degradation of HBpin and other hydridoboranes is well-documented.¹ It is important to note that complexes containing both boron-containing WCAs [BAr^F₄]⁻ and [B(3,5-ditBu-cat)₂]⁻ gave excellent conversions. Finally, reactions employing rhenium complexes in higher oxidation states were also examined. It is not really surprising that the oxo-rhenium complex 1 proceeded with complete conversion (Entries 12) in both solvent systems as Fernandes and co-workers have previously demonstrated the utility of this species in catalysed hydroborations.²⁰

Encouraged by these promising results, we decided to expand the scope of these hydroborations and initially examine various aldehyde, ketone and aldimine derivatives (Table 1) using the

readily-available high-oxidation state complex [(PPh,3)2Re(Q)2] (1) as the representative catalyst precursor Read to how the set of the set o substrates were carried out using HBpin and 5 mol% precatalyst in C₆D₆ at 80°C for 2 h. Benzaldehyde, 4-4-nitrobenzaldehyde fluorobenzaldehvde and were investigated to see what effect varying the electronics on the aldehyde had on these reductions. Like 4-anisaldehyde, hydroborations of both benzaldehyde and 4fluorobenzaldehyde proceeded smoothly with complete conversion of the aldehyde, demonstrating that these reactions work well with electron-rich and electron-poor aromatic derivatives. Reactions with 4-nitrobenzaldehyde were of singular interest as both the aldehyde and the nitro group were reduced using excess HBpin (3 equiv) under these conditions. The catalytic reduction of aromatic nitro groups has been the subject of recent considerable interest. 19,28 Hydroborations of

Table 1 Rhenium-Catalyzed Hydroboration of 4-Anisaldehyde with Pinacolborane



Entry	Catalyst Precursor	Conversion ^a	
		C_6D_6	CD_2Cl_2
1	none	30	40
2	[Re ₂ (CO) ₁₀]	30	90
3	[Re(CO)₅Br]	100	100
4	[Re(bipy)(CO)₃Br] ^b	30	95
5	[Re(bipy)(CO)₃(quin)][BAr ^F 4] ^c	45	80
6	[Re(bipy)(CO) ₃ (CH ₃ CN)][BAr ^F ₄]	100	100
7	[Re(bipy)(CO) ₃ (THF)][BAr ^F ₄]	100	100
8	[Re(bipy)(CO) ₃ (CH ₃ CN)][B(3,5-di-tBu-cat) ₂] ^d	100	100
9	[(PMe₂Ph)₃ReCl₃]	30	85
10	[(PPh ₃) ₂ Re(O) ₂ I]	100	100

^aConversion ascertained using ¹H NMR spectroscopy averaged over three runs. Reactions were carried out in a sealed J. Young tube in a water bath heated at 80°C for two hours. ^bbipy = 2,2'-bipyridine. ^cquin = quinoline. ^d3,5-di-tBu-cat = bis-(3,5-di-tert-butyl[1,2-benzenediolato(2-)-O,O']borate

sterically-hindered mesitaldehyde (Entry x) also proved successful (100% conversion) as did the aliphatic derivative hexanal (Entry) . Reactions with thiophene-2-carbaldehyde (Entry) proceeded with complete conversion demonstrating that this reaction is tolerant of heteroaromatic groups. More remarkable, however, is that the rhenium-catalysed of *trans*-cinnamaldehyde hydroboration gave the corresponding 1,2-addition product predominantly with only minor amounts of the 1,4-addition product (< 2%) (Entry x). Boron-element additions with this challenging substrate frequently give mixtures of both 1,2- and 1,4-addition products.²⁸ Similar results were generally observed in reactions employing the low-valent rhenium complexes and variation of the bipy (N-N) groups and labile ligand L had negligible effect on both activities and selectivities (see SI).

Dalton Transactions

Published on 30 May 2017. Downloaded by State University of New York at Binghamton on 31/05/2017 11:35:17.

 $\label{eq:Table 2} {$ Table 2 $ Rhenium-Catalyzed Hydroboration of Aldehydes and Aldimines using 5 mol% [(PPh_3)_2Re(O)_2l] with Pinacolborane.^a $ } }$



Entry	Substrate	Product	Yield ^a
1	°_∓	Bpin	100
2	F F	F F Bpin	100
3	O ₂ N H	(pinB) ₂ N	100
4	С Н Н	Bpin	100
5	✓	O Bpin	100
6	С ^S Н	S Bpin	100
7	С Н	Bpin	98
8	Me	Me	40
9	Ph	O-Bpin Ph	15
10	H	pinB	100
11	NM H	pinB, NMe	100
12	NPh	pinB, NPh	25
13	N H	pinB N	100
14	H MB MB	Me Me	100

^aConversion ascertained using ¹H NMR spectroscopy averaged over three runs. Reactions were carried out in a sealed J. Young tube in a water bath heated at 80°C for two hours. We then turned our attention to the rhenium-catalysed hydroboration of ketones using 5 mol% 1 as this precataryst was the most efficient at reducing the more challenging substrates. Unfortunately, reactions of HBpin with acetophenone only gave 40% (by ¹H NMR spectroscopy) conversion after 2 h at 80 °C and even lower conversions (15%) were observed in reactions with benzophenone. Taking advantage of these reduced activities able to reduce 4-acetylbenzaldehyde we were chemoselectively, as hydroboration occurred solely at the aldehyde group. Although aldimines derived from primary aliphatic amines, such as N-benzylidenemethanamine, were readily reduced under these conditions, attempts to affect the hydroboration of aldimines derived from aromatic primary amines proved more challenging. Indeed, only 25% of the bulkier aldimine, N-benzylidene-4-methoxyaniline, was actually reduced under these conditions.

More significant is the observation that hydroborations of (E)-*N*-benzylideneprop-2-yn-1-amine proceeded with complete selectivity for the imine C=N bond as the alkyne group was left intact. Likewise, rhenium-catalysed reactions of (E)-1-(4-((prop-2-yn-1-ylimino)methyl)-phenyl)ethanone with HBpin were remarkably selective and only reduction of the imine double bond was observed. In the absence of any catalyst, only the aldehydes seemed to react with HBpin under these conditions but only gave the corresponding reduction products in around 10-40% yield.

Considering the wealth of recent interest into the catalysed hydroboration of pyridines using transition metals, ³⁰ lanthanides,³¹ main-group compounds,³² as well as metal-free methodologies,³³ we decided to examine the potential of using rhenium complexes to facilitate this remarkable reduction. Although no reaction was observed in the case of pyridine itself, 5 mol% of [(PPh₃)₂Re(O)₂I] can be employed along with 2 eq of HBpin at 80°C for 2 h to afford partially reduced quinoline products while preserving the annulated carbocyclic aromatic ring (Scheme 4). When the reaction was carried out in C_6D_6 , a mixture of the 1,2-addition product Q1, the 1,4-addition product Q2 as well as the partially-saturated derivative Q3 was observed in a ratio of 70:20:10 as ascertained by ¹H NMR spectroscopy. The borylated guinoline derivative Q3 presumably arises from a subsequent hydrogenation of **Q1** and Q2. Dihydrogen and B₂pin₃ are well-known decomposition products frequently observed in catalysed hydroboration reactions.³⁴ Reactions of quinoline using only one equivalent of HBpin did not improve selectivities and resulted in incomplete conversion. Changing the solvent to CD₂Cl₂ gave increased amounts of both Q2 and Q3 in a ratio of 15:55:30. Although other rhenium complexes were also examined (see SI section), the highest selectivities were achieved using [Re(CO) 5Br] in C6D6 which gave a mixture of the two addition products in a ratio of 90:10:0. Unfortunately, the low valent rhenium complexes all showed poor conversion of this substrate. These results might explain why hydroborations using precatalyst 2 gave lower conversions (Table 1, Entry 5) for all substrates as HBpin could also be reducing the labile quinoline ligand.



While it is likely that hydroborations using $[(PPh_3)_2Re(O)_2I]$ proceed via a mechanism similar to that outlined by Fernandes and co-workers,²⁰ where the B–H bond in HBpin is activated by a sigma-bond metathesis step to generate a reactive rhenium hydride intermediate, along with a [Re]-OBpin group, it is unclear yet how these reactions occur using [Re(CO) 5Br] and the low-valent rhenium cationic complexes. In analogous rheniumcatalysed hydrogenation and hydrosilylation reactions, 35 possible mechanisms have been postulated whereby the active catalytic species involves a low-valent rhenium hydride.³⁶ Indeed, monitoring all reactions upon completion of catalysis by ^{1}H NMR spectroscopy in reactions employing [Re(bipy)(CO)₃(THF)][BAr^F₄] and [Re(bipy)(CO)₃(CH₃CN)][B(3,5di-tBu-cat)₂] as pre-catalysts showed resonances at -9.1 and -8.8 ppm, respectively. Likewise, reactions using [Re(CO) 5Br] and a slight excess of HBpin (1.5 eq) showed a major hydride at -13.5 ppm in the ¹H NMR spectra. Further investigation is currently underway to elucidate the mechanism of these unique reactions, the results of which will be reported in due course.

Conclusions

Published on 30 May 2017. Downloaded by State University of New York at Binghamton on 31/05/2017 11:35:17.

In this study, we report the first examples for the rheniumcatalysed hydroboration of aldehydes, ketones and aldimines, including heteroaromatic quinoline. Reactions are remarkably chemoselective and tolerant of several functional groups. A wide array of rhenium complexes were efficient pre-catalysts for these hydroborations including the low-valent complexes of the formula [Re(bipy)(CO)₃(L)]X (L = labile ligand, and X = [BAr^F₄]⁻ and [B(3,5-di-tBu-cat)₂]⁻) prepared in this study. A new silver spiroboronate ester Ag[B(3,5-di-tBu-cat)₂](NCCH₃)₃ was prepared and characterized fully and used to make one of the new rhenium complexes. The possibility that low valent rhenium hydrides may play a part in some of these catalysed reactions is under further investigation.

Experimental

General conditions and methods

All reagents and solvents, unless otherwise noted, were purchased from commercial sources and used without further purification. Solvents were distilled over appropriate drying agents under dinitrogen: CH_2Cl_2 over CaH_2 , THF over Na/benzophenone, and hexane and toluene over freshly wired sodium. CD_2Cl_2 was purchased from Cambridge Isotope Laboratories, degassed by three freeze-pump-thaw cycles, kept in a J. Young tube with potassium carbonate and stored in a dark place over oven-activated 4Å molecular sieves. Pinacolborane

ARTICLE

(HBpin) was purified by vacuum distillation prior Artoe usee B2(3,5-di-tBu-cat)3,25 DOI: 10.10[Re(CO)]B71,37 Compounds [Re(bipy)(CO)₃Br],³⁷ [Re(bipy)(CO)₃(THF)][BAr^F₄],³⁸ [Re(bipy)-(CO)₃OTf],³⁸ $[Re(bipy)(CO)_3(CH_3CN)][BAr_4]^{38}$ [Re(phen)-(CO)₃Br],³⁹ [Re(phen)(CO)₃OTf],³⁹ [Re(bipy)(CO)₃(MeIm)][OTf],⁴⁰ and Na[BAr^F₄] (Ar^F = 3,5-bis(trifluoromethyl)phenyl),²⁴ were prepared following the reported procedures. Compounds [Re(4,4'-X₂bipy)(CO)₃Br], [Re(3,4,7,8-Me₄phen)(CO)₃Br] were prepared in similar way to that described for [Re(bipy)(CO) 3Br], and [Re(4,4'-X₂bipy)(CO)₃OTf] and [Re(3,4,7,8-Me₄phen)-(CO)₃OTf] to that for [Re(bipy)(CO)₃OTf].³⁸ Please see the SI section for a complete list of new complexes, prepared, characterized and examined in this study.

NMR spectra were recorded on Bruker AV 300, Bruker AV 400 (¹H: 400.54 MHz and ¹³C: 100.72 MHz), or JEOL JNM-GSX400 (¹H: 400 MHz; ¹¹B: 128 MHz; and ¹³C: 100 MHz) spectrometers. Chemical shifts (δ) are reported in ppm [relative to residual solvent peaks (¹H and ¹³C) or external BF₃·OEt₂ (¹¹B)]. Multiplicities are reported as singlet (s), doublet (d), triplet (t), quartet (q), quintet (quint), multiplet (m), overlapping (ov), and broad (br). Melting points were measured uncorrected with a Stuart SMP30 apparatus. IR solution spectra were obtained using a Perkin-Elmer Spectrum 100 FT-IR spectrometer using 0.2 mm CaF₂ cells or in the solid state using a Thermo Fisher Scientific Nicolet iS5 FT-IR spectrometer in ATR mode and are reported in cm⁻¹. All manipulations were performed under an atmosphere of dinitrogen using standard Schlenk techniques or in an MBraun LabMaster glovebox.

Preparation of compounds

[Re(bipy)(CO)₃(quin)][BAr^F₄] (1). Quinoline (14 μ L, 0.12 mmol) was added to a solution of [Re(CO)₃(bipy)(THF)][BAr^F₄] (160 mg, 0.12 mmol) in CH₂Cl₂ (10 mL). The reaction mixture was stirred for 5 hours and then concentrated to 5 mL. Addition of hexane (15 mL) caused the precipitation of a yellow solid which was washed with hexane $(2 \times 10 \text{ mL})$ and dried under vacuum. Compound 1 was isolated as a yellow microcrystalline solid in a 91% yield (151 mg). Slow diffusion of hexane (8 mL) into a concentrated solution of 1 in CH₂Cl₂ (4 mL) at -20°C yielded yellow crystals of 1 one of which was employed for X-ray diffraction. Yield: 89% (148 mg). Anal. Calcd. for C₅₄H₂₇BF₂₄N₃O₃Re (1418.79) (%): C 45.71, H 1.92, N 2.96. Found: C 45.05, H 1.82, N 2.93. IR (CH₂Cl₂, cm⁻¹): 2039, 1936 (v_{CO}). ¹H NMR (CD₂Cl₂, 298 K, 400 MHz): δ 9.38 (d, J = 5.6 Hz, 2H, H₆ bipy), 8.49 (m, 2H, quin), 8.30 (d, J = 8.4 Hz, 1H, quin), 8.13 (m, 4H, H₄ bipy), 8.09 (m, 2H, H₃ bipy), 7.89([m, 2H, quin), 7.79 (m, 2H, H₅ bipy {obscured by BAr^F₄ signal, can be seen in COSY and HSQC spectrum}), 7.66 (m, 1H, quin), 7.24 (dd J = 8.3, 5.3 Hz, 1H, quin). ¹³C{¹H} NMR (CD₂Cl₂, 298 K, 100 MHz): δ 195.2 (2 CO), 189.3 (CO), 155.6 (C₂ bipy), 153.9 (C₆ bipy), 153.8 (CH quin), 147.4 (QC quin), 141.3 (CH quin), 141.1 (C₄ bipy), 132.3, 129.9 (CH quin), 128.6 (C₅ bipy {obscured by BAr^F₄ signal, can be seen in HSQC and HMBC spectra}), 128.4, 126.8 (CH quin), 125.9 (QC quin), 124.1 (C₃ bipy), 121.6 (CH quin).

Published on 30 May 2017. Downloaded by State University of New York at Binghamton on 31/05/2017 11:35:17.

Dalton Transactions

The resonances due to the anion $[BAr^{F_4}]^-$ are: ¹H NMR (CD₂Cl₂, 298 K, 400 MHz): δ 7.77 (s, 8H, H₀), 7.52 (s, 4H, H_p). ¹³C{¹H} NMR (CD₂Cl₂, 298 K, 400 MHz): δ 161.7 (q, J_{CF} = 49.1 Hz, C_i), 134.8 (C₀), 129.0 (q, J_{CF} = 31.3 Hz, C_m), 125.1 (q, J_{CF} = 271.2 Hz, CF₃), 117.5 (C_p).

Ag[B(3,5-di-tBu-cat)₂](NCCH₃)₃ (3). B₂(3,5-di-tBu-cat)₃ (1.50 g, 2.20 mmol) was added in small portions as a solid to a stirred CH₃CN (20 mL) suspension of Ag(acac) (0.50 g, 2.42 mmol). The reaction mixture was stirred in the dark at RT for 18 hours. The reaction mixture was filtered to remove residual solids and the volume of solvent was reduced to 10 mL under vacuum and the clear red solution was stored at -30 °C. The initial precipitate [acacB(3,5-di-tBu-cat)] was collected by suction filtration and discarded. The amount of CH₃CN was reduced to 5 mL under vacuum and the solution stored once again at -30 °C. The second precipitate was collected by suction filtration to afford 9 as a white crystalline solid. Yield: 61% (0.75 g); mp: 112-113 °C (decomposition). Anal. Calcd. for C₃₄H₅₈AgBN₃O₄ (691.53) (%): C 59.84, H 7.24, N 6.16. Found: C 60.08, H 7.19, N 5.98. IR (ATR, cm⁻¹): 2961, 2271 (v_{CN}). ¹H NMR data (CD₃CN, 298 K, 400 MHz): δ 6.54 (d, J = 1.8 Hz, 2H, Ar), 6.53 (d, J = 1.8 Hz, 2H, Ar), 1.31 (s, 18H, tBu), 1.24 (s, 18H, tBu). ¹¹B NMR data (CD₃CN, 298 K, 128 MHz): δ 13.1 (sharp). ¹³C{¹H} NMR data (CD₃CN, 298 K, 100 MHz): δ 151.6 (C cat), 147.1 (C cat), 139.4 (C cat), 128.9 (C cat), 110.7 (CH cat), 104.0 (CH cat), 34.1 (C tBu), 33.7 (C tBu), 31.4 (CH₃ tBu), 28.9 (CH₃ tBu).

[Re(bipy)(CO)₃(CH₃CN)][B(3,5-di-tBu-cat)₂] (4). To a stirred CH₃CN (5 mL) yellow suspension of [Re(bipy)(CO)₃Br] (200 mg, 0.39 mmol) in a foil covered 25 mL RB was added a colourless CH₃CN (2 mL) solution of 3 (220 mg, 0.47 mmol). The reaction was allowed to proceed for 18 hours at RT at which point a grey solid (AgBr) was removed by filtration and discarded. The resulting orange filtrate was collected and the solvent was removed under vacuum to afford an oily orange solid. Trituration of the oily solid with hexanes (3 x 10 mL) afforded an orange-yellow solid which was collected by suction filtration and washed with hexanes (2 x 5 mL) to afford [Re(bipy)(CO)₃(CH₃CN)][B(3,5-di-tBu-cat)₂]. Yield: 88% (305 mg); mp: 189-191°C (decomposes to a brown solid). Anal. Calcd. for C43H51N3BO7Re (918.90) (%): C 56.20, H 5.59, N 4.57. Found: C 56.27, H 5.52, N 4.48. IR (ATR, cm⁻¹): 2951, 2036 (v_{co}), 1925 (v_{CO}). ¹H NMR (CD₂Cl₂, 298 K, 400 MHz): δ 8.86 (dd, J = 5.5, 1.4 Hz, 2H, Ar), 8.37 (d, J = 8.2 Hz, 2H, Ar), 7.78 (ov ddd, J = 8.2, 7.8, 1.4 Hz, 2H, Ar), 7.40 (ddd, J = 7.8, 5.5, 1.4 Hz, 2H, Ar), 6.54 (d, J = 1.4 Hz, 2H, Ar), 6.24 (d, J = 1.4 Hz, 2H, Ar), 1.83 (s, 3H, CH ₃CN), 1.27 (br s, 36H, *t*Bu). ¹¹B NMR (CD₂Cl₂, 298 K, 120 MHz): δ 13.1 (sharp). ¹³C{¹H} NMR (CD₂Cl₂, 298 K, 100 MHz): δ 193.8 (2 CO), 190.1 (CO), 156.2 (C bipy), 152.7 (CH bipy), 151.3 (C cat), 147.2 (C cat), 141.1 (CH bipy), 139.3 (C cat), 129.4 (C cat), 127.7 (CH bipy), 125.3 (CH bipy), 122.2 (C CH₃CN), 111.0 (CH cat), 104.2 (CH cat), 34.4 (C tBu), 33.9 (C tBu), 31.9 (CH₃ tBu), 29.3 (CH₃ tBu), 3.0 (CH₃ CH₃CN).

General rhenium catalysed hydroborations

To a stirred solution of substrate (0.18 mmol), and the appropriate rhenium complex (5 mol%) in $CC_{10} + CD_{2} + CD_$

X-ray crystallography

Data collection was performed on a Bruker X8 KappaAPEXII single crystal diffractometer, using Cu-Kα radiation (I = 1.54184 Å). Images were collected at a 65 mm fixed crystal-detector distance, using the oscillation method, with 1° oscillation and variable exposure time per image (4-16 s). Data collection strategy was calculated with the program CrysAlisPro CCD.⁴¹ Data reduction and cell refinement was performed with the program CrysAlisPro RED.⁴² An empirical absorption correction was applied using the SCALE3 ABSPACK algorithm as implemented in the program CrysAlisPro RED⁴¹ Crystal structure was solved by direct methods, using the program SIR-92.42 Anisotropic least-squares refinement was carried out with SHELXL-97.43 All non-hydrogen atoms were anisotropically refined. Hydrogen atoms were geometrically placed riding on their parent atoms with isotropic displacement parameters set to 1.2 times the Ueg of the atoms to which they are attached (1.5 for methyl groups). Crystallographic data have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication numbers 1544620 and 1544621.

Acknowledgements

The Oviedo authors thank MINECO and FEDER (grant CTQ2015-70231-P) and Principado de Asturias (grant FC-15-GRUPIN14-103) for funding, and Ministerio de Educación for an FPU predoctoral fellowship (to RA). SAW is grateful to the Natural Sciences and Engineering Research Council, the CRC programme and Mount Allison University for funding and Mr. Dee Durant for expert technical assistance and anonymous reviewers are thanked for their very helpful comments.

Notes and references

- (a) C. M. Crudden and D. Edwards, *Eur. J. Org. Chem.*, 2003, 4695–4712.
 (b) C. M. Vogels and S. A. Westcott, *Curr. Org. Chem.*, 2005, **9**, 687–699.
- (a) I. P. Query, P. A. Squier, E. M. Larson, N. A. Isley and T. B. Clark, *J. Org. Chem.*, 2011, **76**, 6452–6456. (b) C. C. Chong and R. Kinjo, *ACS Catal.*, 2015, **5**, 3238–3259.
- 3 N. Eedugurala, Z. Wang, U. Chaudhary, N. Nelson, K. Kandel, T. Kobayashi, I. I. Slowing, M. Pruski and A. D. Sadow, ACS Catal., 2015, 5, 7399–7414.
- 4 (a) C. W. Lindsley and M. DiMare, *Tetrahedron Lett.*, 1994, 35, 5141–5144. (b) F. Almqvist, L. Torstensson, A. Gudmundsson and T. Frejd, *Angew. Chem. Int. Ed. Engl.*, 1997, 36, 376–377. (c) I. Sarvary, F. Almqvist and T. Frejd, *Chem. Eur. J.*, 2001, 7, 2158–2166. (d) A. A. Oluyadi, S. Ma and C. N. Muhoro, *Organometallics*, 2013, 32, 70–78.

Dalton Transactions

- 5 G. Zhang, H. Zeng, J. Wu, Z. Yin, S. Zheng and J. C. Fettinger, Angew. Chem. Int. Ed., 2016, 55, 14369–14372.
- 6 S. Chakraborty, P. Bhattacharya, H. Dai and H. Guan, Acc. Chem. Res., 2015, 48, 1995–2003.
- 7 (a) J. Guo, J. Chen and Z. Lu, *Chem. Commun.*, 2015, **51**, 5725–5727. (b) T. Zhang, K. Manna and W. Lin, *J. Am. Chem. Soc.*, 2016, **138**, 3241–3249.
- 8 A. E. King, S. C. E. Stieber, N. J. Henson, S. A. Kozimor, B. L. Scott, N. C. Smythe, N. C. Sutton and J. C. Gordon, *Eur. J. Inorg. Chem.*, 2016, **11**, 1635–1640.
- 9 S. Bagherzadeh and N. P. Mankad, *Chem. Commun.*, 2016, **52**, 3844–3846.
- 10 (a) M. Bandini, P. G. Cozzi, M. de Angelis and A. Umani-Ronchi, *Tetrahedron Lett.*, 2000, 41, 1601–1605. (b) S.-G. Roh, Y.-C. Park, D.-K. Park, T.-J. Kim and J.-H. Jeong, *Polyhedron*, 2001,
 20, 1961–1965. (c) M. Locatelli and P. G. Cozzi, *Angew. Chem. Int. Ed.*, 2003, 42, 4928–4930. (d) S.-G. Roh, J.-U. Yoon and J.-H. Jeong, *Polyhedron*, 2004, 23, 2063–2067. (e) P. A. Lummis, M. R. Momeni, M. W. Lui, R. McDonald, M. J. Ferguson, M. Miskolzie, A. Brown and E. Rivard, *Angew. Chem. Int. Ed.*, 2015, 53, 9347–9351.
- 11 A. Y. Khalimon, P. Farha, L. G. Kuzmina and G. I. Nikonov, *Chem. Commun.*, 2012, **48**, 455–457.
- (a) L. Koren-Selfridge, H. N. Londino, J. K. Vellucci, B. J. Simmons, C. P. Casey and T. B. Clark, *Organometallics*, 2009, 28, 2085–2090. (b) A. Kaithal, B. Chatterjee and C. Gunanathan, *Org. Lett.*, 2015, 17, 4790–4793.
- 13 M. W. Drover, L. L. Schafer and J. A. Love, Angew. Chem. Int. Ed., 2016, 55, 3181–3186.
- 14 T. J. Hadlington, M. Hermann, G. Frenking and C. Jones, J. Am. Chem. Soc., 2014, **136**, 3028–3031.
- 15 C. C. Chong, H. Hirao and R. Kinjo, Angew. Chem. Int. Ed., 2015, 54, 190–194.
- 16 (a) Z. Yang, M. Zhong, X. Ma, S. De, C. Anusha, P. Parameswaran and H. W. Roesky, *Angew. Chem. Int. Ed.*, 2015, 54, 10225–10229. (b) V. K. Jakhar, M. K. Barman and S. Nembenna, *Org. Lett.*, 2016, 18, 4710–4713.
- 17 A. J. Blake, A. Cunningham, A. Ford, S. J. Teat and S. Woodward, *Chem. Eur. J.*, 2000, 6, 3586–3594.
- 18 (a) M. Arrowsmith, T. J. Hadlington, M. S. Hill and G. Kociok-Köhn, *Chem. Commun.*, 2012, **48**, 4567–4569. (b) D. Mukherjee, A. Ellern and A. D. Sadow, *Chem. Sci.*, 2014, **5**, 959–964.
- 19 V. L. Weidner, C. J. Barger, M. Delferro, T. L. Lohr and T. J. Marks, *ACS Catal.*, 2017, **7**, 1244–1247.
- 20 (a) A. C. Fernandes, J. A. Fernandes, F. A. A. Paz and C. C. Romão, *Dalton Trans.*, 2008, 6686–6688. (b) I. Cabrita, S. C. A. Sousa and A. C. Fernandes, *Tetrahedron Lett.*, 2010, **51**, 6132–6135. (c) A. C. Fernandes, J. A. Fernandes, C. C. Ramão, L. F. Veiros and M. J. Calhorda, *Organometallics*, 2010, **29**, 5517–5525. (d) S. C. A. Sousa, I. Cabrita and A. C. Fernandes, *Chem. Soc. Rev.*, 2012, **41**, 5641–5653.
- 21 (a) M. E. Viguri, M. A. Huertos, J. Pérez, L. Riera and I. Ara, J. Am. Chem. Soc., 2012, 134, 20326–20329. (b) M. A. Huertos, J. Pérez and L. Riera, Chem. Eur. J., 2012, 18, 9530–9533. (c) M. E. Viguri, M. A. Huertos, J. Pérez, L. Riera and I. Ara, Chem. Eur. J., 2013, 19, 12974–12977. (d) R. Arévalo, J. Pérez and L. Riera, Inorg. Chem., 2013, 52, 6785–6787. (e) M. E. Viguri, J. Pérez and L. Riera, Chem. Eur. J., 2014, 20, 5732–5740. (f) H. Martínez-García, D. Morales, J. Pérez, M. Puerto and I. del Río, Chem. Eur. J., 2014, 20, 5821–5834. (g) R. Arévalo, J. Pérez and L. Riera, Chem. Eur. J., 2015, 21, 3546–3549.
- 22 E. N. Daley, C. M. Vogels, S. J. Geier, A. Decken, S. Doherty and S. A. Westcott, *Angew. Chem. Int. Ed.*, 2015, **54**, 2121–2125.
- 23 (a) J. J. Kennedy-Smith, K. A. Nolin, H. P. Gunterman and F. D. Toste, J. Am. Chem. Soc., 2003, 125, 4056–4057. (b) W. Huang and H. Berker, Chimia, 2005, 59, 113–115. (c) B. Royo and C. C. Romão, J. Mol. Catal. A: Chem., 2005, 236, 107–112. (d) G.

Du and M. M. Abu-Omar, *Organometallics*, 2006, **25**, 4920– 4923. (e) W.-G. Zhao and R. Hua, *Eur D* (*Qrg*) *Greatine Orline*, 5495–5498. (f) K. A. Nolin, J. R. Krumper, M. D. Pluth, R. G. Bergman and F. D. Toste, *J. Am. Chem. Soc.*, 2007, **129**, 14684– 14696. (g) H. Dong and H. Berke, *Adv. Synth. Catal.*, 2009, **351**, 1783–1788. (h) K. A. Nolin, R. W. Ahn, Y. Kobayashi, J. J. Kennedy-Smith and F. D. Toste, *Chem. Eur. J.*, 2010, **16**, 9555– 9562. (i) J. L. Smeltz, P. D. Boyle and E. A. Ison, *Organometallics*, 2012, **31**, 5994–5997. (j) P. Gu, W. Wang, Y. Wang and H. Wei, *Organometallics*, 2013, **32**, 47–51. (k) L. Huang, Y. Zhang and H. Wei, *Eur. J. Inorg. Chem.*, 2014, 5714– 5723. (l) H. Dong, Y. Jiang and H. Berke, *J. Organomet. Chem.*, 2014, **750**, 17–22.

- 24 M. Brookhart, B. Grant and A. F. Volpe Jr., Organometallics, 1992, 11, 3920–3922.
- 25 N. R. Halcovitch, C. M. Vogels, A. Decken and S. A. Westcott, *Can. J. Chem.*, 2009, **87**, 139–145.
- 26 (a) G. L. Ning, M. Munakata, L. P. Wu, M. Maekawa, Y. Suenaga, T. Kurodo-Sawa and K. Sugimoto, *Inorg. Chem.*, 1999, **38**, 5668–5673. (b) J. M. López-de-Luzuriaga, M. Monge, M. E. Olmos and D. Pascual, *Organometallics*, 2015, **34**, 3029–3038.
- 27 C. E. Tucker, J. Davidson and P. Knochel, J. Org. Chem., 1995, 57, 3482–3485.
- 28 (a) C. M. Vogels, A. Decken and S. A. Westcott, *Tetrahedron Lett.*, 2006, 47, 2419–2422. (b) S. M. Ansar and C. L. Kitchens, *ACS Catal.*, 2016, 6, 5553–5560. (c) K. Yang, F. Zhou, Z. Kuang, G. Gao, T. G. Driver and Q. Song, *Org. Lett.*, 2016, 18, 4088–4091.
- (a) E. C. Neeve, S. J. Geier, I. A. I. Mkhalid, S. A. Westcott and T. B. Marder, *Chem. Rev.*, 2016, **116**, 9091–9161. (b) A. B. Cuenca, R. Shishido, H. Ito and E. Fernández, *Chem. Soc. Rev.*, 2017, **46**, 415–430.
- 30 (a) K. Oshima, T. Ohmura and M. Suginome, J. Am. Chem. Soc.,
 2012, 134, 3699–3902. (b) A. Kaithal, B. Chatterjee and C. Gunanathan, Org. Lett., 2016, 18, 3402–3405.
- 31 A. S. Dudnik, V. L. Weidner, A. Motta, M. Delferro and T. J. Marks, *Nature Chem.*, 2014, 6, 1100–1107.
- 32 (a) M. Arrowsmith, M. S. Hill, T. Hadlington and G. Kociok-Köhn, Organometallics, 2011, **30**, 5556–5559. (b) J. Intemann, M. Lutz and S. Harder, Organometallics, 2014, **33**, 5722–5729.
- 33 (a) P. Eisenberger, A. M. Bailey and C. M. Crudden, J. Am. Chem. Soc., 2012, 134, 17384–17387. (b) K. Oshima, T. Ohmura and M. Suginome, Chem. Commun., 2012, 48, 8571– 8573 (c) X. Fan, J. Zheng, Z. H. Li and H. Wang, J. Am. Chem. Soc., 2015, 137, 4916–4919.
- 34 E. C. Keske, B. D. Moore, O. V. Kenkina, R. Wang, G. Schatte and C. M. Crudden, *Chem. Commun.*, 2014, **50**, 9883–9886.
- (a) E. A. Ison, R. A. Corbin and M. M. Abu-Omar, J. Am. Chem. Soc., 2005, 127, 11938–11939. (b) G. Alcaraz and S. Sabo-Etienne, Coord. Chem. Rev., 2008, 252, 2395–2409. (c) H. Dong and H. Berke, Adv. Synth. Catal., 2009, 351, 1783–1788. (d) Y. Jiang, J. Hess, T. Fox and H. Berker, J. Am. Chem. Soc., 2010, 132, 18233–18247. (e) Y. Jiang, O. Blacque, T. Fox and H. Berke, J. Am. Chem. Soc., 2013, 135, 7751–7760.
- 36 (a) D. Guisto, *Inorg. Chim. Acta Rev.*, 1972, **6**, 91–101. (b) M. A. Andrew, S. W. Kirtley and H. D. Kaesz, *Inorg. Chem.*, 1977, **16**, 1556–1561. (c) G. Jia, A. J. Lough and R. H. Morris, *J. Organomet. Chem.*, 1993, **461**, 147–156. (d) M. Bergamo, T. Beringhelli, G. D'Alfonso, G. Ciani, M. Moret and A. Sironi, *Organometallics*, 1996, **15**, 3876–3884. (e) Y. Ruiz-Morales, G. Schreckenbach and T. Ziegler, *Organometallics*, 1996, **15**, 3920–3923. (f) G. Albertin, S. Antoniutti, S. García-Fontán, R. Carballo and F. Padoan, *J. Chem. Soc., Dalton Trans.*, 1998, 2071–2081. (g) S. Ghosh, X. Lei, C. L. Cahill and T. P. Fehlner, *Angew. Chem. Int. Ed.*, 2000, **39**, 2900–2902. (h) N. Masciocchi, G. D'Alfonso, L. Garavaglia and A. Sironi, *Angew. Chem. Int. Ed.*, 2000, **39**, 4478–4480. (i) I. del Rosal, M. R.

Dalton Transactions Accepted Manuscript

Dalton Transactions

View Article Online DOI: 10.1039/C7DT01549F

Poteau and F. Jolibois, *Dalton Trans.*, 2008, 3959–3970. (j) S. Bolaño, J. Bravo, J. Castro, M. del Carmen Marín and S. García-Fontán, *Inorg. Chem. Commun.*, 2009, **12**, 916–918. (k) S. Bolaño, J. Bravo, F. Fernández-García, S. García-Fontán and M. del Carmen Marín, *J. Organomet. Chem.*, 2009, **694**, 2994–2999. (l) N. Li, Y. Xie, R. B. King and H. F. Schaeffer III, *J. Phys. Chem. A*, 2010, **114**, 11670–11680. (m) N. Li, Y. Xie, R. B. King and H. F. Schaeffer III, *Eur. J. Inorg. Chem.*, 2011, 4626–4636.

- 37 S. P. Schmidt, W. C. Trogler and F. Basolo, *Inorg. Synth.*, 1990, 28, 160–165.
- 38 E. Hevia, J. Pérez, V. Riera, D. Miguel, S. Kassel and A. Rheingold, *Inorg. Chem.*, 2002, **41**, 4673–4679.
- 39 E. Hevia, J. Pérez, L. Riera, V. Riera, I. del Río, S. García-Granda and D. Miguel, *Chem. Eur. J.*, 2002, 8, 4510–4521.
- 40 M. A. Huertos, J. Pérez and L. Riera, *J. Am. Chem. Soc.*, 2008, **130**, 5662–5663.
- 41 Oxford Diffraction (2008). CrysAlisProCCD, CrysAlisProRED. Oxford Diffraction Ltd., Abingdon, Oxfordshire, UK.
- 42 A. Altomare, G. L. Cascarano, C. Giacovazzo, A. Guagliardi, M. C. Burla, G. Polidori, M. Camalli, J. Appl. Cryst., 1994, 27, 435.
- 43 G. M. Sheldrick, SHELXL97, University of Göttingen, Germany, 1997.

The first examples of the catalysed hydroboration of aldehydes and aldimines using low- and high-valent rhenium complexes are reported.

