

Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: M. Pawliczek, T. Hashimoto and K. Maruoka, *Chem. Sci.*, 2017, DOI: 10.1039/C7SC04854H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Alkylative Kinetic Resolution of Vicinal Diols under Phase-Transfer Conditions: A Chiral Ammonium Borinate Catalysis

Martin Pawliczek, Takuya Hashimoto* and Keiji Maruoka*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Herein, we report the first alkylative kinetic resolution of vicinal alcohols realized by cooperative use of a chiral quaternary ammonium salt and an achiral borinic acid. In addition, a catalytic regioselective alkylation of a secondary alcohol in the presence of an unprotected primary one is presented, emphasizing the unique selectivity and potential of this ammonium borinate catalysis.

Introduction

Chiral phase-transfer catalysis has been extensively studied in the past three decades due to its operational simplicity and benign reaction conditions. A large body of work focusing on the development of new chiral catalysts and reactions, was reported during this period.¹ Despite these continuing progresses, one simple reaction, namely the asymmetric functionalization of unprotected alcohols, has remained intractable in chiral phase-transfer catalysis as a way to provide enantioenriched molecules. It is in contrast with the fact that the functionalization of alcohols has been known to be accelerated in the presence of quaternary ammonium salts long before the emergence of chiral phase-transfer catalysis.² From our experience in this field,^{1a,b} we surmised that an obstacle to achieve this goal is the formation of a localized, hard alkoxide anion. Compared to a delocalized and flat anion like an enolate,³ an alkoxide is structurally more flexible, rendering the recognition by a chiral cation difficult. The strong basicity of an alkoxide is also a problem since such a strong base is known to degrade the phase-transfer catalyst and is incompatible with base-sensitive substrates.

Taking these facts into consideration, we planned to realize an enantioselective functionalization of alcohols by merging the chiral ammonium salt with an additional catalyst which is able to interact directly with an alcohol and modify its properties. In this regard, Taylor *et al.* recently reported the use of achiral borinic acids for the mono-functionalization of vicinal diols via an intermediary formed borinate (Fig. 1).^{4,5} In such an anionic species, the alkoxide is structurally fixed and its basicity is attenuated. Deploying a chiral quaternary ammonium salt to pair up with this anionic species is expected to create a chiral

environment around the diol, facilitating asymmetric functionalization of the diol (Figure 1).⁶

We report herein our primary investigations on this subject using an alkylative kinetic resolution of vicinal diols. Despite the fact that alkylation of alcohols directly forges ubiquitous ether moieties,⁷ such transformation has remained unexplored in asymmetric catalysis.^{8,9} High enantioselectivities were achieved in the kinetic resolution of secondary as well as tertiary diols by exploiting this chiral ammonium borinate catalysis. The availability of the optimal chiral quaternary ammonium salt, a persistent bottleneck in the binaphthyl-based chiral phase-transfer catalysts,^{1a,b} was enhanced by applying a metal-catalyzed C-H arylation. To further highlight the unique reactivity and selectivity of this catalysis, we also demonstrate a catalyst-controlled, regioselective alkylation of an innately less reactive secondary alcohol of a terminal vicinal diol.




Fig. 1 Cooperative catalysis for the asymmetric functionalization of vicinal diols.

Results and discussion

The first challenge to achieve our goal was the confirmation of reaction conditions to facilitate the alkylation only when both the chiral quaternary ammonium salt and the borinic acid are present. The optimization was implemented by use of (*rac*)-1-phenyl-1,2-ethanediol **2a** and benzyl bromide as reactant in combination with 2 mol% of a simple chiral ammonium salt **4** and 10 mol% of a nitrogen-tethered borinic acid **8** as catalyst pair (Table 1).¹⁰ Initially we found out that a biphasic system


* Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo, Kyoto, 606-8502, (Japan). E-mail: takuya@kuchem.kyoto-u.ac.jp, maruoka@kuchem.kyoto-u.ac.jp

† Electronic Supplementary Information (ESI) available: See DOI: 10.1039/x0xx00000x



ARTICLE

consisting of CH_2Cl_2 and aqueous base suppresses independent catalyst activities and ensures enantioselective product formation through a cooperative catalytic pathway (entries 1–3). Screening of a collection of chiral ammonium salts (see Supplementary Information) revealed the effectiveness of catalyst **6** (entries 4 and 5),¹¹ which was further optimized by attaching 3,5-bis(heptafluoroisopropyl)-phenyl substituents (**7**) (entries 6 and 7).¹² A final optimization resulted in the use of 4 mol% of **7**, 5 mol% of **8**, a CH_2Cl_2 -toluene co-organic solvent system, an excess amount of benzyl bromide, 2 equivalent of Cs_2CO_3 and NaI as additive. Under these conditions, product **2aa** was obtained in 44% yield with 91% ee (entry 8). Interestingly, a considerable amount of side-product **3aa**, derived from alkylation of the secondary alcohol, was also observed with the preferential uptake of (*R*)-**1a**, indicating the divergent nature of this catalysis.

Table 1 Optimization of the reaction conditions.^{a-c}


Chemical structures for Table 1:


- Ar = 3,4,5-F₃C₆H₂ (4)**
- Ar = 3,4,5-F₃C₆H₂ (5)**
- Ar = 3,4,5-F₃C₆H₂ (6)**
- Ar = 3,5-(iC₃F₇)₂C₆H₃ (7)**
- 8**

entry	PTC	2aa		1a		3aa	
		(% yield, % ee)		(% yield, % ee)		(% yield, % ee)	
1	4	17	21	72	4	-	-
2 ^d	4	-	-	97	0	-	-
3	-	-	-	97	0	-	-
4	5	14	3	78	1	-	-
5	6	15	45	75	8	-	-
6 ^e	6	4	54	90	2	-	-
7 ^e	7	7	87	86	4	-	-
8 ^{f,g}	7	44	91	49	74	6	22

^a Performed with (*rac*)-**1a** (0.10 mmol), benzyl bromide (0.06 mmol), phase-transfer catalyst (2 mol%) and **8** (10 mol%) in $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ (0.35 mL/0.65 mL) at rt. ^b NMR yield. ^c Ee determined by chiral HPLC. ^d Performed without **8**. ^e Performed at 0 °C for 48 h. ^f **7** (4 mol%), **8** (5 mol%), Cs_2CO_3 (2 eq.) and NaI (5 eq.) in CH_2Cl_2 /toluene/ H_2O (0.15 mL/0.20 mL/0.65 mL) at 0 °C for 96 h. ^g Isolated yield.


At this stage, we decided to streamline the synthesis of the structurally complex phase-transfer catalyst **7** with the help of the state-of-the-art C-H arylation.¹³ Starting from readily available 1,1'-binaphthyl-2,2'-dicarboxylic acid,¹⁴ direct mono

C-H arylation of the 3-position was investigated using several methods. It was revealed that the ruthenium-catalyzed arylation of mono ester **9** with 3,5-(iC₃F₇)₂C₆H₃ is a robust and scalable procedure (Scheme 1).^{15,16} Arylated compound **10** was further transformed into dibromide **11** by conventional synthetic methods, which upon treatment with aqueous ammonia yielded the desired catalyst **7**.

**Scheme 1** Preparation of catalyst **7**. Conditions: a) [(*p*-cym)RuCl₂]₂, Bu₃P, 3,5-(iC₃F₇)₂C₆H₃, K₂CO₃, NMP, 110 °C, 78% yield; b) LiOH aq., THF, reflux; c) BH₃·Me₂S, THF, reflux, 86% yield (2 steps); d) PBr₃, THF, rt, 88% yield; e) NH₃ aq., CH₃CN, 80 °C, 69% yield.

With the optimized reaction conditions and the secure supply of the catalyst in hand, we investigated the substrate scope with various alkyl halides (Table 2, entries 1–13). As for benzylic bromides, regardless of the substituent pattern on the aromatic ring, the alkylation proceeded with good enantioselectivities at nearly half conversion of the diol (**2ab–2aj**). The reaction could be scaled up to 1.0 mmol without difficulty as shown in the preparation of **2ad** (entry 3). Efficient kinetic resolution took place with prenyl and cinnamyl bromides to afford products **2ak** and **2al** with high enantioselectivities (entries 10 and 11). The mild reaction conditions allowed the use of base-sensitive methyl 4-bromocrotonate, to give **2am** bearing an unsaturated ester (entry 12). The reaction using phenylpropargyl bromide was found to be slow, giving **2an** in 36% yield with 90% ee after 6 d (entry 13). In these experiments, we isolated some amount of side-products derived from the alkylation of the secondary alcohol (e.g. **3aa** in Table 1) in a range from 10% to 30% yield in addition to the recovered (*R*)-diols **1** (see SI for detail).

We then turned our attention to the use of various terminal vicinal diols as nucleophile (Table 2, entries 14–22). A variety of aromatic diols were found to be applicable irrespective of the substituent pattern and the electronic property (**2ba–2ia**). The use of an aliphatic diol gave the benzylated product **2ja** with lower enantioselectivity, implying the necessity of catalyst re-optimizations for this class of substrates.

Table 2 Asymmetric alkylation of terminal vicinal diols.^{a-c}


entry	PTC	2		1		3	
		(% yield, % ee)		(% yield, % ee)		(% yield, % ee)	
1	4	17	21	72	4	-	-
2 ^d	4	-	-	97	0	-	-
3	-	-	-	97	0	-	-
4	5	14	3	78	1	-	-
5	6	15	45	75	8	-	-
6 ^e	6	4	54	90	2	-	-
7 ^e	7	7	87	86	4	-	-
8 ^{f,g}	7	44	91	49	74	6	22




entry	R	R'	% yield	% ee	
1 ^d	Ph	2-MeC ₆ H ₄	51	87	2ab
2	Ph	3-MeC ₆ H ₄	46	82	2ac
3 ^{d,e}	Ph	4-MeC ₆ H ₄	48(48)	87(86)	2ad
4	Ph	2-FC ₆ H ₄	47	90	2ae
5	Ph	2-BrC ₆ H ₄	45	85	2af
6	Ph	2-CNC ₆ H ₄	49	87	2ag
7	Ph	4-PhC ₆ H ₄	46	91	2ah
8	Ph	3,5-(CF ₃) ₂ C ₆ H ₄	36	89	2ai
9	Ph	2-naphthyl	36	91	2aj
10	Ph	(Me) ₂ C=CH	40	89	2ak
11	Ph	(E)-PhCH=CH	39	84	2al
12	Ph	(E)-MeOC ₆ H ₄ =CH	39	80	2am
13	Ph	PhC≡C	36	90	2an
14	2-MeC ₆ H ₄	Ph	45	83	2ba
15	3-MeC ₆ H ₄	Ph	40	87	2ca
16	4-MeC ₆ H ₄	Ph	34	97	2da
17	2-FC ₆ H ₄	Ph	42	87	2ea
18	3-FC ₆ H ₄	Ph	40	87	2fa
19	4-FC ₆ H ₄	Ph	44	91	2ga
20	4-BrC ₆ H ₄	Ph	33	82	2ha
21	4-MeOC ₆ H ₄	Ph	46	88	2ia
22	Cy	Ph	38	48	2ja

^a Performed with **7** (4 mol%), **8** (5 mol%), Cs₂CO₃ (2 eq.) and NaI (5 eq.). ^b Isolated yield. ^c Ee determined by chiral HPLC. ^d Performed with 2 mol% of **7**. ^e Yield and ee in parentheses are the results of 1 mmol scale experiment.

As more challenging substrates, terminal vicinal diols (*rac*)-**12** with a tertiary alcohol moiety were examined under our reaction conditions (Table 3).^{9b} Gratifyingly, exactly the same co-catalytic system worked for substrates having an additional methyl group (entries 1–8). At nearly half conversion, benzylated products **13a–13h** were obtained with enantioselectivities ranging around 80% to 90%. Attachment of other alkyl substituents instead of the methyl group led to lower enantioselectivity as exemplified by **13i** (entry 9). A general requirement for tertiary alcohols is a higher borinic acid loading and longer reaction time due to the slow turnover of the catalysis.

Table 3 Alkylation of tertiary vicinal diols.^{a,c}



entry	R	R''	% yield	% ee	
1	Ph	Me	47	84	13a
2	2-MeC ₆ H ₄	Me	42	82	13b
3	3-MeC ₆ H ₄	Me	38	85	13c
4 ^d	4-MeC ₆ H ₄	Me	41	83	13d
5	4-FC ₆ H ₄	Me	30	91	13e
6	4-MeOC ₆ H ₄	Me	46	87	13f
7	2-naphthyl	Me	33	78	13g
8	3-thienyl	Me	44	72	13h
9	Ph	CH ₂ CH=CH ₂	25	69	13i

^a Performed with **7** (4 mol%), **8** (10 mol%), Cs₂CO₃ (2 eq.) and NaI (5 eq.). ^b Isolated yield. ^c Ee determined by chiral HPLC. ^d Performed with 15 mol% of **8**.

We then carried out NMR experiments of the ammonium borinate prepared from 1:1:2 ratio of **7**, **8** and (*rac*)-**1a**, and Cs₂CO₃ in CDCl₃ (see SI for detail). The analysis of the ¹H and ¹¹B NMRs confirmed the formation of two diastereomeric ammonium borinates favoring the ion pair **IP_S** complexed with (*S*)-**1a** over **IP_R** (Fig. 2). Addition of more of (*rac*)-**1a** to this solution shifted the ratio of ion pairs within minutes, indicating that these two diastereomeric pairs are in a rapid equilibrium relative to the reaction time frame. This observation suggested that the alkylation, not the ion pairing, is the enantio-determining step. Therefore, the enantioselectivity arguably stems from the reactivity difference of **IP_S** and **IP_R** for alkylation (path *a* > path *c*). As a result of deceleration of the reaction via path *c*, **IP_R** is considered to react eventually at the innately less reactive secondary alcohol to give **3aa**.




Fig. 2 The reaction pathways of the ammonium borinate catalyzed alkylation.


Intrigued by the unique regioselectivity toward secondary alcohol observed with **IP_R**, we decided to take a closer look on this phenomenon (Scheme 2a).^{6b,17,18} Such a transformation is in contrast to the studies conducted by Taylor *et al.*, in which the borinic acid exclusively functionalized the primary alcohol of a terminal diol.^{4b} It was thus surmised that the judicious choice of an ammonium salt would reveal a novel, catalytic regioselective alkylation of the secondary alcohol of terminal vicinal diols. By analyzing the results of the previous catalyst screening (Table 1 and SI), we could quickly identify the optimal catalyst **14** for this purpose. Starting from commercially available (*R*)-**1a** with 87% ee, the catalyst controlled, regioselective alkylation proceeded with a ratio of 3.4:1 in favor of the secondary alcohol. A slight increase in enantiomeric excess for the desired product **3aa** was observed as well.

Finally, we applied our catalyst system to the desymmetrization of a meso-diol (Scheme 2b). Treatment of meso-diol **15** with 2-methylbenzyl bromide gave the desired compound **16** in 96% yield with 52% ee paving the way for further optimizations in the future.



ARTICLE

Journal Name



Scheme 2. a) Catalyst-controlled, regioselective alkylation of the secondary alcohol of (R)-1a. b) Desymmetrization of meso-diol 15.

Conclusions

In conclusion, we developed herein the cooperative catalytic system capable of performing asymmetric functionalization of alcohols under phase-transfer conditions as demonstrated by the first alkylative kinetic resolution of secondary and tertiary diols. In addition, this method allowed a catalyst-controlled, regioselective alkylation of the less accessible secondary alcohol in the presence of an unprotected primary one. Our ongoing research is focused on the application of this chiral ammonium borinate catalysis to other electrophiles and more complex diols like saccharides.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements


M.P. thanks Deutsche Forschungsgemeinschaft (DFG) for postdoctoral fellowship. K.M. and T.H. thank JSPS KAKENHI Grant Number JP26220803. T.H. thanks JSPS KAKENHI Grant Number JP 16H01021 in Precisely Designed Catalysts with Customized Scaffolding.

Notes and references

- (a) T. Hashimoto and K. Maruoka, *Chem. Rev.*, 2007, **107**, 5656; (b) S. Shirakawa and K. Maruoka, *Angew. Chem., Int. Ed.*, 2013, **52**, 4312; (c) J. Tan and N. Yasuda, *Org. Process Res. Dev.*, 2015, **19**, 1731.
- (a) H. H. Freedman and R. A. Dubois, *Tetrahedron Lett.*, 1975, **16**, 3251; (b) J. de la Zerda, G. Barak and Y. Sasson, *Tetrahedron*, 1989, **45**, 1533; (c) Y. Liu, L. Ciszewski, L. Shen and M. Prasad, *Org. Process Res. Dev.*, 2014, **18**, 1142.
- For chiral phase-transfer catalyzed O-alkylation of enolates, see; J. D. Jolliffe, R. J. Armstrong, and M. D. Smith, *Nat. Chem.* 2017, **9**, 558.

- (a) L. Chan and M. S. Taylor, *Org. Lett.*, 2011, **13**, 3090; (b) D. Lee, C. L. Williamson, L. Chan and M. S. Taylor, *J. Am. Chem. Soc.*, 2012, **134**, 8260; (c) E. Dimitrijevic and M. S. Taylor, *Chem. Sci.*, 2013, **4**, 3298; (d) K. A. D'Angelo and M. S. Taylor, *J. Am. Chem. Soc.*, 2016, **138**, 11058; (e) K. A. D'Angelo and M. S. Taylor, *Chem. Commun.*, 2017, **53**, 5978.
- S. O. Bajaj, E. U. Sharif, N. G. Akhmedov, and G. A. O'Doherty, *Chem. Sci.* 2014, **5**, 2230
- The use of a borinic acid toward catalytically-activated prochiral electrophiles was reported during the course of our study, see; (a) R. Z. Li, H. Tang, K. R. Yang, L. Q. Wan, X. Zhang, J. Liu, Z. Fu and D. Niu, *Angew. Chem., Int. Ed.*, 2017, **56**, 7213. (b) R.-Z. Li, H. Tang, L. Wan, X. Zhang, Z. Fu, J. Liu, S. Yang, D. Jia and D. Niu, *Chem.*, 2017, **3**, 834.
- For arylation, see; N. Hamaguchi, M. Kuriyama and O. Onomura, *Tetrahedron: Asymmetry*, 2016, **27**, 177.
- For reviews, see; (a) M. D. Díaz-de-Villegas, J. A. Gálvez, R. Badorrey, and M. P. López-Ram-de-Viú, *Chem. Eur. J.* 2012, **18**, 13920. (b) L. W. Xu, Y. Chen and Y. Lu, *Angew. Chem. Int. Ed.*, 2015, **54**, 9456; (c) A. Borissov, T. Q. Davies, S. R. Ellis, T. A. Fleming, M. S. W. Richardson and D. J. Dixon, *Chem. Soc. Rev.*, 2016, **45**, 5474.
- For selected examples of catalytic kinetic resolutions of vicinal diols, see; acylation (a) S. Kuwano, S. Harada, B. Kang, R. Oriez, Y. Yamaoka, K. Takasu and K. Yamada, *J. Am. Chem. Soc.*, 2013, **135**, 11485; (b) H. Mandai, K. Fujii, H. Yasuhara, K. Abe, K. Mitsudo, T. Korenaga and S. Suga, *Nat. Commun.*, 2016, **7**, 11297; (c) K. Fujii, K. Mitsudo, H. Mandai and S. Suga, *Adv. Synth. Catal.*, 2017, **359**, 2778; silylation (d) Y. Zhao, A. W. Mitra, A. H. Hoveyda and M. L. Snapper, *Angew. Chem. Int. Ed.*, 2007, **46**, 8471-8474. (e) N. Manville, H. Alite, F. Haeffner, A. H. Hoveyda and M. L. Snapper, *Nat. Chem.*, 2013, **5**, 768. (f) X. Sun, A. D. Worthy and K. L. Tan, *J. Org. Chem.*, 2013, **78**, 10494.; acetalization (g) J. H. Kim, I. Čorić, C. Palumbo and B. List, *J. Am. Chem. Soc.*, 2015, **137**, 1778.
- The use of other reported borinic acids (ref 4) gave only a trace of the product with less than 10% ee even under optimized reaction conditions.
- T. Kano, Q. Lan, X. Wang and K. Maruoka, *Adv. Synth. Catal.*, 2007, **349**, 556.
- (a) J. Guin, C. Rabalakos and B. List, *Angew. Chem., Int. Ed.*, 2012, **51**, 8859; (b) A. Arlt, H. Toyama, K. Takada, T. Hashimoto and K. Maruoka, *Chem. Commun.* 2017, **53**, 4779.
- P. Nareddy, F. Jordan, and M. Szostak, *ACS Catal.*, 2017, **7**, 5721.
- (a) T. Ooi, Y. Uematsu, and K. Maruoka, *J. Org. Chem.*, 2003, **68**, 4576; (b) T. Hashimoto and K. Maruoka, *J. Am. Chem. Soc.*, 2007, **129**, 10054.
- A. Biafora, T. Krause, D. Hackenberger, F. Berlitz and L. J. Gooßen, *Angew. Chem., Int. Ed.*, 2016, **55**, 14752.
- T. Furuta, M. Nikaido, J. Yamamoto, T. Kuribayashi and T. Kawabata, *Synthesis*, 2013, **45**, 1312.
- For a review, see; N. Funken, Y. Q. Zhang and A. Gansäuer, *Chem. Eur. J.*, 2017, **23**, 19.
- For catalyst-controlled, regioselective functionalization of vicinal diols, see; (a) J. M. Rodrigo, Y. Zhao, A. H. Hoveyda and M. L. Snapper, *Org. Lett.*, 2011, **13**, 3778. (b) A. D. Worthy, X. Sun and K. L. Tan, *J. Am. Chem. Soc.*, 2012, **134**, 7321. (c) X. Sun, H. Lee, S. Lee and K. L. Tan, *Nat. Chem.*, 2013, **5**, 790. (d) E. Mensah, N. Camasso, W. Kaplan and P. Nagorny, *Angew. Chem., Int. Ed.*, 2013, **52**, 12932. (e) G. Xiao, G. A. Cintron-Rosado, D. A. Glazier, B.-m. Xi, C. Liu, P. Liu and W. Tang, *J. Am. Chem. Soc.*, 2017, **139**, 4346.





Alkylative kinetic resolution of vicinal alcohols is realized by a cooperative chiral ammonium borinate catalysis.

