

A Novel Type of Chiral Diphosphine Ligand, trans-2,3-Bis(diphenylphosphino)-1-methyl-1-cyclopropanecarboxylic Acid and Asymmetric Allylic Alkylation by the Use of Its Palladium Complex

Yoshiharu OKADA, Toru MINAMI,* Tsutomu YAMAMOTO, and Junji ICHIKAWA
Department of Applied Chemistry, Kyushu Institute of Technology,
Sensuicho, Tobata, Kitakyushu 804


Optically active trans-2,3-bis(diphenylphosphino)-1-methyl-1-cyclopropanecarboxylic acid was synthesized from trans-1,2-bis(diphenylphosphinyl)ethene *via* resolution of the racemic diphosphine oxide. Asymmetric allylic alkylation of 2-cyclohexenyl acetate with *l*-menthyl sodiodiethylphosphonoacetate was achieved in good optical yields by the use of its palladium complex.

A variety of chiral diphosphines has been recently synthesized as chelating agents for homogeneous metal catalysts.^{1, 2)} We have previously reported a simple method for the synthesis of optically active cycloalkylphosphines bearing a carboxyl group at the β -position and its important role in inducing high stereoselectivity for the asymmetric allylic alkylation catalyzed by phosphine-palladium complexes.^{3, 4)} We report here the synthesis of a novel type of chiral diphosphine bearing a carboxyl group and its application to asymmetric alkylation.

The synthesis of trans-2,3-bis(diphenylphosphino)-1-methyl-1-cyclopropanecarboxylic acid (**5**) was accomplished as follows (Scheme 1). To a solution of LDA (2.0 mmol) in THF (15 ml) was added *t*-butyl α -chloropropionate (**2**) (2.0 mmol) at -70 °C. After stirring for 30 min, trans-1,2-bis(diphenylphosphinyl)ethene (**1**) (1.0 mmol) was added to the solution. The mixture was stirred overnight at room temperature and refluxed for an additional 1 h. The reaction mixture was quenched with dil. HCl aqueous solution and extracted with CHCl₃. The extracts were washed with H₂O, dried over anhydrous Na₂SO₄, and evaporated. Column chromatography of the residue on silica gel with CHCl₃-ethyl acetate (1:1) gave *t*-butyl trans-2,3-bis(diphenylphosphinyl)-1-methyl-1-cyclopropanecarboxylate (**3**) in 81% yield.⁵⁾

Similar to the resolution of ($\pm$)-DPCB oxide,²⁾ treatment of the racemic mixture of **3** with a stoichiometric quantity of D-(+)-dibenzoyltartaric acid (DBT) in ethanol resulted in the formation of the precipitate, in which diastereoisomer (-)-**3**•(+)-DBT was enriched. After filtration, recrystallization of the precipitate from ethanol gave


pure (-)-**3**•(+)-DBT, mp 123-126 °C. Treatment of (-)-**3**•(+)-DBT with aqueous NaOH led to optically pure (-)-**3**.⁶⁾ Optically pure (+)-**3**⁶⁾ was obtained by a similar resolution of (±)-**3** with (-)-DBT.



Scheme 1.

Reduction of optically pure (-)-**3** (2.0 mmol) was carried out with trichlorosilane (20 mmol) in benzene (10 ml) at 110 °C for 5 h in a sealed tube to give the *t*-butyl diphosphinocarboxylate (-)-**4**⁷⁾ in 97% yield. Subsequent treatment of (-)-**4** with *p*-toluenesulfonic acid in benzene at reflux for 2 h gave optically active (-)-trans-2,3-bis(diphenylphosphino)-1-methyl-1-cyclopropanecarboxylic acid [(-)-**5**]⁸⁾ in 71% yield, $[\alpha]_D^{24} -55.39^\circ$ (c 6.60, CH₂Cl₂). The optical purities of both (-)- and (+)-**5** were over 99% ee, which was determined by HPLC analysis of their diastereomeric amides.⁹⁾

The reaction of 2-cyclohexenyl acetate (**6**) (1.0 mmol) with *l*-menthyl sodiodi-



ethylphosphonoacetate (1.5 mmol) generated from *l*-menthyl diethylphosphonoacetate (**7**)¹⁰⁾ (1.7 mmol) and sodium hydride (60% dispersion in mineral oil, 1.5 mmol) was carried out at 65 °C for 5 h, in the presence of the palladium complex (1.5 mol%) prepared *in situ* by mixing the chiral ligand **5** with palladium acetate (0.15 mmol) in THF (10 ml), which gave the optically active allylic alkylation product **8**. The enantiomeric purity of **8** was similarly determined by HPLC analysis of its conversion into the corresponding diastereomeric amide.⁹⁾ The results are summarized in Table 1.

The reaction of **6** with **7** using the (-)-**5**-palladium complex as a catalyst led to **8** in 58% ee, while the use of (+)-**5** instead of (-)-**5** as a ligand decreased the optical yield of **8**.

Table 1. Asymmetric Allylic Alkylation of 2-Cyclohexenyl Acetate (**6**) with *l*-Menthyl Diethylphosphonoacetate (**7**) Catalyzed by Chiral Phosphine-Palladium Complexes^{a)}

Entry	Ligand	Pd(OAc) ₂ / L*	Product (% yield) ^{b)}	%ee ^{c)}
1	(-) - 5	1 / 0.8	(+) - 8 (100)	61
2	(-) - 5	1 / 1.1	(+) - 8 (86)	58
3	(-) - 5	1 / 1.8	(+) - 8 (88)	48
4	(-) - 5	1 / 2.2	(+) - 8 (94)	36
5	(+) - 5	1 / 1.1	(-) - 8 (79)	26
6	(-) - 4 ^{d)}	1 / 1.0	(+) - 8 (94)	48
7	(-) - DPCBC ^{e)}	1 / 2.6	(+) - 8 (100)	32
8	(+) - DPCBC ^{e)}	1 / 2.2	(-) - 8 (100)	40
9	PPh ₃	1 / 2.4	(+) - 8 (92)	31
10	(+) - DIOP	1 / 1.2	(-) - 8 (85)	7
11	(-) - DIOP	1 / 0.8	(+) - 8 (100)	19

a) Reaction of 1 mmol of **6** with 1.5 mmol of **7** in 10 ml of THF was carried out in the presence of 0.015 mmol of Pd(OAc)₂ at 65 °C for 5 h. b) Isolated yield based on the acetate **6**. c) The enantiomeric purity of **8** was determined by HPLC analysis of diastereomeric amide prepared from **9**¹¹⁾ and (-)-PEA, with a stationary phase column (Nomura Chemical Co., Develosil). d) [α]_D²² -33.11° (c 3.66, CH₂Cl₂). e) DPCBC = (2-Diphenylphosphino)cyclobutanecarboxylic acid.³⁾

to 26% ee (entries 2 and 5). These results indicate that the ligand (-)-**5** matched but the ligand (+)-**5** ill-matched with the nucleophile **7** in this catalytic allylic alkylation. The highest optical yield was obtained by the use of the palladium complex prepared from (-)-**5** and Pd(OAc)₂ in the ratio of *ca* 1:1 (entries 1-4). The replacement of the carboxyl substituent of (-)-**5** with ester remarkably decreased optical yield (entry 6). In contrast, the use of the monophosphine-carboxylic acid (-)- or (+)-DPCBC as a ligand resulted in decreasing optical yields (entries 2, 7, and 8) and exhibited a slight double stereodifferentiation. Furthermore, the use of PPh₃ or DIOP in this alkylation afforded **8** with low enantioselectivity (7-31% ee) (entries 9-11). On the basis of these results, it is evident that the diphosphine bearing the carboxyl group plays an important role in determining enantioselectivity.

Thus, trans-2,3-bis(diphenylphosphino)-1-methyl-1-cyclopropanecarboxylic acid **5** was easily prepared from trans-1,2-bis(diphenylphosphinyl)ethene in optically pure form and this new type of chiral diphosphine is applicable to asymmetric allylic alkylation as an effective ligand for palladium catalyst.

We are grateful for financial support of this work by a Grant-in Aid for Scientific Research on Priority Areas from the Japan Ministry of Education, Science and Culture.

References

- 1) J. M. Brown and P. A. Chaloner, "Homogeneous Catalysis with Metal Phosphine Complexes," ed by L. H. Pignolet, Plenum Press, New York (1983); A. Miyashita, A. Yasuda, T. Ito, H. Takaya, T. Souchi, K. Toriumi, and R. Noyori, *J. Am. Chem. Soc.*, **102**, 7932 (1980).
- 2) T. Minami, Y. Okada, R. Nomura, S. Hirota, Y. Nagahara, and K. Fukuyama, *Chem. Lett.*, **1986**, 613.
- 3) Y. Okada, T. Minami, Y. Sasaki, Y. Umezu, and M. Yamaguchi, *Tetrahedron Lett.*, **31**, 3905 (1990); Y. Okada, T. Minami, Y. Umezu, S. Nishikawa, R. Mori, and Y. Nakayama, *Tetrahedron Asymmetry*, **2**, 667 (1991).
- 4) For Palladium-catalyzed asymmetric allylic alkylation, see: B. M. Trost and P. E. Strege, *J. Am. Chem. Soc.*, **99**, 1649 (1979); T. Hayashi, A. Yamamoto, T. Hagihara, and Y. Ito, *Tetrahedron Lett.*, **27**, 191 (1986).
- 5) **3**: Mp 192-197 °C; IR (KBr) 1720, 1435, 1190, 1120 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.30 (s, 9H, t-Bu), 1.71 (s, 3H, CH_3), 2.10-3.10 (m, 2H, CH), 7.00-8.20 (m, 20H, phenyl H); Anal. Found: C, 70.91; H, 6.19%. Calcd for $\text{C}_{33}\text{H}_{34}\text{O}_4\text{P}_2$: C, 71.21; H, 6.16%.
- 6) (-)-**3**: $[\alpha]_{\text{D}}^{23}$ -55.86° (c 1.13, CH_2Cl_2). (+)-**3**: $[\alpha]_{\text{D}}^{23}$ +54.32° (c 1.04, CH_2Cl_2).
- 7) **4**: Mp 119-121 °C; IR (KBr) 1710, 1430, 1130 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.30 (s, 9H, t-Bu), 1.65 (s, 3H, CH_3), 1.72-2.08 (m, 1H, CH), 2.32-2.82 (m, 1H, CH), 6.90-7.64 (m, 20H, phenyl H); Anal. Found: C, 75.40; H, 6.58%. Calcd for $\text{C}_{33}\text{H}_{34}\text{O}_2\text{P}_2$: C, 75.56; H, 6.53%.
- 8) **5**: Mp 174-179 °C; IR (KBr) 1690, 1430, 690 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.65 (s, 3H, CH_3), 1.76-2.15 (m, 1H, CH), 2.49-2.90 (m, 1H, CH), 6.80-7.60 (m, 20H, phenyl H), 9.59 (br s, 1H, COOH); Anal. Found: C, 74.09; H, 5.67%. Calcd for $\text{C}_{29}\text{H}_{26}\text{O}_2\text{P}_2$: C, 74.35; H, 5.59%.
- 9) The enantiomeric purities of (-)-**5**, (+)-**5**, and **8** were determined by HPLC analysis of the corresponding diastereomeric amides prepared from the carboxylic acids (-)-**5**, (+)-**5**, and **9**¹¹⁾ and (-)- α -methylbenzylamine (PEA) by the use of 2-chloro-pyridinium iodide and triethylamine as condensing agents.
- 10) *l*-Menthyl diethylphosphonoacetate **7** was prepared from *l*-menthol and diethylphosphonoacetic acid by using 2-chloropyridinium iodide and triethylamine, bp 125-130 °C/0.4 mmHg, $[\alpha]_{\text{D}}^{24}$ -44.29° (c 1.00, CHCl_3); IR (neat) 2940, 1730, 1270, 1025, 970 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.76 (d, $J=6.89$ Hz, 3H, CH_3), 0.90 (d, $J=6.74$ Hz, 6H, CH_3), 1.34 (t, $J=7.18$ Hz, 6H, CH_3), 0.70-2.24 (m, 9H, CH_2 and CH), 2.95 (d, $J=21.68$ Hz, CH_2), 3.90-4.42 (m, 4H, CH_2), 4.44-4.96 (br m, 1H, CH).
- 11) The Wittig reaction of **8** with paraformaldehyde in the presence of NaH in THF, followed by hydrolysis with sodium hydroxide in aqueous ethanol gave α -(2-cyclohexenyl)acrylic acid (**9**) in good yield.

(Received January 13, 1992)