

Progress towards viridin: synthesis of the pentacyclic furanosteroid ring system via *o*-benzoquinonoid cycloadditions

Fabio E. S. Souza^a and Russell Rodrigo^{*b}

^a Department of Chemistry, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1


^b Department of Chemistry, Wilfrid Laurier University, Waterloo, Ontario, Canada N2L 3C5.

E-mail: rrodrigo@wlu.ca

Received (in Corvallis, OR, USA) 8th July 1999, Accepted 17th August 1999


The pentacyclic ring system of viridin is synthesised in nine steps from 4-methylguaicol by means of successive cycloadditions involving *o*-benzoquinonoid intermediates generated *in situ*.

The viridin family of pentacyclic furanosteroidal antibiotics¹ isolated from various species of fungi possess several unusual structural features. Viridin **1** contains an aromatic ring C, a



highly oxygenated ring A and a furan ring in a 'triterpenoid' location, betraying its biogenetic origins from lanosterol. Virone **2**, wortmannolone **3**² and wortmannin **4**³ have an additional 'angular' methyl group at C-13 and *trans* C/D ring fusion. Viridin has powerful species specific anti-fungal activity, and wortmannin has attracted some attention as an anti-inflammatory agent and, more recently, as a potent inhibitor of phosphatidylinositol 3-kinase in neutrophils of the guinea pig.⁴

Apart from a long and low yielding conversion⁵ of hydrocortisone to wortmannin no synthesis of any of these compounds has been achieved, but two other studies in this area have been reported.⁶ Our recent explorations in the use of *o*-benzoquinonoid intermediates for natural product synthesis had led to the preparation⁷ of the pentacyclic system **5** of viridin from benzindanone **6** and penta-2,4-dienol in one step (Scheme 1). However, we were unable to selectively reduce the C-10 thiol ester of **5** to the desired methyl group and even worse, an attempt at prior installation of the methyl group as in the model




Scheme 1


substrate 2-methoxy-4-methylnaphthol resulted in failure of the intramolecular Diels–Alder (IMDA) reaction with the pentadienol under the usual oxidative conditions. We now report that the problems associated with the incorporation of the C-10 methyl group have been overcome and the first synthesis of the desired pentacycle **7** has been achieved in nine steps from 4-methylguaicol **8**.

The acid **9** was conveniently prepared on a multi-gram scale from **8** by Kolbe–Schmitt carboxylation^{8,9} followed by BBr₃ demethylation (72% overall). This methylcatechol containing the carbon skeleton of ring B will serve as a linchpin for the attachment of rings CD and AE in two separate annulations employing novel *o*-benzoquinonoid protocols recently developed in our laboratory. When treated with an excess of 1-nitrohepta-4,6-diene¹⁰ in the presence of bis(trifluoroacetoxy)iodobenzene (BTIB) it underwent a sequence of reactions (oxidation to the *o*-quinone, regiospecific Diels–Alder cycloaddition, decarboxylation of the β -keto acid adduct and rearomatization) to provide the dihydronaphthalene **10** in 86% yield. This remarkable 'one-pot' process, in which the evanescent carboxy group of **9** plays a key role,¹¹ is the equivalent of a regiospecific cycloaddition of the 5,6-benzynes derived from 4-methylcatechol. Methylation of **10** with Me₃OBf₄ is both selective and quantitative. The methyl ether **11** was acetylated and subjected to the intramolecular nitrile oxide cycloaddition¹² by treatment with *p*-chlorophenyl isocyanate and Et₃N to produce the isoxazoline **13** after hydrolytic work-up with aqueous base. Catalytic hydrogenolysis (10% Pd/C, boric acid) then generated the benzindanone **14**, representing the BCD rings of viridin, as a single diastereomer¹³ (52% overall, Scheme 2).

Rings A and E were now attached to the benzene ring of **14** by our *o*-benzoquinone monoketal procedure.⁷ Treatment of **14** with BTIB in the presence of excess penta-2,4-dienol produced the monoketal **15** which reacted *in situ* by both possible IMDA pathways to produce a 1:1 mixture of two inseparable *endo*-adducts, pentacycles **16** and **17** (path A, quinonoid double bond as dienophile), and bridged adduct **18** (path B, quinonoid ring as diene). The assignments of relative stereochemistry to **16** and **17** are based on our previous experience^{7,14} with the IMDA reactions of *o*-quinone monoketals, and the *endo* stereochemistry of the adducts is confirmed by the fact that both lose MeOH very readily to form the dienone unit of ring B. *Exo* adducts of this type do not eliminate MeOH.¹⁴ The stereochemical issue with **18** was decided on the assumption that one face of **15** is more crowded than the other because of the *cis* CD ring fusion of this compound; the reaction is therefore presumed to occur at the less congested face of the diene unit of **15**. We are endeavouring to confirm this assignment by an X-ray crystal structure. Bridged adduct **18** was easily separated from the mixture of two diastereomeric pentacycles and when subjected to the Cope rearrangement in refluxing tetrachloroethane provided the target **7** in 65% yield. The diastereomers **16** and **17** when treated with TsOH in benzene with exposure to air provided the same compound (**7**) by dehydration and spontaneous aromatisation of ring C.



Scheme 2 Reagents and conditions: i, K_2CO_3 , CO_2 (800 psi), 200 °C, 4 h; ii, BBr_3 , CH_2Cl_2 , -78 °C to room temp. (72%, two steps); iii, 1-nitrohepta-4,6-diene (5 equiv.), BTIB, THF (86%); iv, K_2CO_3 , Me_3OBF_4 , CH_2Cl_2 (100%); v, Ac_2O , pyridine; vi, $p\text{-ClC}_6\text{H}_4\text{NCO}$, Et_3N , C_6H_6 , room temp., 36 h; vii, NaOH , H_2O -THF (84%, three steps); viii, H_2 , Pd/C , H_3BO_3 , MeOH -THF- H_2O (100%).



Scheme 3 Reactions and conditions: i, BTIB, THF (path A: 33%, path B: 31%); ii, TsOH , C_6H_6 , 45 °C, 4 h (50%); iii, $\text{Cl}_2\text{CHCHCl}_2$, reflux, 48 h (65%); iv, $p\text{-chloranil}$, xylenes, reflux, 36 h (60%).

The nine-step sequence described above is not only the first synthesis of the ring system of viridin, but it also makes **7** available in sufficient quantity (19% overall, Scheme 3) to enable a comprehensive investigation for installation of the oxygen substituents in ring A, by derivatisation of the $\text{C}_2\text{-C}_3$ alkene, to be undertaken. Dehydrogenation of the dihydrofuran

is facile and can be effected at any convenient stage; in fact, the conversion **7** $\rightarrow$ **19** has already been achieved in 60% yield. The benzindanone **14** offers possibilities for modification into the C/D non-aromatic analogues (e.g. **2**, **3** and **4**). Many obvious problems remain before any of these antibiotics can be synthesised, but we regard this short and flexible route to the pentacyclic system¹⁵ as an important first step on the way to members of the viridin family.

We thank the Natural Sciences and Engineering Research Council of Canada for support of this work.

Notes and references

- For a recent review, see J. R. Hanson, *Nat. Prod. Rep.*, 1995, **12**, 381.
- M. M. Blight and J. F. Grove, *J. Chem. Soc., Perkin Trans. 1*, 1986, 1317.
- T. J. Pecher, H.-P. Weber and S. Kis, *J. Chem. Soc., Chem. Commun.*, 1972, 1061.
- T. Okada, L. Sakuma, Y. Fukui, O. Hazeki and M. Ui, *J. Biol. Chem.* 1994, **269**, 3563.
- S. Sato, M. Nakada and M. Shibasaki, *Tetrahedron Lett.*, 1996, **37**, 6141.
- S. Honzawa, T. Mizutani and M. Shibasaki, *Tetrahedron Lett.*, 1999, **40**, 311; C. A. Broka and B. Ruhland, *J. Org. Chem.*, 1992, **57**, 4888.
- R. Carlini, K. Higgs, C. Older, S. Randhawa and R. Rodrigo, *J. Org. Chem.*, 1997, **62**, 2330.
- O. Baine, G. F. Adamson, J. W. Barton, J. L. Fitch, D. R. Swayampati and H. Jeskey, *J. Org. Chem.*, 1954, **19**, 510. We thank Warren Wallace for laboratory assistance with the preparation of **9**.
- The acid **9** had been previously prepared by the CuSO_4 catalysed reaction of 3-bromo-5-methylsalicylic acid with NaOH : D. D. Weller and E. P. Stirchak, *J. Org. Chem.*, 1983, **48**, 4873. We were unable to obtain good yields of **9** by this method in spite of several attempts.
- Prepared from 1-iodohepta-4,6-diene and silver nitrite in 68% yield; see E. Vedejs, T. H. Eberlein and R. G. Wilde, *J. Org. Chem.*, 1988, **53**, 2220 for the synthesis of the iodide.
- R. Carlini, C.-L. Fang, D. Herrington, K. Higgs, R. Rodrigo and N. Taylor, *Aust. J. Chem.*, 1997, **50**, 271.
- A.P. Kozikowski and C.-S. Li, *J. Org. Chem.*, 1987, **52**, 3541 and pertinent references therein.
- D. P. Curran, *J. Am. Chem. Soc.*, 1983, **105**, 5826.
- R. Carlini, K. Higgs, R. Rodrigo and N. Taylor, *Chem. Commun.*, 1998, 65.
- Selected data for **7**: δ_{H} (500 MHz, CDCl_3) 1.51 (s, Me), 2.18 (dd, J 17.8, 3.1, H-1 α), 2.67 (m, H-15), 2.74 (dd, J 17.8, 2.3, H-1 β), 3.67 (m, H-16), 4.00 (app dt, J 10.3, 2.6, H-4), 4.06 (dd, J 10.7, 8.1, H-20 α), 4.87 (dd, J 9.8, 8.1, H-20 β), 5.74 (app s, H-2, H-3), 7.55 (d, J 8.1, H-11), 7.89 (d, J 8.1, H-12). Full details will be published later.

Communication 9/05731E