

ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: C. A. Smith, S. E. Motika, L. Wojtas and X. Shi, *Chem. Commun.*, 2017, DOI: 10.1039/C6CC09794D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

Accessing Alternative Reaction Pathways of the Intermolecular Condensation between Homo-Propargyl Alcohols and Terminal Alkynes through Divergent Gold Catalysis

Courtney A. Smith, Stephen E. Motika, Lukasz Wojtas, Xiaodong Shi*

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

An intermolecular condensation of alkynols and terminal alkynes is reported. Using IPrAuNTf₂, an efficient Au-catalyzed cyclization-alkynylation strategy furnishes (2-arylalkynyl) cyclic ethers in moderate to excellent yields (up to 94%). This strategy is extended to the synthesis of functionalized 2,3-dihydrooxepines via the sequential Au-catalyzed ring expansion of the cyclic ether substrates.


Gold(I) catalysis has developed rapidly in recent years and the resulting methodologies have proved useful for the facile installation of molecular complexity.¹ Gold's intrinsic π -acid mode of activation has rendered it synthetically valuable in promoting reactions involving substrates bearing unsaturated C-C moieties (alkyne, allene, and alkene; Scheme 1A). Consequently, the design of selective catalytic systems is often problematic when multiple reactive functional groups are present.² Furthermore, additional challenges in attaining selectivity arise if these catalytic systems are a composite of intra- and inter- molecular reactions.³

Over the past several years, our group has demonstrated the application of 1,2,3-triazole gold(I) (TA-Au) catalysts in promoting these challenging transformations.⁴ One relevant example is the intermolecular condensation between homo-propargyl alcohols and terminal aliphatic alkynes to afford 2,3-dihydrooxepines (Scheme 1B).⁵ Within this reaction pathway, a vinyl ether intermediate is selectively formed via the intermolecular gold-catalyzed hydroalkoxylation of the terminal alkyne. Moreover, in exploring the reactivity of vinyl ethers with terminal alkynes under triazole gold-catalytic conditions, we later developed an intermolecular alkynylation of vinyl ethers.⁶

Department of Chemistry, University of South Florida, Tampa, FL 33620, USA

* E-mail: xmshi@usf.edu

† Electronic Supplementary Information (ESI) available: Experimental procedures and spectroscopic data of new compounds. CCDC 1521359, 1521360, and 1521361. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c000000x/




Scheme 1. Gold-catalyzed C-C multiple bond activation

Despite these recent advances, certain concerns regarding the mechanisms of the reactions have not been completely addressed. First, the enyne cycloisomerization reaction between homo-propargyl alcohol **1** and terminal alkyne **2** was tolerated only by aliphatic alkynes (Scheme 1B). This result was unexpected considering that gold π -acid activation of aryl-substituted terminal alkynes is extensively documented.⁷ A messy reaction mixture was obtained when aryl-substituted terminal alkynes such as phenylacetylene were used. Second, for the alkynylation of vinyl ether **4**, aliphatic alkynes such as 1-hexyne resulted in slightly lower yields in comparison with aromatic alkynes (Scheme 1C). Based on these observations, changing the substrate of either reaction has the potential to dramatically alter the reaction pathway, and this phenomenon warrants further investigation.

Our strategy in exploring this unique catalytic system was focused on: 1) extending the reaction scope of the 2,3-dihydrooxepine synthesis to include aryl-substituted terminal

COMMUNICATION

alkynes and, 2) exploring the possibility of a cascade process for the alkylation of vinyl ether **B** (Scheme 2).⁸⁻⁹



Scheme 2. Proposed reaction pathways

As shown in Table 1, the reaction of **1a** with aromatic alkyne **2b** under our previously reported conditions resulted in a messy reaction, with no formation of dihydrooxepine **3a** (entry 2). To explore the plausibility of a cascade process, we screened various gold catalysts (see ESI[†]). Interestingly, when 5% RuPhosAuNTf₂ was used as the catalyst, alkyne **6b** was observed, albeit in low yield (31%, entry 3). Since the formation of **6b** requires gold π -acid activation of the internal alkyne of **1a**, we increased the reaction temperature with the objective of promoting the intramolecular cyclization of **1a** over intermolecular hydroalkoxylation. Gratifyingly, the reaction yield was improved to 61% (entry 4). Finally, ligand screening revealed IPrAuNTf₂ as the optimal catalyst, affording the desired product **6b** in excellent yield (5% loading, 91% yield, entry 5; 4% loading, 87% yield, entry 6). Having identified the optimal conditions, we turned our attention to exploring the reaction scope (Table 2).

Table 1. Optimization of reaction conditions^{a,b}


Entry	Conditions	2 (eq)	convn (%)	yield (%)	
				3	6
1	5% [XPhosAu(TA)]OTf, 1% Cu(OTf) ₂ , toluene, rt, 24h	2a (3)	100	84%	n.d.
2	Same conditions as entry 1, messy reaction	2b (3)	100	<5%	n.d.
3	5% RuPhosAuNTf ₂ , CHCl ₃ , rt, 12h	2b (3)	100	n.d.	47%
4	5% RuPhosAuNTf ₂ , CHCl ₃ , 40 °C, 4h	2b (3)	100	n.d.	61%
5	5% IPrAuNTf ₂ , CHCl ₃ , 40 °C, 4h	2b (3)	100	n.d.	91%
6	4% IPrAuNTf ₂ , CHCl ₃ , 40 °C, 4h	2b (3)	100	n.d.	87%
7	1% IPrAuNTf ₂ , CHCl ₃ , 40 °C, 4h	2b (3)	100	n.d.	34%
8	5% IPrAuNTf ₂ , CHCl ₃ , 40 °C, 4h	2b (2)	100	56%	36%
Other catalysts (5) including AuCl ₃ , HOTf,					
9	pTsOH•H ₂ O, In(OTf) ₃ , Pd(OAc) ₂ , CuBr, Cu(OTf) ₂ , ZnBr ₂ , AgOTf	2b (3)	<30%	n.d.	<5%

^a General conditions: **1** (0.2 mmol), **2** (eq), and gold catalyst (mol %) in CHCl₃ (0.8 mL) under Ar; ^b The yield was determined by ¹H-NMR using 1,3,5-trimethoxybenzene as the internal standard.

A variety of phenylacetylene derivatives were subjected to the optimized reaction conditions and subsequently afforded the desired products in good yield (**6b-6l**). Other aromatic alkynes such as ferrocenyl (**6m**, structure confirmed by X-ray), thiophenyl (**6n**), and pyrenyl (**6o**) derivatives could also be utilized in the reaction. Additionally, alkynes bearing adamantyl (**6p**), indole (**6s**), and benzofuran (**6t**) moieties were

tolerated without significant decomposition of the product. The amino acid derivative **6q** and estrone derivative **6r** were also prepared, demonstrating the tolerance of the reaction conditions to a variety of functional groups. Conversely, non-aromatic terminal alkynes such as cyclopentyl acetylene, 1-hexyne, and TMS acetylene, did not form the desired alkylation product.

Table 2. Reaction scope of alkylation cascade^{a,b,c}




^a General conditions: **1** (0.2 mmol), **2** (3 eq), IPrAuNTf₂ (3 mol %) in CHCl₃ (0.8 mL) under Ar at 40 °C. Yields of isolated products are given. Reference the ESI for detailed conditions. ^b Reaction heated at 60 °C. ^c Ratio determined via ¹H-NMR.

To fully investigate the scope of this alkylation reaction, different alkynols were also assessed. Substitution at the *para* (**6aa-6ae**), *meta* (**6af-6ah**), and *ortho* (**6ai**) position of the benzene ring on the homopropargyl alcohol afforded good yields. Additionally, the naphthyl derivative **6aj** could be furnished in good yield (81%). To probe the *endo/exo* selectivity of the cyclization step, we used a series of extended-chain internal alkynols bearing aliphatic or phenyl substitution. Aliphatic substitution selectively yielded the *exo*

product (**6ak**), while phenyl substitution resulted in a 10:3 mixture of *exo:endo* isomers (**6al**). Terminal alkynes were tolerated under the reaction conditions, yielding the 5-*exo* (**6am**), 6-*exo* (**6an**), and 7-*exo* (**6ao**) products. In contrast, the terminal alkyne substrate for 8-*exo* cyclization gave trace formation of the desired product. To determine the diastereoselectivity of the alkylation reaction, we utilized enantiomerically pure alkynols. However, when an α -methyl homopropargyl alcohol is used (**6aq**, dr = 3:2) or an α,β -disubstituted homopropargyl alcohol (**6ap**, dr = 1:1), low diastereoselectivity was exhibited. Next, we diverted our focus to the development of a strategy for the synthesis of the corresponding 2,3-dihydrooxepine **3b**.

During optimization of the reaction conditions (Table 1), we observed that if 2 equivalents of terminal alkyne **2b** were used, the yield of the desired product **6b** was reduced. Monitoring the reaction using $^1\text{H-NMR}$ revealed the formation of dihydrooxepine **3b** as the major by-product (see ESI⁺). Reducing the amount of phenylacetylene to 1.1 equivalents resulted in the formation of **3b** as the major product in 74% yield. Notably, **3b** was unstable upon chromatographic isolation. Therefore, the dienophile tetracyanoethylene was used to trap the product via a Diels Alder cycloaddition¹⁰ and the adduct **7b** was obtained in 71% isolated yield over two steps. The structure of **7b** was unambiguously characterized by X-ray crystallography (see ESI⁺). With these new optimized conditions for the synthesis of 2,3-dihydrooxepines, various aromatic alkynes were subjected to the reaction and moderate yields of the derivatives were obtained (Table 2).

Table 2. Reaction scope of oxepine adducts^a



^a General conditions: **1** (0.4 mmol), **2** (1.1 eq), and IPrAuNTf₂ (5 mol %) in dry CHCl₃ (1.6 mL) under Ar atmosphere at 40 °C. The crude was passed through silica, **8** (1.5 eq) was added, and the reaction was heated at 75 °C. Yields of isolated products are given.

Accordingly, to explore the progression of this process, we monitored the kinetics of the reaction with $^1\text{H-NMR}$. Interestingly, within approximately 15 minutes, homopropargyl alcohol **1a** was completely converted to the

corresponding ketone **1a'** via gold-catalyzed hydrolysis.¹¹ Thus, **1a'** serves as a viable intermediate in the formation of **6b**. After the full conversion of **1a'** to **6b**, a slow conversion of **6b** occurs with concomitant formation of **3b**. Monitoring the analogous reaction in the presence of 3 equivalents of phenylacetylene indicated low yield of **3b** (<30%) after 48 hours (Figure 1).




Figure 1. Reaction kinetic profile

Therefore, this kinetic profile and the intermediary nature of **6b** suggest that, unlike the analogous reaction involving aliphatic terminal alkynes, oxepine **3b** was formed via rearrangement of alkyne **6b**.¹² To confirm this hypothesis, **6b** was isolated and subjected to the reaction conditions (5% IPrAuNTf₂, 40 °C). Within 6 hours, full conversion of **6b** was attained, with oxepine **3b** formed in 63% yield (Figure 2A). Notably, treating **6b** with a catalytic amount of HNTf₂ (5%) led to decomposition of the substrate with no observation of oxepine (Figure 2A). These results confirmed that this ring-expansion rearrangement is a gold-catalyzed process.




Figure 2. Exploration of reaction mechanism

An additional mechanistic concern is the reactivity difference between aliphatic and aryl-substituted terminal alkynes. As shown in Figure 2B, treating either alkyne **1a** (Figure 2C) or ketone **1a'** (Figure 2D) with an equimolar ratio of aliphatic and aromatic alkyne afforded alkyne **6b** as the major product, with no aliphatic oxepine or alkynylation observed. This data suggests that gold-acetylides formed from either aliphatic or aryl-substituted terminal alkynes exhibit divergent reactivity within the catalytic system based on alkyne substitution. Therefore, the intramolecular cyclization of homo-propargyl alcohol **1a** efficiently out-competes the intermolecular hydroxylation of the terminal alkyne, selectively forming the alkynylation product **6b**.

This result contrasts with our previously reported system in which intermolecular hydroalkoxylation was preferred. Furthermore, by utilizing the N-heterocyclic carbene gold catalyst IPrAuNTf₂,¹³ the internal alkyne functionality of compound **6** can be activated to form oxepine **3** through a ring-expansion rearrangement (a detailed mechanistic study is currently under investigation). Consequently, aryl-substituted terminal alkynes are successfully implemented into a highly efficient gold-catalyzed synthesis of 2,3-dihydrooxepines for the first time.

Herein, we report the gold-catalyzed intermolecular condensation of alkynols and aromatic terminal alkynes. It was discovered that the initial alkynylated substrates undergo a subsequent gold-catalyzed ring expansion to afford the corresponding 2,3-dihydrooxepines. This reaction represents a complementary synthesis to our previously reported enyne cycloisomerization. Additionally, the transformation is an illustration of reaction divergence that is tuned by altering the structure of the substrate. As a result of the mild reaction conditions, expedient functional group modification, and mechanistic insight gained, this study holds significant value in the advancement of gold-catalysis for the synthesis of complex molecules.

We thank the NIH (1R01GM120240-01), NSF (CHE-1362057) and NSFC (21629201) for financial support. CAS thanks NSF (HRD-1400837) for financial support.

Notes and references

- For a collection of recent reviews, see: (a) S. Pflasterer and A. S. K. Hashmi, *Chem. Soc. Rev.*, 2016, **45**, 1331–1367; (b) R. Dorel and A. M. Echavarren, *Chem. Rev.*, 2015, **115**, 9028; (c) M. Rudolph and A. S. K. Hashmi, *Chem. Soc. Rev.*, 2012, **41**, 2448; (d) N. Krause and C. Winter, *Chem. Rev.*, 2011, **111**, 1994; (e) A. Fürstner, *Chem. Soc. Rev.*, 2009, **38**, 3208.
- (a) G-Q. Chen, W. Fang, Y. Wei, X-Y. Tang and M. Shi, *Chem. Sci.*, 2016, **7**, 4318; (b) W. Fang, X-Y. Tang and M. Shi, *RSC Adv.*, 2016, **6**, 4047; (c) E. Tomás-Mendivil, P. Y. Toullec, J. Díez, S. Conejero, V. Michelet and V. Cadierno, *Org. Lett.*, 2012, **14**, 2520; (d) B. Alcaide, P. Almendros, S. Cembellín, T. M. del Campo and I. Fernández,

Chem. Commun., 2013, **49**, 1282; (e) R. Meiss, K. Kumar, and H. Waldmann, *Chem. Eur. J.*, 2015, **21**, 13526; (f) J. Miró, M. Sánchez-Roselló, J. González, C. del Pozo and S. Fustero, *Chem. Eur. J.*, 2015, **21**, 5459; (g) W. D. Rao, M. J. Koh, P. Kothandaraman and P. W. H. Chan, *J. Am. Chem. Soc.*, 2012, **134**, 10811; (h) P. B. Brady, E. M. Carreira, *Org. Lett.*, 2015, **17**, 3350.

- Y. Wei and M. Shi, *ACS Catal.*, 2016, **6**, 2515.
- (a) S. E. Motika, Q. Wang, N. G. Akhmedov, L. Wojtas and X. Shi, *Angew. Chem., Int. Ed.*, 2016, **55**, 11582; (b) Y. Yang, A. Qin, K. Zhao, D. Wang and X. Shi, *Adv. Synth. Catal.*, 2016, **358**, 1433. (c) Q. Wang, S. E. Motika, N. G. Akhmedov, J. L. Petersen and X. Shi, *Angew. Chem., Int. Ed.*, 2014, **53**, 5418; (d) H. Duan, S. Sengupta, J. L. Petersen, N. G. Akhmedov and X. Shi, *J. Am. Chem. Soc.*, 2009, **131**, 12100.
- (a) S. Hosseyni, L. Wojtas, M. Li and X. Shi, *J. Am. Chem. Soc.*, 2016, **138**, 3994; (b) V. Snieckus and L. C. Frota, *Synfacts*, 2016, **12**, 573.
- S. Hosseyni, C. A. Smith and X. Shi, *Org. Lett.*, 2016, ASAP.
- (a) W. Wang, G. B. Hammond and B. Xu, *J. Am. Chem. Soc.*, 2012, **134**, 5697; (b) D. J. Gorin and F. D. Toste, *Nature*, 2007, **446**, 395.
- For selected examples of metal-catalyzed oxonium alkynylation, see: (a) X-Q. Yao and C-J. Li, *Org. Lett.*, 2006, **8**, 1953; (b) C. Obradors and A. M. Echavarren, *Chem. Eur. J.*, 2013, **19**, 3547; (c) M. Michalska, O. Songis, C. Taillier, S. P. Bew and V. Dalla, *Adv. Synth. Catal.*, 2014, **356**, 2040; (d) H. Ueda, M. Yamaguchi, H. Kameya, K. Sugimoto and H. Tokuyama, *Org. Lett.*, 2014, **16**, 4948; (e) H. Ueda, M. Yamaguchi and H. Tokuyama, *Chem. Pharm. Bull.*, 2016, **64**, 824.
- For selected examples of metal-catalyzed iminium alkynylation, see: (a) C. Koradin, K. Polborn and P. Knochel, *Angew. Chem., Int. Ed.*, 2002, **41**, 2535; (b) T. F. Knöpfel, P. Zarotti, T. Ichikawa and E. M. Carreira, *J. Am. Chem. Soc.*, 2005, **127**, 9682. (c) J. Han, B. Xu and G. B. Hammond, *J. Am. Chem. Soc.*, 2010, **132**, 916.
- (a) J. Yan, G. L. Tay, C. Neo, B. R. Lee and P. W. H. Chan, *Org. Lett.*, 2015, **17**, 4176; (b) N. V. Borrero, L. G. DeRatt, L. F. Barbosa, K. A. Abboud and A. Aponick, *Org. Lett.*, 2015, **17**, 1754; (c) A. Z. Gonzalez and D. F. Toste, *Org. Lett.*, 2010, **12**, 200.
- T. X. M. Nguyen, B. Son-Meniel, J-C. Jullian and B. Figadere. *Lett. Org. Chem.*, **6**, 630.
- For recent examples of gold-catalyzed ring expansions, see: (a) L-L. Zhu, X-X. Li, W. Zhou, X. Li and Z. Chen, *J. Org. Chem.*, 2011, **76**, 8814; (b) B. Bolte and F. Gagosz, *J. Am. Chem. Soc.*, 2011, **133**, 7696; (c) K-D. Kim, H-S. Yeom, S. Shin, and S. Shin, *Tetrahedron*, 2012, **68**, 5241; (d) W. Yuan, X. Dong, Y. Wei and M. Shi, *Chem. Eur. J.*, 2012, **18**, 10501; (e) J-H. An, H. Yun, S. Shin and S. Shin, *Adv. Synth. Catal.*, 2014, **356**, 3749; (f) J. Zhao, J. Liu, X. Xie, S. Li and Y. Liu, *Org. Lett.*, 2015, **17**, 5926; (g) M. Chen, N. Sun, W. Xu, J. Zhao, G. Wang and Y. Liu, *Chem. Eur. J.*, 2015, **21**, 18571.
- S. Díez-González, N. Marion and S. P. Nolan, *Chem. Rev.*, 2009, **109**, 3612.