

View Article Online View Journal

ChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Z. Qiao, N. Ge and X. Jiang, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC03038B.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

Published on 12 May 2015. Downloaded by Freie Universitaet Berlin on 14/05/2015 01:46:26.



ChemComm

COMMUNICATION

CO₂-Promoted Oxidative Cross-Coupling Reaction for C-S Bond Formation *via* Masked Strategy in an Odourless Way

Received 00th January 20xx, Accepted 00th January 20xx

Zongjun Qiao,^a Nanyang Ge,^a and Xuefeng Jiang^{*ab}

DOI: 10.1039/x0xx00000x

www.rsc.org/

Cu-catalyzed direct oxidative cross-coupling between boronic acids and masked sulfides delivering thioethers was described, in which the SO_3^- , as a mask, has shown a distinctive effect on oxidative cross-coupling condition. Disulfide could be suppressed efficiently via masked strategy under CO_2 atmosphere. A broad scope of aromatics and scalable processes indicates its practicality, which could be further applied to drug late-stage modification and unsymmetrical dibenzothiophenes (DBTs) synthesis.

Transition-metal-catalyzed oxidative cross-coupling¹ between boronic acids and heteroatom nucleophiles has been well known as the Chan-Lam reaction² and emerged as a powerful protocol for the construction of carbon-nitrogen bonds³ or carbon-oxygen bonds,⁴ due to a variety of readily available boronic acids and their derivatives.⁵ However, direct cross-coupling between metallic nucleophile and sulfide has been less studied,⁶ despite its importance in the pharmaceutical industry,⁷ material science,⁸ and food chemistry⁹ in the race toward organosulfur compounds synthesis. There are three main reasons for this: a) strong coordination of the lone pair electrons on sulfur to the transition metal; b) preferential oxidization to disulfide when thiol meets oxidant; and c) unpleasant smell during the experimental process. Several couplings have been tried to construct carbon-sulfur bonds through boronic acids mainly in virtue of stoichiometric metal, special ligand, or by formation from disulfide or mercaptan with unpleasant smell.¹⁰ Our group has focused on introducing a designed mask to naked sulfur, which should possess certain properties including electron effect, steric hindrance and resonance stabilization (Fig. 1). Based on our odourless sulfur atom transfer (SAT) findings¹¹ and inspired by the first single crystal X-ray diffraction analysis¹² of the structure I (Bunte salts¹³), we envisioned that SO_3^- should be the right candidate for the mask because of its unique properties: a) an electron withdrawing group preventing the strong coordination to metal; b) a sustained bulky group preventing homo-coupling; and c) a conjugated group tuning the electron effect on sulfur. In this manuscript, we will describe an odourless oxidative cross-coupling reaction through the concept of the combination of mask-effect and enabling atmosphere (Scheme 1).





Scheme 1. Oxidative sulfurating cross-coupling reaction

We commenced our study with sodium *S*-*p*-tolyl sulfothioate $1a^{14}$ and phenylboronic acid 2a under standard conditions, (see ESI) which afforded desired product 3a in 85% yield with less than 2% of disulfide 4a under 80 °C (Fig. 2). However, when 4-methylbenzenethiol served as a sulfur source, it tended to oxidize to 4a in 49% yield. Both copper trifluoromethanesulfonate and 1,10-phenanthroline are necessary in this system, which indicated the reaction is not capable of undergoing un-catalyzed nucleophilic substitution. The consumption of trifluoromethanesulfonic acid was

^{a.} Shanghai Key Laboratory of Green Chemistry and Chemical Process, Department of Chemistry, East China Normal University, 3663 North Zhongshan Road, Shanghai, 200062, P. R. China.

^{b.} State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Lu, Shanghai 200032, P. R. China.

⁺ Electronic Supplementary Information (ESI) available: Experimental procedures and spectroscopic data for all new compounds. CCDC: 1034135 and 1034136. For ESI and crystallographic data in CIF or other electronic format See DOI: 10.1039/x0xx00000x

COMMUNICATION

addressed depending on different reactant. During the course of condition optimization, the unique effect of carbon dioxide was detected, which could give better conversion and yield compared to inert gas (N_2), reducing gas (CO, H_2) or oxidizing gas (O_2).



Figure 2. Dramatic atmosphere effect in oxidative sulfurating cross-coupling reaction

The interaction between organic thiosulfate salt and carbon dioxide was preliminarily investigated. Carbon dioxide was bubbled into sodium S-3-methoxyphenyl sulfothioate in methanol at ambient temperature for 1 h, affording CO₂-bubbled thiosulfate salt. There is no sharp difference between bubbled and unbubbled organic thiosulfate salt in the ¹H NMR signal, which meant that a chemical change has not taken place. Two kinds of thiosulfate salts were investigated through ¹³C NMR spectroscopy as well, which showed that one more peak (at about 126.3 ppm) was detected when using CO₂-bubbled (see ESI). Followed by thermogravimetric analysis (TGA) in the temperature range of 30-800 °C under a flow of nitrogen, the TG curve of CO₂-bubbled thiosulfate salt displayed one more stage, which showed a rapid initial weight loss of 2.0 wt % before the release of water molecules in the temperature range of 90–100 $^{\circ}$ C (Fig. 3A). N₂, O₂, H₂, or CO was bubbled into organic thiosulfate salt, which has not shown similar stage (Fig. 3B). These evidences implied that CO₂ could partially prevent the formation of disulfide in the form of physical absorption but not in a chemical way.





Figure 3. TGA curves for sodium S-3-methoxyphenyl sulfothioate (A) DSC-TG spectra of CO₂-bubbled organic thiosulfate salt (B) DSC-TG spectra of organic thiosulfate salt bubbled by different gases.

Control experiments showed that the desired product **3a'** could be obtained in a yield of 63% by CO₂-bubbled thiosulfate salt (Table 1, entry 1), which was similar to unbubbled salt under the CO₂ condition (Table 1, entry 2). However, unbubbled thiosulfate salt under a N₂ atmosphere only achieved 37% yield, in which more disulfide has been produced in the yield of 27% (Table 1, entry 3). Alternatively, additives, which could generate CO₂ in situ, were subjected to the reaction under a N₂ atmosphere, inhibiting disulfide in different levels (Table 1, entries 4-6). These current observations from the physical absorption phenomenon and control experiments demonstrated that the enabling atmosphere played the role of stabilization and slow oxidation of the revealed thiolate to disulfide, perhaps by the formation of carbonate, bicarbonate, or carbonic acid. Table 1. Control Experiments

Phenylboronic acid 4(OTf)₂, 1,10-ph Ag₂CO₃, DTBP ____K₃PO₄ Cu(OŤ SO₂Na Disulfid DMF, 80 °C Gas, Additive 39 entry thiosulfat gas additive 3a' disulfide CO₇-bubbled N₂ 63% <2% 1 <2% 2 unbubbled CO₂ 1 69% unbubbled N_2 37% 27% N_2 K₂CO₃ <2% unbubbled 33% unbubbled N₂ Na₂CO₂ 46% <2% 5 unbubbled N_2 NaHCO₃ 55% <2% 6

The scope of this catalytic sulfuration was investigated in Table 2. Generally, a variety of arylboronic acids, bearing both electronwithdrawing and -donating groups at the para-, meta-, or orthopositions afford moderate to excellent yields (**3a-3l**). All halogen substituted substrates proceed smoothly, such as F (**3m**), Cl (**3n-3p**), Br (**3q**), and even I (**3r**), which was rarely benign in traditional crosscoupling. On the other hand, diversity of organic thiosulfate salts were also examined, which indicated that the electronic effect from sulfur sources did not affect the efficiency of the transformations either. The reaction was further extended to scalable processes, in which sodium *S*-3-methoxyphenyl sulfothioate (**1.2** g) could be efficiently converted to **3a'** in 84% yield with even reduced loading of catalyst (see ESI). Remarkably, the boronic acids embedded in the tocopherol framework could provide thioether-substituted

DOI: 10.1039/C5CC03038B

ChemComm

^aSodium S-p-tolyl sulfothioate served as sulfur source. ^b4-methylbenzenethiol served as sulfur source.

Published on 12 May 2015. Downloaded by Freie Universitaet Berlin on 14/05/2015 01:46:26.

View Article Online DOI: 10.1039/C5CC03038B COMMUNICATION

Table 2. Substrate Scope^{a, b}



^aReaction conditions: **1** (0.1 mmol), **2** (0.15 mmol), Cu(OTf)₂ (20 mol%), 1, 10-phen (40 mol%), Ag₂CO₃ (0.12 mmol), K₃PO₄ (0.14 mmol), DTBP (0.1 mmol), and DMF (2.0 mL); 80 °C; 7 h. ^bIsolated yields. ^c0.03 mmol of CF₃SO₃H was added.



^aReaction conditions: **1** (0.1 mmol), **2** (0.2 mmol), Cu(OTf)₂ (20 mol%), 1, 10-phen (40 mol%), Ag₂CO₃ (0.12 mmol), K₃PO₄ (0.14 mmol), DTBP (0.1 mmol), and DMF (2.0 mL); 80 °C; 7 h. ^bIsolated yield. ^cCu(OTf)₂ (20 mol%) was replaced by CuCl₂. ^dCu(OTf)₂ (0.1 mmol), 1, 10-phen (0.1 mmol).

Table 4. Application in Parmaceutical and Material



 $R^{1} \underbrace{f_{1}}_{3} \underbrace{f_{2}}_{3} \underbrace{ref.19}_{R^{2}} \underbrace{f_{1}}_{R^{1}} \underbrace{f_{2}}_{R^{2}} \underbrace{f_{2}} \underbrace{f_{2}}_{R^{2}} \underbrace{f_{2}}_{R^{2}} \underbrace{f_{2}} \underbrace{f_{2}}_{R^{2}} \underbrace{f_{2}} \underbrace{f_{$

DOI: 10.1039/C5CC03038B

ChemComm

COMMUNICATION

tocopherol **3b'** in 73% yield, which has shown potential value in complex molecule modification.

Heteroaromatics are always crucial and challenge in drug discovery,¹⁵ whose coupling was also demonstrated in Table 3. Compounds containing nitrogen (**5a**, **5b**, and **5c**), oxygen (**5d**), and sulfur (**5e** and **5f**) atoms all performed well. Ferrocene-containing product **5g**¹⁶ could be achieved in moderate yield, which provides the opportunity for developing new sulfur-containing ferrocene ligands.

This method could be readily applied to drug late-stage diversification (Table 4A).¹⁷ Estratrien, a kind of female hormone drug, could be easily transformed to the corresponding organic thiosulfate salt in 94% yield, to afford sulfur-modified estratrien 6a followed by oxidative sulfurating cross-coupling. Sulfur-decorated Fenofibrat 6b could derive from its boronic acid ester efficiently through a similar transformation. This method offers great potential for the generation of drug analogue libraries. From another aspect, this method could be carried out to establish unsymmetrical dibenzothiophenes (DBTs), which consititute a core framework in certain kinds of organic compounds including pharmaceuticals, photoactive compounds, dyes, liquid crystals, and conducting polymers (Table 4B).^{8,18} For example, benzo[*a*]naphtho[1,2b]thiophene which was substituted by OMe, F, or Cl could be obtained untrammeled through this protocol followed by oxidative dehydrogenative cyclization.¹⁹

In conclusion, relying on the versatile effect of SO_3^- and CO_2 , sulfur could be compatible with oxidizing condition in an odourless way. A dramatic masked strategy and an interesting CO_2 accelerated phenomenon were uncovered, which have been studied by TGA. The extensive adaptability of this sulfuration present in both reactants renders this reaction highly efficient and practical for the synthesis of medicinally important and materially unobtainable units using readily available boronic acids. Further explorations for more synthetic applications are ongoing in our laboratory.

Financial support was provided by NBRPC (973 Program, 2015CB856600), NSFC (21472050, 21272075), NCET (120178), DFMEC (20130076110023), Fok Ying Tung Education Foundation (141011), the program for Shanghai Rising Star (15QA1401800), Professor of Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Learning, and the Changjiang Scholar and Innovative Research Team in University.

Notes and references

- (a) S. V. Ley, A. W. Thomas, Angew. Chem. Int. Ed., 2003, 42, 5400.
 (b) E. M. Beccalli, G. Broggini, M. Martinelli, S. Sottocornola, Chem. Rev., 2007, 107, 5318.
 (c) C. Liu, H. Zhang, W. Shi, A. Lei, Chem. Rev., 2011, 111, 1780.
- (a) D. M. T. Chan, K. L. Monaco, R. Wang, M. P. Winters, *Tetrahedron Lett.*, **1998**, *39*, 2933. (b) D. A. Evans, J. L. Katz, T. R. West, *Tetrahedron Lett.*, **1998**, *39*, 2937. (c) P. Y. S. Lam, C. G. Clark, S. Saubern, J. Adams, M. P. Winters, D. M. T. Chan, A. Combs, *Tetrahedron Lett.*, **1998**, *39*, 2941. (d) J. X. Qiao, P. Y. S. Lam, *Syn.* **2011**, *6*, 829.
- 3 C-N bonds formation through boronic acids: (a) J. C. Antilla,
 S. L. Buchwald, Org. Lett., 2001, 3, 2077. (b) R. E. Shade, A.
 M. Hyde, J. C. Olsen, C. A. Merlic, J. Am. Chem. Soc., 2010,
 132, 1202. (c) C. Zhu, G. Li, D. H. Ess, J. R. Falck, L. Kurti, J.
 Am. Chem. Soc., 2012, 134, 18253.

- 4 C-O bonds formation through boronic acids: (a) A. Kina, H. Iwamura, T. Hayashi, *J. Am. Chem. Soc.*, **2006**, *128*, 3904. (b) M. Dowlut, D. G. Hall, *J. Am. Chem. Soc.*, **2006**, *128*, 4226. (c) S. N. Mlynarski, A. S. Karns, J. P. Morken, *J. Am. Chem. Soc.*, **2012**, *134*, 16449.
- 5 D. G. Hall, Boronic Acids, 2nd edn., Wiley-VCH: Weinheim, 2011. For reviews, please see: (b) N. Miyaura, A. Suzuki, *Chem. Rev.*, **1995**, *95*, 2457. (c) R. Martin, S. L. Buchwald, *Acc. Chem. Res.*, **2008**, *41*, 1461. (d) C. Zhu, J. R. Falck, *Adv. Synth. Catal.*, **2014**, *356*, 2395.
- 6 (a) T. Kondo, T. A. Mitsudo, *Chem. Rev.*, 2000, 100, 3205. (b)
 I. P. Beletskaya, V. P. Ananikov, *Chem. Rev.*, 2011, 111, 1596. (c)
 (c) H. Liu, X. Jiang, *Chem. Asian J.*, 2013, *8*, 2546. (d) X. Xu, J. Liu, J. Zhang, Y. Wang, Y. Peng, *Org. Lett.*, 2013, 15, 550. (e)
 C. Lee, Y. Liu, S. S. Badsara, *Chem. Asian J.*, 2014, 9, 706.
- 7 (a) A. Q. Acton, Ed. Sulfur Compounds: Advances in Research and Application; Scholarly Editions: Atlanta, GA, 2012. (b) E. A. Ilardi, E. Vitaku, J. T. Njardarson, J. Med. Chem., 2014, 57, 2832. (c) Sulfur-containing pharmaceuticals can be found at: http://www.cbc.arizona.edu/njardarson/group/.
- 8 (a) A. Mishra, C. Q. Ma, P. Bäuerle, *Chem. Rev.*, **2009**, *109*, 1141. (b) H. Bürchstümmer, A. Weissenstein, D. Bilalas, F. Würthner, *J. Org. Chem.*, **2011**, *76*, 2426. (c) K. Takimiya, I. Osaka, T. Mori, M. Nakano, *Acc. Chem. Res.*, **2014**, *47*, 1493.
- 9 (a) E. Block, Angew. Chem. Int. Ed., 1992, 31, 1135. (b) D. Y. Lin, S. Zhang, E. Block, L. C. Katz, Nature, 2005, 434, 470. (c) N. I. Joyce, C. C. Eady, P. Silcock, N. B. Perry, J. W. van Klink, J. Agric. Food Chem., 2013, 61, 1449.
- 10 (a) P. S. Herradura, K. A. Pendola, R. K. Guy, Org. Lett., 2000, 2, 2019. (b) C. Savarin, J. Srogl, L. S. Liebeskind, Org. Lett., 2002, 4, 4309. (c) N. Taniguchi, J. Org. Chem., 2007, 72, 1241. (d) H. Xu, Y. Zhao, T. Feng, Y. Feng, J. Org. Chem., 2012, 77, 2878.
- (a) Z. Qiao, H. Liu, X. Xiao, Y. Fu, J. Wei, Y. Li, X. Jiang, Org. Lett., 2013, 15, 2594. (b) Z. Qiao, J. Wei, X. Jiang, Org. Lett., 2014, 16, 1212. (c) Y. Li, J. Pu, X. Jiang, Org. Lett., 2014, 16, 2692. (d) Y. Zhang, Y. Li, X. Zhang, X. Jiang, Chem. Commun., 2015, 51, 941. (e) X. Xiao, M. Feng, X. Jiang, Chem. Commun., 2015, 51, 4208.
- 12 CCDC-1034136 (I): See the ESI⁺ for details.
- (a) H. Bunte, *Chem. Ber.*, **1874**, *7*, 646. (b) J. T. Reeves, K. Camara, Z. S. Han, Y. Xu, H. Lee, C. A. Busacca, C. H. Senanayake, *Org. Lett.*, **2014**, *16*, 1196.
- 14 The reaction proceeded from blue liquid to a viscous mixture under N₂ atmosphere, the salts could be obtained by recrystallization in mixed solvent (volume ratio 1:1) of NaCl_(aq) and *n*-hexane.
- 15 (a) A. John, K. M. Joule, Heterocyclic Chemistry, 5th ed.; Wiley: New York, 2010. (b) D. J. S. Jean, Jr., C. Fotsch, *J. Med. Chem.*, **2012**, 55, 6002. (c) M. A. Düfert, K. L. Billingsley, S. L. Buchwald, *J. Am. Chem. Soc.*, **2013**, *135*, 12877.
- 16 CCDC-1034135 (5g): See the ESI⁺ for details.
- 17 (a) E. Lee, A. S. Kamlet, D. C. Powers, C. N. Neumann, G. B. Boursalian, T. Furuya, D. C. Choi, J. M. Hooker, T. Ritter, *Science*, **2011**, *334*, 639. (b) J. Wencel-Delord, F. Glorius, *Nat. Chem.*, **2013**, *23*, 369. (c) T. Kang, Y. Kim, D. Lee, Z. Wang, S. Chang, *J. Am. Chem. Soc.*, **2014**, *136*, 4141.
- 18 (a) S. Allard, M. Forster, B. Souharce, H. Thiem, U. Scherf, Angew. Chem. Int. Ed., 2008, 47, 4070. (b) K. Niimi, S. Shinamura, I. Osaka, E. Miyazaki, K. Takimiya, J. Am. Chem. Soc., 2011, 133, 8732. (c) A. Fukazawa, D. Kishi, Y. Tanaka, S. Seki, S. Yamaguchi, Angew. Chem. Int. Ed., 2013, 52, 12091. (d) T. Mori, T. Nishimura, T. Yamamoto, I. Doi, E. Miyazaki, I. Osaka, J. Am. Chem. Soc., 2013, 135, 13900.
- (a) H. Petersen, K. P. Zeller, *Synthesis*, **1975**, 532. (b) R. Sanz,
 Y. Fernández, M. P. Castroviejo, A. Pérez, F. J. Fañanás, *J. Org. Chem.*, **2006**, *71*, 6291. (c) R. Che, Z. Wu, Z. Li, H. Xiang,
 X. Zhou, *Chem. Eur. J.*, **2014**, *20*, 7258.