

133. Reassignment of the Configuration of Several Keto-cyclolignans Prepared from Podophyllotoxin


by José María Miguel del Corral, Marina Gordaliza, José-Luis López*, Esther del Olmo, M. Angeles Castro,
and M. Luisa López

Departamento de Química Orgánica, Facultad de Farmacia, Universidad de Salamanca,
Avda. Campo Charro s/n, E-37007 Salamanca

(6. IV. 95)

The configuration of several keto-cyclolignans related to podophyllotoxin has been reviewed. Under basic catalysis, the configuration at the C-atom in α -position to the lactone carbonyl group in podophyllotoxone is inverted instead of the C-atom in α -position to the ketone group, as it has been reported.

Last year, in a paper published in this Journal, Höfert and Matusch reported a 'novel rearrangement of podophyllotoxone' [1]. They described the epimerization of podophyllotoxone (**1**) at C(3) α to the C(4)=O, and the formation of isopicropodophyllone (**2**), in equilibrium with **1**, when treating the former substance with BuLi in refluxing Et₂O. Subsequently, preparation of the unsaturated keto-acid **3** and several other derivatives were reported.



It is well known that the *trans*-fused lactone podophyllotoxin (**4**) and its derivatives readily epimerize at C(2), leading to the more stable *cis*-fused picro analogues, in basic or even neutral media [2]. However, those authors assigned the structure of **2** to the epimerization product, based on the speculative assumption that the greater acidic character of H at C(3) as compared to H at C(2) is sufficient to favor the transformation of **1** into **2**. They did not provide any experimental proof or calculations supporting that assumption.

We believe that Höfert and Matusch did not realize that the spectral properties they observed for compounds **2** and **3** almost perfectly matched those published by us for

picropodophyllone (**6**) and thuriferic acid (**5**), respectively [3], because no reference to our paper, reporting the structure assignment of **5** and related compounds, was mentioned in [1].


The aim of this communication is to correct the structures proposed by *Höfert* and *Matusch* for most of the compounds reported in [1].

Thus, podophyllotoxone (**1**), prepared by oxidation of podophyllotoxin (**4**) [4], was equilibrated in refluxing AcOH to obtain a mixture containing the C(3)-epimer, isopodophyllotoxone (**2**; *Scheme*). Further, picropodophyllin (**7**), prepared by epimerization of podophyllotoxin (**4**), was transformed into picropodophyllone (**6**) through PDC oxidation [5]. NMR Data for the keto derivatives are shown in *Tables 1* and *2*.

Spectral data of **2** coincide with those reported in [6] for isopodophyllotoxone, whereas data for compound **6** is in agreement with those reported by *Höfert* and *Matusch* [1] for the assigned structure of isopodophyllotoxone (named isopodophyllotoxone in [1]).

Finally, the treatment of either **1** and **6** with 1% KOH in MeOH afforded thuriferic acid (**5**) as the only product in a fair yield, demonstrating that epimerization at C(2) occurred preferentially with respect to C(3) under basic treatment of podophyllotoxone.

It seems obvious that the arguments presented by *Höfert* and *Matusch* failed to justify the structure of the product of the reaction of podophyllotoxone with BuLi. It is evident from calculations¹⁾ (*Fig.*) that H–C(3), α to the keto group, is more acidic than H–C(2),




¹⁾ Atomic charges have been calculated from the lowest-energy conformer using *Stewart's* Hamiltonian in MOPAC [7].

Table 1. $^1\text{H-NMR}$ Data (CDCl_3) for Keto-cyclolignans **1**, **2**, and **6**

H-Atom	1	2	6
H–C(1)	4.85 (<i>d</i> , <i>J</i> = 4.2)	4.57 (<i>d</i> , <i>J</i> = 5.2)	4.70 (<i>s</i>)
H–C(2)	3.33 (<i>dd</i> , <i>J</i> = 15.5, 4.2)	3.60–3.63 (<i>m</i>)	3.29 (<i>m</i>)
H–C(3)	3.52 (<i>ddd</i> , <i>J</i> = 15.2, 9.6, 7.3)	3.60–3.63 (<i>m</i>)	3.29 (<i>m</i>)
H–C(3a)	4.35 (<i>t</i> , <i>J</i> = 9.6); 4.55 (<i>dd</i> , <i>J</i> = 9.6, 7.3)	3.87 (<i>m</i>); 4.48 (<i>m</i>)	4.36 (<i>dd</i> , <i>J</i> = 9.3, 3.9); 4.77 (<i>d</i> , <i>J</i> = 9.3)
H–C(5)	7.52 (<i>s</i>)	7.38 (<i>s</i>)	7.50 (<i>s</i>)
H–C(8)	6.71 (<i>s</i>)	6.66 (<i>s</i>)	6.70 (<i>s</i>)
H–C(2')	6.34 (<i>s</i>)	6.28 (<i>s</i>)	6.21 (<i>s</i>)
H–C(6')	6.34 (<i>s</i>)	6.28 (<i>s</i>)	6.21 (<i>s</i>)
OCH ₂ O	6.07 (<i>d</i> , <i>J</i> = 1.5); 6.10 (<i>d</i> , <i>J</i> = 1.5)	6.05 (<i>d</i> , <i>J</i> = 1.5); 6.10 (<i>d</i> , <i>J</i> = 1.5)	6.00 (<i>s</i>)
MeO–C(3')	3.75 (<i>s</i>)	3.72 (<i>s</i>)	3.75 (<i>s</i>)
MeO–C(4')	3.81 (<i>s</i>)	3.79 (<i>s</i>)	3.81 (<i>s</i>)
MeO–C(5')	3.75 (<i>s</i>)	3.72 (<i>s</i>)	3.75 (<i>s</i>)

Table 2. $^{13}\text{C-NMR}$ Data (CDCl_3) for Keto-cyclolignans **1**, **2**, and **6**

C-Atom	1	2	6	C-Atom	1	2	6
C(1)	44.8	44.8	43.3	C(10)	128.3	128.8	127.1
C(2)	46.7	45.1	46.3	C(1')	132.2	133.9	137.3
C(2a)	173.5	175.2	175.3	C(2')	108.0	107.0	104.8
C(3)	43.6	44.3	43.1	C(3')	153.2	153.4	153.6
C(3a)	67.0	69.4	70.2	C(4')	138.0	138.9	137.9
C(4)	192.4	194.1	193.3	C(5')	153.2	153.4	153.6
C(5)	106.3	106.1	105.8	C(6')	108.0	107.0	104.8
C(6)	148.2	148.4	148.2	OCH ₂ O	102.4	102.2	102.0
C(7)	153.2	153.4	153.6	MeO–C(3')	56.6	56.2	56.0
C(8)	109.7	108.6	109.2	MeO–C(4')	60.8	60.8	40.5
C(9)	141.6	139.1	139.4	MeO–C(5')	56.6	56.2	56.0

Figure. Comparative electronic and stereochemical aspects for podophyllotoxone (**1**).
Calculated with semiempirical MOPAC 6.0.

α to the lactone C=O. Moreover, from the stereochemical point of view, the access of bases to H at C(3) is hindered due to the presence of the trimethoxyphenyl group at C(1), which is pseudoaxially disposed.

In conclusion, practically all the structures proposed for those compounds reported in [1] by Höfert and Matusch should be corrected in the configurations at C(2) and C(3).

Financial support came from the Junta de Castilla y León (SA-64/12/92).

Experimental Part

General. Column chromatography: performed over silica gel (0.063–0.2 mm). Flash chromatographies (FC): with 3–85 ml/min flow rates, over silica gel (0.040–0.063 mm). TLC was performed on precoated silica-gel polyester plates (0.25-mm thickness) with fluorescent indicator UV_{254} . Solns. of 10% phosphomolybdic acid in EtOH or 10% H_2SO_4 in EtOH were used for visualization, after heating at 110°. PLC was developed on silica gel F_{254} plates. M.p.: determined in silicone bath; uncorrected. IR Spectra: in $CHCl_3$ soln. NMR Spectra: recorded at 200/50 MHz ($^1H/^{13}C$) in $CDCl_3$ soln., chemical shifts (δ) in ppm, referred to internal TMS, and coupling constants (J) in Hz. Mass spectra (EI): recorded under ionization energy of 70 eV.

Picropodophyllin (7). Compound **4** (150 mg) in 10 ml of 1% KOH in MeOH was stirred for 30 min at r.t. After neutralization and extraction with AcOEt, 140 mg (93%) of **7** were obtained.

Podophyllotoxone (1). Compound **4** (500 mg) in 15 ml of dry CH_2Cl_2 was treated with 600 mg of PDC. The suspension was stirred for 3 h at r.t. Usual workup afforded after FC (CH_2Cl_2 /AcOEt 1:1) 416 mg (84%) of **1**.

Picropodophyllone (6). By the same method described for **1**, 200 mg of **6** were obtained from the treatment of 220 mg of **7** with 280 mg of PDC.

Isopicropodophyllone (2). Compound **1** (100 mg) in 7 ml of AcOH was refluxed for 1 h. After addition of H_2O and extraction with AcOEt, the resulting material was chromatographed (Cl_2CH_2 /AcOEt 95:5) yielding 70 mg (70%) of **1** and 22 mg of **2**.

Thuriferic Acid (5). Compound **1** (110 mg) was treated with 5 ml of 1% KOH/MeOH. The mixture was left 30 min at r.t., yielding, after usual workup and FC, 90 mg of **5**. M.p. 92–96° (Et_2O). Spectroscopic and physical data are identical to those reported in [3].

REFERENCES

- [1] P. H. Höfert, R. Matusch, *Helv. Chim. Acta* **1994**, *77*, 771.
- [2] W. J. Gensler, C. D. Gatsonis, *J. Am. Chem. Soc.* **1966**, *31*, 3224.
- [3] A. San Feliciano, J. L. López, M. Medarde, J. M. Miguel del Corral, B. de Pascual-Teresa, P. Puebla, *Tetrahedron* **1988**, *44*, 7255.
- [4] W. J. Gensler, F. Johnson, *J. Am. Chem. Soc.* **1955**, *77*, 3674.
- [5] D. E. Jackson, P. M. Dewick, *Phytochemistry* **1981**, *20*, 2277.
- [6] D. E. Jackson, P. M. Dewick, *Phytochemistry* **1984**, *23*, 1147.
- [7] A General Molecular Orbital Package (Version 6.0). Quantum Chemistry Program Exchange Catalogue (QCPE), 1992.