J. CHEM. SOC., CHEM. COMMUN., 1993

# A Radical Cyclisation Strategy to $\alpha$ -Spiro- $\beta$ -methylene- $\gamma$ -butyrolactones. Total Synthesis of (±)-Norbakkenolide-A

## Adusumilli Srikrishna,\* Sankuratri Nagaraju and G. Veera Raghava Sharma

Department of Organic Chemistry, Indian Institute of Science, Bangalore-560 012, India

The 5-*exo-dig* radical cyclisation reaction of the bromoacetals **9**, obtained from the enol ethers **8**, and treatment of the resulting hemiacetal **10** with Jones' reagent transforms the cyclic ketones **7** and **12** into  $\alpha$ -spiro- $\beta$ -methylene- $\gamma$ -butyrolactones **11** and norbakkenolide-A **2**.

Bakkenolide-A 1, first isolated<sup>1</sup> from the bud of *Petasites japonicus* subsp. *giganteus Kitam*, is the simplest member of the bakkane class of sesquiterpenes containing a novel  $\alpha$ -spirolactone fused hydrindane framework. Interestingly bakkenolide-A, which possesses an unconjugated  $\beta$ -methyl-

ene- $\gamma$ -butyrolactone grouping, shows cytotoxic and antifeedant effects<sup>2</sup> normally associated with the conjugated  $\alpha$ -methylene- $\gamma$ -butyrolactones. In contrast to the many procedures that exist for obtaining  $\alpha$ -methylene- $\gamma$ -butyrolactones,<sup>3</sup> there are very few general methods available for the synthesis of

### J. CHEM. SOC., CHEM. COMMUN., 1993

β-methylene-γ-butyrolactones.<sup>4</sup> The last decade has witnessed a rapid growth in the use of free radical cyclisation reactions in organic synthesis.<sup>5</sup> Stork and coworkers have developed a convenient method for the construction of butyrolactones *via* the 5-*exo-trig* radical cyclisation of mixed bromoacetals of allyl alcohols.<sup>5</sup> Herein we report a four step, efficient and general method for the transformation of cyclic ketones into α-spiroβ-methylene-γ-butyrolactones [eqn. (1)] employing a 5-*exodig* radical cyclisation reaction as the key step, and its extension to the total synthesis of (±)-norbakkenolide-A **2**.



**Table 1** Synthesis of  $\alpha$ -spiro- $\beta$ -methylene- $\gamma$ -butyrolactones<sup>*a*</sup>

It was anticipated that the 5-exo-trig (or dig) radical cyclisation of a suitable cyclic tertiary bromide will generate the requisite spiro centre. It was first tested for the construction of  $\alpha$ -spirobutyrolactone 3 as shown in Scheme 1. Thus, reaction of the enol ether 4, readily available from cyclohexanone, with N-bromosuccinimide (NBS) in the presence of dimethylallyl alcohol furnished the mixed bromoacetal 5. The 5-exo-trig radical cyclisation of the bromoacetal 5 using in situ generated catalytic Bun<sub>3</sub>SnH (Bun<sub>3</sub>SnCl, NaCNBH<sub>3</sub>, ButOH)6 in the presence of a catalytic amount of azoisobutyronitrile (AIBN) furnished, regiospecifically, the spirohemiacetal 6. The hemiacetal 6 was converted into the  $\alpha$ -spirobutyrolactone 3 by sonochemically accelerated reaction with Jones' reagent.7 The methodology has been extended to various  $\beta$ -methylene- $\alpha$ -spirobutyrolactones, as shown in Scheme 2. Thus, Wittig reaction on a cyclic ketone 7 with methoxymethylenetriphenylphosphorane followed by NBS bromination of the resultant enol ethers 8 in the presence of prop-2-ynyl alcohol (or dimethylprop-2-ynyl alcohol) generated the requisite radical precursors, acetylenic bromoacetals

Entry	Ketone	Enol ether <sup>b</sup> (% yield)	Bromoacetal <sup>b</sup> 9 (% yield)	Spiroacetal <sup>b</sup> 10 (% yield)	Lactone (% yield)	
i	<b>○</b> =°	4 (90)	<b>9</b> a† (92)	<b>10a</b> † (86)	<b>11a</b> † (96)	$\sim$
ii	u		<b>9b</b> (52)	<b>10b</b> (70)	<b>11b</b> (63)	$\sim$
iii	◯>=∘	<b>8c</b> (80)	<b>9c</b> (90)	<b>10c</b> (82)	11c (79)	C∱.
iv		<b>8d</b> (90)	<b>9d</b> (97)	<b>10d</b> (91)	11d (92)	$\sim$
v		<b>8e</b> (94)	<b>9e</b> (92)	<b>10e</b> (90)	<b>11e</b> (88)	$\bigvee$
vi	$\sum$	<b>8f</b> (90)	<b>9f</b> (86)	<b>10f</b> (79)	<b>11f</b> ° (90)	
vii		<b>8</b> g (83)	<b>9</b> g (59)	<b>10g</b> (70)	<b>11g</b> <sup>c,d</sup> (70)	
viii	. A H	<b>8h</b> (82)	<b>9h</b> (77)	<b>10h</b> (62)	11h <sup>c,e</sup> (78)	SALS
ix	12	<b>8i</b> (96)	<b>9i</b> (93)	<b>10i</b> (90)	<b>2</b> † (94) <sup>f</sup>	$\sim$

<sup>*a*</sup> All the compounds exhibited satisfactory spectral data. Yields refer to isolated and chromatographically pure products. <sup>*b*</sup> Mixture of stereoisomers. <sup>*c*</sup> Only one stereoisomer is formed. <sup>*d*</sup>  $[\alpha]_D^{24}$  -41.3 (*c*, 0.75, CHCl<sub>3</sub>). <sup>*e*</sup> M.p. 160-162 °C,  $[\alpha]_D^{26}$  63.6 (*c*, 1.4; CHCl<sub>3</sub>). <sup>*f*</sup> 5:1 mixture of norbakkenolide **2** and its C-7 epimer.

NBS

9. The 5-exo-dig radical cyclisation reaction of the bromoacetals 9 using *in situ* generated catalytic  $Bun_3SnH$  in the presence of AIBN followed by sonochemically accelerated reaction of the resultant spirohemiacetals 10 with Jones' reagent furnished  $\alpha$ -spiro- $\beta$ -methylene- $\gamma$ -butyrolactones 11. The results

OMe

OMe

AIBN

Scheme 1

sonication

Bun<sub>3</sub>SnCl

NaCNBH<sub>3</sub>



Scheme 2 Reagents and conditions: i,  $Ph_3P^+$ - $CH_2OMe^-Cl$ ,  $EtC(Me_2)$ O<sup>-</sup> +K, tetrahydrofuran (THF)-EtC(Me\_2) OH, room temp., 2-5 h; ii, NBS, HC=C-CR<sub>2</sub>OH (R = H or Me),  $CH_2Cl_2$ , -40 °C, 0.5 h; iii, Bu<sup>n</sup><sub>3</sub>SnCl (0.15 equiv.), NaCNBH<sub>3</sub> (2 equiv.), AIBN (catalytic), Bu<sup>t</sup>OH, reflux, 1.5 h; iv, 1.6 mol dm<sup>-3</sup> Jones' reagent, acetone, sonication, 5-15 min



Scheme 3 Reagents and conditions: i, LiAlH<sub>4</sub>, Et<sub>2</sub>O, -50 °C, 2 h, 95%; ii, (a) MeC(OEt)<sub>3</sub>, EtCO<sub>2</sub>H (catalytic), 180 °C, 6 days; (b) 10% aq. NaOH-MeOH (1:1), reflux, 7 h, 60% from 14; iii (a) C<sub>6</sub>H<sub>6</sub>, (COCl)<sub>2</sub>, room temp., 1 h; (b) CH<sub>2</sub>N<sub>2</sub>, Et<sub>2</sub>O, 2 h; (c) anhydrous CuSO<sub>4</sub>, cyclohexane, reflux (W-lamp), 4 h, 48% from the acid 15; iv, Li, liquid NH<sub>3</sub>, 15 min, 80%; v, same as Scheme 2, for yields see Table 1

are summarised in Table 1. It is worth mentioning that in entries vi-viii, the radical cyclisation reaction proceeded with a high degree of stereoselectivity *via* the preferred less crowded axial radical, and yielded only one isomer of the butyrolactone after reaction with Jones' reagent.

For the synthesis of norbakkenolide 2, the hydrindanone<sup>8</sup> 12 was identified as the requisite precursor, for which a new synthesis has been developed as depicted in Scheme 3. Regiospecific reduction of 3-methylcyclohexenone 13 with  $LiAlH_4$  gave the allyl alcohol 14. The ortho ester Claisen rearrangement9 of the alcohol 14 with triethyl orthoacetate in the presence of a catalytic amount of propionic acid, followed by hydrolysis of the resultant ester furnished the acid 15 in 60% yield. Treatment of the acid chloride derived from the acid 15, with an excess of ethereal diazomethane generated the diazo ketone 16. Anhydrous copper sulfate catalysed decomposition<sup>10</sup> of the diazo ketone 16 in refluxing cyclohexane (W-lamp) resulted in the cyclopropyl ketone 17 (48% from the acid 15).† Regiospecific cleavage of the cyclopropane ring in 17 with lithium in liquid ammonia produced the hydrindanone<sup>8</sup> 12 in 80% yield. Finally, spirannulation as described above (Scheme 2), transformed the hydrindanone 12 into a 5:1 mixture of norbakkenolide 2<sup>+</sup> and its C-7 epimer. The structure of the norbakkenolide was established based on the expected thermodynamically less crowded transition state in the radical cyclisation reaction, which was further confirmed by the NMR comparison with that of bakkenolide [two triplet (J ca. 2 Hz) signals due to olefinic protons appeared for **2** at  $\delta$  4.99 and 5.06; for the minor epimer at 5.05 and 5.2; whereas for bakkenolide<sup>1</sup> at 5.0 and 5.07].

In conclusion, an efficient method for transforming cyclic ketones into  $\alpha$ -spiro- $\beta$ -methylene- $\gamma$ -butyrolactones using a radical cyclisation reaction as the key step, and its extension to the synthesis of (±)-norbakkenolide have been achieved. Since the 3-methylcyclohexenone can be enantioselectively reduced<sup>11</sup> to the 3-methylcyclohexenol by LiAlH<sub>4</sub> in the presence of chiral *N*-methylephedrine, the present route provides a convenient route to chiral norbakkenolide as well. We thank Professor A. E. Greene for providing the <sup>1</sup>H NMR (200 MHz) spectrum of bakkenolide. S. N. and

† Spectral data for bromoacetal **9a**:  $v_{max}/cm^{-1}$  (neat) 3290, 2120, 1449, 1194, 1122.  $\delta_H$  (200 MHz, CDCl<sub>3</sub>) 4.54 (1H, s, O-CH-O), 4.43 (2H, d, J 2.4 Hz, O-CH<sub>2</sub>), 3.66 (3H, s, OMe), 2.49 (1H, t, J 2.4 Hz, C=CH), 1.5-2.2 (10H, m).  $\delta_{\rm C}$  (22.5 MHz, CDCl<sub>3</sub>) 107.8 (d, O-CH-O), 78.9 (s, C=CH), 75.6 (s, C-Br), 75.0 (C=CH), 58.9 (q, OMe), 56.1 (t, O-CH<sub>2</sub>), 34.4 (t), 34.0 (t), 25.2 (t), 21.8 (2C, t). For hemiacetal **10a**:  $v_{max}/cm^{-1}$  (CCl<sub>4</sub>) 3076, 1182, 1095, 1023, 945, 933, 882. δ<sub>H</sub> (200 MHz, CDCl<sub>3</sub>) 4.88 (3H, m, O-CH-O and =CH<sub>2</sub>), 4.49 and 4.39 (2H, t of AB q, J 13 and 2 Hz, 0-CH<sub>2</sub>), 3.38 (3H, s, OMe), 0.8-2.0 (10H, m), δ<sub>C</sub> (67.5 MHz, CDCl<sub>3</sub>) 156.0 (C=CH<sub>2</sub>), 108.8 (C=CH<sub>2</sub>), 102.8 (O-CH-O), 69.9 (O-CH<sub>2</sub>), 55.5 (OMe), 50.9 (spiro C), 37.1, 30.0, 26.5, 24.3 (2C). For lactone 11a:  $v_{max}/cm^{-1}$  (neat) 1776, 1671, 1452, 1353, 1170, 1113, 1029, 900.  $\delta_{\rm H}$  (200 MHz, CDCl<sub>3</sub>) 5.17 (1H, t, J 2.1 Hz) and 5.08 (1H, t, J 2 Hz) (C=CH<sub>2</sub>), 4.76 (2H, t, J 2 Hz, O–CH<sub>2</sub>), 1.35–2.0 (10H, m).  $\delta_{\rm C}$  (22.5 MHz, CDCl<sub>3</sub>) 179.7 (s, O=C–O), 149.1 (s, C=CH<sub>2</sub>), 107.1 (t, C=CH<sub>2</sub>), 69.1 (t, O–CH<sub>2</sub>), 44.6 (s, spiro C), 33.2 (2C, t), 25.0 (t), 20.9 (2C, t). For cyclopropyl ketone 17:  $v_{max}$  cm<sup>-1</sup> (neat) 1722.  $\delta_{H}$  (90 MHz, CDCl<sub>3</sub>) 2.3 and 2.0 (2H, AB q, J 16 Hz, O=CCH<sub>2</sub>), 1.0–2.1 (9H, m), 1.37 (3H, s, Me).  $\delta_{C}$  (100  $\begin{array}{l} MHz, \ CDCl_3) \ 214.9 \ (s, \ C=O), \ 56.4 \ (t, \ O=CCH_2), \ 35.4 \ (d), \ 35.3 \ (d), \\ 32.7 \ (d), \ 32.4 \ (s, \ C-Me), \ 30.3 \ (q, \ Me), \ 24.3 \ (d), \ 20.3 \ (t), \ 18.1 \ (t). \ For \end{array}$ **2**:  $v_{max}/cm^{-1}$  (neat) 1776, 1674, 1467, 1449, 1350, 1233, 1146, 1128, 1113, 1026, 894.  $\delta_{\rm H}$  (400 MHz, CDCl\_3) 5.06 (1H, t, J 2.1 Hz) and 4.99 (1H, t, J 1.6 Hz) (C=CH<sub>2</sub>), 4.78 (2H, t of AB q, J ca. 14 and 2 Hz, O-CH<sub>2</sub>), 2.4 (1H, t, J 12.6 Hz, H-9a), 2.13 and 1.67 (2H, AB q, J 13.7 Hz, H-6), 1.9 (1H, m, H-10), 1.72 (1H, dd, J 12.7 and 6.2 Hz, H-9b), 1.2–1.7 (8H, m), 1.11 (3H, s, Me).  $\delta_{\rm C}$  (50 MHz, CDCl<sub>3</sub>) 182.0 (O=C–O), 151.6 (*C*=CH<sub>2</sub>), 105.4 (C=CH<sub>2</sub>), 69.9 (OCH<sub>2</sub>), 52.6 (C-6), 49.7 (spiro C), 44.7 (C-10), 41.7 (C-9), 40.7 (C-5), 33.0 (C-4), 24.9 (Me), 23.5, 22.3, 20.3. MS: *m/z* 220 (M<sup>+</sup>, 75%), 133 (30), 111 (64), 110 (100), 95 (45), 81 (45).

#### J. CHEM. SOC., CHEM. COMMUN., 1993

G. V. R. S. thank the CSIR, New Delhi for the award of research fellowships.

Received, 23rd October 1992; Com. 2/05666F

#### References

- N. Abe, R. Onoda, K. Shirahata, T. Kato, M. C. Woods and Y. Kitahara, *Tetrahedron Lett.*, 1968, 369. So far only two total synthetic approaches are known for Bakkenolide-A: D. A. Evans, C. L. Sims and G. C. Andrews, *J. Am. Chem. Soc.*, 1977, 99, 5453; A. E. Greene, J.-P. Depres, F. Coelho and T. J. Brocksom, *J. Org. Chem.*, 1985, 50, 3943; *Tetrahedron Lett.*, 1988, 29, 5661.
- 2 K. Kano, K. Hayashi and H. Mitsuhashi, *Chem. Pharm. Bull.*, 1982, **30**, 1198 and references cited therein.
- 3 N. Petragnani, H. M. C. Ferraz and G. V. J. Silva, Synthesis, 1986, 157; A. Srikrishna, J. Chem. Soc., Chem. Commun., 1987, 587; A. R. Sidduri and P. Knochel, J. Am. Chem. Soc., 1992, 114, 7579 and references cited therein.
- 4 O. Moriya, M. Okawara and Y. Ueno, *Chem. Lett.*, 1984, 1437; A. Srikrishna and G. V. R. Sharma, *Tetrahedron Lett.*, 1988, 29, 6487; A. E. Greene, F. Coelho and J.-P. Depres, *J. Org. Chem.*, 1985, 50, 1973 and references cited therein.
- 5 G. Stork and R. Mook, Jr., J. Am. Chem. Soc., 1983, 105, 3720; G. Stork, R. Mook, Jr., S. A. Biller and S. D. Rychnovsky, J. Am.

Chem. Soc., 1983, 105, 3741; G. Stork and P. M. Sher, J. Am. Chem. Soc., 1983, 105, 6765; G. Stork and M. Kahn, J. Am. Chem. Soc., 1985, 107, 500; G. Stork and R. Mook, Jr., J. Am. Chem. Soc., 1987, 109, 2829; G. Stork and R. Mah, Tetrahedron Lett., 1989, 30, 3609; G. Stork, Bull. Chem. Soc. Jpn., 1988, 61, 149; Bull. Soc. Chim. Fr., 1990, 127, 675; see also Y. Ueno, O. Moriya, K. Chino, M. Watanabe and M. Okawara, J. Chem. Soc., Perkin Trans. I, 1986, 1351 and references cited therein. For general reviews on radical cyclisation reactions, see: B. Giese, Radicals in organic synthesis: Formation of carbon-carbon bonds, Pergamon Press, Oxford, 1986; M. Ramaiah, Tetrahedron, 1987, 43, 3541; D. P. Curran, Synthesis, 1988, 417; 489, C. P. Jasperse, D. P. Curran and T. L. Fevig, Chem. Rev., 1991, 91, 1237.

- 6 G. Stork and P. M. Sher, J. Am. Chem. Soc., 1986, 108, 303; A. Srikrishna and G. Sundarababu, *Tetrahedron*, 1990, 46, 7901 and references cited therein.
- A. Srikrishna, S. Nagaraju and G. V. R. Sharma, Synth. Commun., 1992, 22, 1127; 1221.
  R. Fraisse-Jullien, C. Frejaville and V. Toure, Bull. Soc. Chim.
- 8 R. Fraisse-Jullien, C. Frejaville and V. Toure, *Bull. Soc. Chim. Fr.*, 1966, 3725; Y. Ito, H. Aoyama and T. Saegusa, *J. Am. Chem. Soc.*, 1980, **102**, 4519.
- 9 W. S. Johnson, L. Werthemann, W. R. Bartlett, T. Li, D. J. Faulkner and M. R. Petersen, J. Am. Chem. Soc., 1970, 92, 741.
- 10 G. Stork and J. Ficini, J. Am. Chem. Soc., 1961, 83, 4678.
- 11 M. Kawasaki, Y. Suzuki and S. Terashima, Chem. Lett., 1984, 239.