

Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: G. Erker, L. Wang, G. Kehr, C. G. Daniliuc, M. Brinkkötter, T. Wiegand, A. Wübker, H. Eckert, L. Liu, J. G. Brandenburg and S. Grimme, *Chem. Sci.*, 2018, DOI: 10.1039/C8SC01089G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Chemical Science

EDGE ARTICLE

Solid State Frustrated Lewis Pair Chemistry

Long Wang,^a Gerald Kehr,^a Constantin G. Daniliuc,^a Melanie Brinkkötter,^b Thomas Wiegand,^c Anna-Lena Wübker,^b Hellmut Eckert,^{b,d,*} Lei Liu,^f Jan Gerit Brandenburg,^{e,f} Stefan Grimme,^{f,*} Gerhard Erker^{a,*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

In solution the PCy₃/B(C₆F₅)₃ pair is rapidly deactivated by nucleophilic aromatic substitution. In the solid state deactivation is effectively suppressed and the active frustrated phosphane/borane Lewis pair splits dihydrogen or adds to sulfur dioxide. A variety of phosphane/B(C₆F₅)₃ pairs have been used to carry out active FLP reactions in the solid state. The reactions were analyzed by DFT calculations and by solid state NMR spectroscopy. The solid state dihydrogen splitting reaction was also carried out under near to ambient conditions with suspensions of the non-quenched phosphane/borane mixtures in the fluorous liquid perfluoromethylcyclohexane.

Introduction

Lewis acids and bases when brought together in solution typically undergo rapid formation of strong adducts. Similar to the neutralization reaction of Brønsted acids and bases this leads to an annihilation of the typical Lewis acid and Lewis base properties.^{1,2} This situation can be changed if one effectively hinders the Lewis pair from the neutralizing adduct formation, e.g. by electronic means³ or, more commonly, by attaching very bulky substituents at the core atoms of the pair.⁴⁻⁶ This invariably leads to situations where active Lewis acids and active Lewis bases are present in a solution at the same time, opening possibilities for cooperative reactions with added substrates. Such “frustrated Lewis pairs” (FLPs)⁷ have very successfully been used within the last decade for small molecule binding and activation, most notably among this the metal-free splitting and activation of dihydrogen.^{8,9} A number of new reaction types have been found in this way^{10,11} and a great variety of different FLPs have been devised,

characterized and their reactions reported.¹²⁻²³ The vast majority of them relies on avoiding the effective neutralizing Lewis acid/Lewis base adduct formation by steric hindrance, and this has led to the discovery and development of a great variety of interesting reactions in solution.²⁴⁻²⁶

We thought that we should not confine ourselves to searching for FLP reactions in the liquid phase. One may safely assume that large molecules in the solid state are more or less rigidly confined to their positions in the crystal lattice. Therefore, Lewis acids and Lewis bases should effectively be hindered from adduct formation or other deactivating reaction pathways even in the absence of efficient steric hindrance by their substituents as long as we keep them in the solid state. This may actually set the scene for possibly finding new frustrated Lewis pairs and, consequently, new FLP reactions, by exposing such Lewis pairs to suitable reagents in the solid state. We have tried this principle and found that FLP chemistry can be done in this way.

There had been some reports about frustrated Lewis pair related behavior at certain heterogeneous catalysts. In these cases the active sites were part of the catalytic solid. Activation of small molecules at such systems had been achieved either thermally or by photolysis.²⁷⁻²⁹ There have also been a few reports about heterogenized Lewis acids, bonded to suitable supports, that have been employed in FLP type reactions.³⁰ Our here reported case is distinctly different: we have employed solid physical mixtures of phosphane Lewis bases with the strong B(C₆F₅)₃ boron Lewis acid and reacted them under suitable conditions with selected small molecules. Activation occurred and the phosphane/borane pair became consumed with selective formation of the FLP reaction products. First examples of this FLP development will be described below in this account.

^a Organisch-Chemisches Institut, Westfälische Wilhelms-Universität Münster, Corrensstraße 40, 48149 Münster, Germany. Email: erker@uni-muenster.de

^b Institut für Physikalische Chemie and Graduate School of Chemistry, Westfälische Wilhelms-Universität Münster, Corrensstraße 30, 48149 Münster, Germany, Email: eckerth@uni-muenster.de

^c Laboratorium für Physikalische Chemie, ETH Zürich, Vladimir-Prelog-Weg 2, 8049 Zürich, Switzerland

^d Institute of Physics in Sao Carlos, University of Sao Paulo, CEP 369, Sao Carlos SP 13566-590, Brazil

^e London Centre for Nanotechnology, University College London, 17-19 Gordon Street London, WC1H 0AH, United Kingdom


^f Mulliken Center for Theoretical Chemistry, Institut für Physikalische und Theoretische Chemie, Universität Bonn, Beringstraße 4, 53115 Bonn, Germany, Email: grimme@thch.uni-bonn.de

† Electronic Supplementary Information (ESI) available: additional experimental details, further spectral and crystallographic data, additional data from the solid state NMR and theoretical studies. CCDC 1515080–1515082. See DOI: xxxxxxxxxxxx



Results and discussion

Tricyclohexylphosphane (**1a**) and tris(pentafluorophenyl)-borane (**2**) represent a Lewis base/Lewis acid pair that is known to react rapidly in solution. It undergoes a nucleophilic aromatic substitution by the phosphane at the para-position of a C₆F₅ group of the borane^{4,31} to eventually yield the reaction product **3a**.^{7,32} This favorable S_NAr reaction eliminates any FLP reactivity of the **1a/2** pair very effectively in solution. Even in the presence of 50 bar H₂ no significant hydrogen splitting is observed, the formation of **3a** even prevails under these conditions. We isolated this compound in 86 % yield and confirmed its formation by X-ray diffraction and by NMR spectroscopy [Scheme 1 (a) (H₂); for details see the ESI].³² The situation is drastically different in the solid state: we mixed equimolar quantities of PCy₃ (**1a**) and B(C₆F₅)₃ (**2**) and exposed it in a glass vial inside a steel autoclave to 50 bar of dihydrogen for a total of 10 days with constant agitation by a Teflon coated magnetic stirring bar. After this time a sample was taken and dissolved in D₂-dichloromethane. The NMR analysis revealed the hydrogen splitting product [HPCy₃⁺][HB(C₆F₅)₃⁻] (**4a**) had been formed as the by far major product [Scheme 1 (b) (H₂)]. Only negligible if any amounts of the S_NAr product **3a**, which would have been formed readily from any residual PCy₃/B(C₆F₅)₃ upon dissolving in dichloromethane, were present in the *in situ* samples. The phosphonium/ hydridoborate product **4a** was identified by its typical ³¹P (δ 33.2, ¹J_{PH} ~ 443 Hz) and ¹¹B (δ -25.3, ¹J_{BH} ~ 92 Hz) NMR signals with correlated ¹H NMR features at δ 5.15 (d, ¹J_{PH} = 444.0 Hz) and δ 3.59 (br 1:1:1:1 q, [B]H), respectively. Workup of a representative sample eventually furnished the salt **4a** isolated in 81% yield on a 100 mg scale. Crystallization from dichloromethane/pentane gave single crystals which were used to confirm the formation of the FLP H₂ splitting product under these special conditions by X-ray diffraction (for details see the ESI).³³ The salt **4a** is an active reducing agent. Its reaction with the bulky N-phenyl-4-methylacetophenon-imine gave the respective secondary amine reduction product (24h at 70°C, 86% conversion, for details see the ESI).



Scheme 1. Reactions of the PR¹R²R³/B(C₆F₅)₃ FLP systems in a dihydrogen and SO₂ atmosphere, respectively, in solution (a) and in the solid state (b). [Cy: cyclohexyl]

We carried out an ample characterization of the solid product material directly (i.e. without ever dissolving it) by solid state NMR spectroscopy. Evidence for the solid state hydrogenation of the PCy₃/B(C₆F₅)₃ mixture comes from the ³¹P and ¹¹B MAS NMR data (Figure 1). In the ³¹P NMR spectrum we recognize the starting material at 7.1 ppm, whereas the phosphonium ion gives a broad signal at 30.1 ppm. In the ¹¹B MAS NMR spectrum the signal of the free B(C₆F₅)₃ gives rise to the previously documented second-order quadrupolar lineshape³⁴ whereas after the hydrogenation a much narrower signal appears at -24.9 ppm. In addition, the spectrum reveals the presence of a minor amount of the substitution product at -2.4 ppm. The latter is the only product formed when the reaction is carried out in solution. The complete NMR characterization of the substitution product both in solution and the solid state is given in the ESI. In a control experiment, the formation of only a very minor amount of the substitution product **3a** was observed in the solid state NMR spectra of a PCy₃/B(C₆F₅)₃ mixture subjected to identical reaction conditions in the absence of H₂.




Figure 1. ¹¹B{¹H} MAS (left) and ³¹P{¹H} CPMAS NMR spectra (right) of the B(C₆F₅)₃/PCy₃ control mixture sample (grey trace) and the hydrogenation sample (black trace) from the solid-state reaction with H₂. Minor side products are labelled by +. The product **3a** of the S_NAr reaction is indicated by an asterisk.

Figure 2 shows the ¹H MAS NMR spectrum of the FLP-H₂ adduct **4a** acquired at 20.0 T and a MAS spinning frequency of 60.0 kHz with the EASY scheme for suppression of background signals from the MAS probe and the MAS rotor cap.³⁵ Under these conditions, the strong ¹H-¹H dipolar couplings are sufficiently suppressed, even though residual line broadening is still detected owing to higher order terms in the Hamiltonian which are not fully eliminated even at 60.0 kHz.³⁶ A distinct doublet (*J*(¹H-³¹P) ~ 430 Hz) can be identified at 5.5 ppm which is assigned to P-bound hydrogen, whereas the singlet at 4.0 ppm is assigned to the B-bound hydrogen (the expected multiplet is not resolved in this case).






Figure 2. ^1H MAS NMR spectrum of a $\text{B}(\text{C}_6\text{F}_5)_3/\text{PCy}_3$ mixture after the solid-state reaction with H_2 acquired using the EASY scheme for probe and rotor cap background suppression measured at 20.0 T and a MAS frequency of 60.0 kHz (a) and corresponding line shape simulation (b). + marks a small signal remaining from the rotor cap.

This assignment is supported by $^{11}\text{B}\{^1\text{H}\}$ and $^{31}\text{P}\{^1\text{H}\}$ heteronuclear correlation experiments (Figure 3) which show intense cross-peaks linking these resonances to the corresponding ^{31}P and ^{11}B NMR signals of the solid-state hydrogenation product. Further support for this assignment comes from $^1\text{H}\{^{11}\text{B}\}$ REAPDOR experiments. The obtained peak assignments are in agreement with solution-state NMR data (vide supra), as well as DFT computations of ^1H NMR chemical shifts for the isolated cationic H-PCy_3^+ and anionic $\text{H-B}(\text{C}_6\text{F}_5)_3^-$ species (5.4 and 4.5 ppm on a B3-LYP/def2-TZVP level of theory, respectively). The absence of an encounter complex is proven by $^{31}\text{P}\{^{11}\text{B}\}$ REAPDOR and $^{11}\text{B}\{^{31}\text{P}\}$ REDOR experiments (see the ESI). As previously discussed, such experiments can probe the B...P distance by measuring the strength of the heteronuclear ^{11}B - ^{31}P dipole-dipole interactions in both FLPs and their reaction products.³⁷ In the present material, no dephasing was observed over a dipolar mixing time of ~ 5 ms. Comparing these experimental data with corresponding two-spin simulations we can conclude that the boron-phosphorus distance must be larger than 600 pm. Thus, all the experimental data are consistent with well-separated phosphonium and borate ions.




Figure 3. $^1\text{H}/^{31}\text{P}$ (left) and $^1\text{H}/^{11}\text{B}$ (right) HETCOR spectra of a $\text{HB}(\text{C}_6\text{F}_5)_3/\text{HPCy}_3^+$ mixture **4a** after the solid-state reaction with H_2 . Cross-peaks denote the B-H and P-H correlations. The additional $^1\text{H}/^{11}\text{B}$ crosspeak seen in the right part of the figure at (1.3/-5.2) ppm arises from the substitution product **3a**.

The reaction of the phosphane PPhCy_2 (**1b**) with the Lewis acid $\text{B}(\text{C}_6\text{F}_5)_3$ (**2**) in the solid state proceeds similarly. The reaction

was carried out analogously as the one described above (r.t., 50 bar H_2 , 3 days). Our analysis of a product sample dissolved in CD_2Cl_2 revealed almost exclusive formation of the dihydrogen FLP splitting product **4b** [Scheme 1b (H_2)]. It shows a characteristic ^1H NMR [P]H doublet at δ 6.13, $^1J_{\text{PH}} = 459.3$ Hz (^{31}P : δ 30.6) and a broad 1:1:1:1 intensity [B]H quartet at δ 3.64 (^{11}B : δ -25.3, d, $^1J_{\text{BH}} \sim 94$ Hz).

Keeping the $\text{PPhCy}_2/\text{B}(\text{C}_6\text{F}_5)_3$ mixture (**1b/2**) in CD_2Cl_2 solution for 12 hours under H_2 (50 bar) gave a different result. The NMR analysis showed the formation of a ca. 1:6 mixture of the salt **4b** with the substitution product **3b**. [Scheme 1a (H_2)] The major zwitterionic phosphonium/fluoroborate product **3b** was eventually isolated as a white solid in 82% yield. It shows a ^{11}B NMR doublet ($^1J_{\text{BF}} \sim 70$ Hz) at δ -0.6 and a sharp ^{31}P NMR signal at δ 33.6. The ^{19}F NMR spectrum shows the [B]F resonance at -192.9, two signals of the bridging C_6F_4 group and three resonances (o, p, m) of the remaining $\text{B}(\text{C}_6\text{F}_5)_2$ unit with a $\Delta\delta^{19}\text{F}_{\text{m,p}}$ shift difference of 4.9 ppm, which is typical for a borate type structure (for details and the depicted NMR spectra see the ESI).

In this case the nucleophilic aromatic substitution by the markedly less nucleophilic phosphane PPhCy_2 (**1b**) relative to PCy_3 (**1a**) seems to allow the FLP reaction to compete as a minor pathway in solution. This becomes continued in the reaction of the much less nucleophilic phosphane $\text{PPh}_2(\text{tBu})$ (**1c**) which in solution together with the $\text{B}(\text{C}_6\text{F}_5)_3$ Lewis acid gave a ratio of the $\text{S}_{\text{N}}\text{Ar}$ and FLP products of **3c** : **4c** \sim 2:1 under our typical reaction conditions (r.t., CH_2Cl_2 solution, 12 hours, 50 bar H_2). [Scheme 1a (H_2)] Without H_2 only the substitution product (**3c**) was formed in solution (isolated in 76% yield, see the ESI for its characterization). On the contrary, the reaction of the $\text{PPh}_2(\text{tBu})/\text{B}(\text{C}_6\text{F}_5)_3$ Lewis base / Lewis acid mixture in the solid state (r.t., 50 bar H_2 , 3 days) gave almost pure H_2 -splitting product **4c**, [Scheme 1b (H_2)] [^{31}P NMR: δ 31.4 ($^1J_{\text{PH}} \sim 474$ Hz), ^{11}B : δ -25.2 ($^1J_{\text{BH}} \sim 91$ Hz), ^{19}F : $\Delta\delta^{19}\text{F}_{\text{m,p}} = 3.0$ ppm] which we isolated from the workup procedure involving recrystallization from CH_2Cl_2 /pentane in 60% yield (details of the characterization of the compounds **3c** and **4c** see the ESI). Quantum chemical simulations were used to investigate the mechanistic details of the different FLP reactivities in all three states of matter, i.e. gas, liquid, and solid phase. The FLP/ H_2 thermochemistry is for the first time investigated here in the solid state using relatively high-level periodic quantum chemistry methods. We employed a hierarchy of theoretical methods, ranging from semi-empirical tight-binding Hamiltonians to accurate London dispersion corrected hybrid density functionals.³⁸⁻⁴⁶ More discussion of methodological points and computational details can be found in our previous benchmark study⁴⁷ and in the ESI. The main representative results of the $\text{PCy}_3/\text{B}(\text{C}_6\text{F}_5)_3$ FLP are shown in Figure 4.






Figure 4. Calculated Gibbs free energies for the reaction of $\text{PCy}_3/\text{B}(\text{C}_6\text{F}_5)_3$ with H_2 in the solution (toluene) and in the solid state (see the supplementary material for computational details). All energy values are given in kcal mol^{-1} . Inset figures are the overlays of the HF-3c calculated crystal structures of $\text{PCy}_3/\text{B}(\text{C}_6\text{F}_5)_3$ (green) and $[\text{HPCy}_3][\text{HB}(\text{C}_6\text{F}_5)_3]$ (blue) (a), and crystal structures of $\text{PCy}_3/\text{B}(\text{C}_6\text{F}_5)_3$ (green) and the $\text{S}_{\text{N}2\text{Ar}}$ product **3a** (orange) (b). Hydrogen atoms except P-H and B-H are omitted for clarity.

All shown reactions are exergonic, and both the solvent and the crystal phase stabilize the products **4a** and **3a** by 10 to 15 kcal/mol compared to the gas phase. In this regard the FLP reaction to **4a** is in fact thermodynamically feasible. However, the competing product **3a** is in solution significantly preferred over **4a** ($-50.7 \text{ kcal mol}^{-1}$ vs. $-15.5 \text{ kcal mol}^{-1}$) and due to the expected high mobility it can readily react and prevent the desired FLP reaction. In contrast, the crystal field provides much more pronounced energy barriers, which kinetically stabilizes the reactant and enables the targeted reaction to product **4a**. These higher energy barriers can be rationalized by a simple geometric comparison (inset of Figure 4). Apparently, the solid state reaction to **4a** requires substantially less rearrangements of the crystal compared to **3a**. Thus, we can identify two key roles that drive the reaction in the solid phase: (1) The crystal environment can adopt the solvent role in enhancing the FLP reactivity, typically explained by both the electrostatic screening and unidirectional London dispersion stabilization of the FLP products. (2) The static crystal field can selectively suppress certain undesired reaction routes, which is not possible in a liquid or gas environment with high molecular mobility.

While point (1) makes the heterogeneous formulation of typical FLP reactions possible, (2) goes beyond it and opens possibilities for new FLP systems as compellingly demonstrated for the here discussed compound. An additional important prerequisite for the discussed reaction is the possible diffusion of H_2 gas through the reactant crystal. Our molecular dynamics (MD) simulation (for details see the ESI) confirm that H_2 can actually move more or less freely through the channels of the crystal thereby generating the correct conditions for the H_2 activation to take place. Moreover, we have conducted MD simulations for a model of the interface between the Lewis acid and base as it may occur

experimentally in a mixture of the solid particles. According to these results which are shown in Figure 5, at the interface the components (in particular the Lewis base) are spatially not constrained, possibly due to a mismatch of the molecular surfaces (a kind of interface strain). This leads partially to a "liquid phase" behavior of the FLPs in the solid state. It is seen that the Lewis acid and base components could move rather freely at the contact surface and adopt molecular FLP conformations enabling hydrogen activation as in solution. (for details see the ESI)




Figure 5. Snapshots of the periodic BO-MD simulation at the DFTB-D3 level of theory for the $\text{PCy}_3 + \text{B}(\text{C}_6\text{F}_5)_3$ FLP (**1a/2**). Color legend: P yellow, B pink, C black, F green and H white.

The new solid state phosphane / borane FLP reactions are not limited to the splitting of dihydrogen. We exposed the 1:1 mixture of PCy_3 (**1a**) and $\text{B}(\text{C}_6\text{F}_5)_3$ (**2**) for 4 hours at r.t. in the solid state to SO_2 gas (1.5 bar).⁴⁸ A sample was dissolved in CD_2Cl_2 and its NMR spectra revealed the almost quantitative formation of the P/B FLP SO_2 addition product **5**. [Scheme 1 (b) (SO_2)] The product was also directly identified from the solid obtained by solid state NMR spectroscopy (see the ESI), indicating essentially quantitative conversion. The distorted four-coordinate boron environment in **5** is characterized by $\delta_{\text{iso}} = -0.6 \text{ ppm}$, and a nearly axially symmetric electric field gradient, with $C_Q = 1.54 \text{ MHz}$ and $\eta_Q = 0.15$. The ^{31}P MAS-NMR spectrum shows a single sharp signal at 51.5 ppm. In this case, $^{11}\text{B}\{^{31}\text{P}\}$ REDOR and $^{31}\text{P}\{^{11}\text{B}\}$ REAPDOR experiments consistently point towards a B...P internuclear distance of 450 pm, which is in good agreement with the distance of 434 pm from the crystal structure.

We performed the reaction on a preparative scale and isolated the product **5** as a white solid after recrystallization from CH_2Cl_2 / pentane in 84 % yield. The product shows the typical ^{11}B ($\delta -0.3$) and ^{19}F NMR features (three resonances, $\Delta\delta^{19}\text{F}_{\text{m,p}} = 6.4 \text{ ppm}$) of the borate section of the molecule and a phosphonium ^{31}P NMR signal at $\delta 50.0$ (for further details see the ESI).






Figure 6. Molecular structure of the solid state P/B FLP SO_2 addition product **5**. Selected bond lengths (Å): S1-O1 1.445(2), S1-O2 1.564(2), P1-S1 2.268(1), B1-O2 1.545(3) and bond angles (°): O1-S1-O2 110.8(1), O1-S1-P1 106.3(1), O2-S1-P1 93.4(1), $\Sigma\text{B1}^{\text{ccc}}$ 334.0, $\Sigma\text{P1}^{\text{ccc}}$ 334.1.

Compound **5** was characterized by X-ray diffraction (Figure 6). The X-ray crystal structure analysis shows the newly formed P1-S1 and O1-B1 bonds. Both the phosphorus and the boron atoms show pseudo-tetrahedral coordination geometries. The sulfur coordination geometry is distorted trigonal-pyramidal. We also exposed the $\text{PCy}_3/\text{B}(\text{C}_6\text{F}_5)_3$ pair to SO_2 in solution but only observed the formation of the substitution product **3a** [Scheme 1 (a) (SO_2); see the ESI for details].

Our study has shown so far that an agitated mixture of particles of the phosphanes **1a-c** with particles of the $\text{B}(\text{C}_6\text{F}_5)_3$ Lewis acid **2** did very effectively evade the deactivating $\text{S}_{\text{N}}\text{Ar}$ reaction that these pairs rapidly undergo in solution. Instead, they retained their frustrated Lewis pair character and, consequently, showed the ability to split dihydrogen heterolytically. While this observation is probably of a far-reaching principal interest, the rather harsh conditions of the solid state FLP H_2 -splitting (50 bars of dihydrogen, 3 to 10 d reaction time) made this far from a conveniently applicable procedure.

Fluorous liquids show some extraordinary properties.⁴⁹⁻⁵⁶ They do not mix with a variety of common organic solvents; they show an enhanced solubility of many gases in them, among them dihydrogen.^{49,56-58} Moreover, many organic and element-organic compounds, among them the phosphanes **1a-c** and $\text{B}(\text{C}_6\text{F}_5)_3$ (**2**) are insoluble in them. Therefore, we decided to carry out the solid state FLP dihydrogen splitting reaction in perfluoromethylcyclohexane ($\text{F}_{11}\text{C}_6\text{-CF}_3$). In a typical experiment (see the ESI for details) we suspended an equimolar mixture of PCy_3 (**1a**) and $\text{B}(\text{C}_6\text{F}_5)_3$ in perfluoromethylcyclohexane and stirred the suspension for 10 h in a dihydrogen atmosphere at near to ambient conditions (r.t., 1.5 bar H_2). Workup was simply done by evaporation of the volatiles. A sample of the obtained white powdery solid was then subjected to NMR analysis in D_2 -dichloromethane solution. It showed that a ca. 60% conversion to the hydrogen splitting product $\text{HPCy}_3^+/\text{HB}(\text{C}_6\text{F}_5)_3^-$ (**4a**) had been achieved. The remaining starting material had become converted to the $\text{S}_{\text{N}}\text{Ar}$ reaction product **3a** under the conditions of the NMR analysis in solution. The solid state NMR spectra of the products obtained after the suspension reaction were

essentially identical to those obtained from the solid-state reaction.

Table 1. A Comparison of the Dihydrogen Splitting Reaction by the Phosphane/Borane FLPs **1**/ $\text{B}(\text{C}_6\text{F}_5)_3$ in Solution, in the Dry Solid State and in the Fluorous Liquid Perfluoromethylcyclohexane.

	No.	solution ^c	dry solid	in $\text{F}_{11}\text{C}_6\text{-CF}_3$ ^b
PCy_3	1a	only 3a	> 90% conv. to 4a ^a	60% conv. to 4a
PPhCy_2	1b	3b : 4b ~ 6 : 1	ca. 95% conv. to 4b ^d	ca. 95% conv. to 4b
PPh_2^tBu	1c	3c : 4c ~ 2 : 1	ca. 95% conv. to 4c ^d	ca. 98% conv. to 4c

^a 10 days, r.t., 50 bar H_2 ; ^b 10 hours, r.t. 1.5 bar H_2 ; ^c d_2 -dichloromethane, 12 hours, r.t.; ^d 3 days, r.t., 50 bar H_2

The reaction of the PPhCy_2 (**1b**)/ $\text{B}(\text{C}_6\text{F}_5)_3$ pair with dihydrogen proceeded at least equally well in this fluorous liquid. Under analogous conditions a ca. 95% conversion to the H_2 -splitting product **4b** was achieved within the 10 h reaction time. The PPh_2^tBu (**1c**)/ $\text{B}(\text{C}_6\text{F}_5)_3$ system even slightly surpassed this result. We obtained a near to quantitative conversion to the $\text{HPPH}_{2t}\text{Bu}^+/\text{HB}(\text{C}_6\text{F}_5)_3^-$ salt within 10 h at near to ambient conditions in the inert perfluoromethylcyclohexane liquid (see the ESI for details).

Conclusions

Our study has shown that frustrated Lewis pair behavior can be achieved by other means than the usual electronic^{3,19} or steric modifications⁴⁻⁹ of Lewis acids and bases. In the cases reported here we have deviated from the ubiquitous method of using sterically bulky substituents at the phosphorus Lewis base in order to prevent neutralizing adduct formation or other deactivating reactions with the strong $\text{B}(\text{C}_6\text{F}_5)_3$ boron Lewis acid. On the contrary, the phosphanes used in this study are rather nucleophilic and in solution they undergo rapid nucleophilic aromatic substitution at one C_6F_5 ring of the Lewis acid replacing fluoride which consequently then becomes bonded to the boron atom³¹ with annihilation of its Lewis acidic features. In all the cases looked at in our study this FLP deactivating reaction is very efficiently suppressed by localizing the individual Lewis acidic and Lewis basic molecules inside their solid state lattices. This prevents them effectively from undergoing the bimolecular deactivation reaction.

From the solid state NMR and the DFT analysis in conjunction with general principles about molecular diffusion in the solid state, we assume that the individual Lewis acid and base components do not easily mix on a molecular level in our experiments, but that initially we are dealing with separate solid state phosphane and borane particles. This lets us assume that the FLP dihydrogen splitting reaction must take place at the surface, respectively the interface between phosphane and borane solid state entities; the ensuing reaction is, however, probably facilitated by the easy permeability of the respective crystal lattices by dihydrogen (and other gases). The dihydrogen splitting reaction may then have formed the phosphonium/hydridoborate salt initially at the surface, but it may be assumed that accumulation of that



ARTICLE

Journal Name

species might create a local situation resembling an ionic liquid, which might facilitate diffusion and mixing since eventually we obtained homogeneous solid samples of the respective dihydrogen splitting products. The MD simulations which support this view are currently due to the large system size too approximate (short) to draw any quantitative conclusions nor are we able to simulate further parts of the solid state reaction dynamically. Nevertheless, the MD and static theoretical results clearly support the above described picture of partially “molten” material at the interface with sufficient molecular flexibility for activation of almost freely diffusing molecular hydrogen.

Although the mechanistic aspects of our solid state FLP reactions must remain somewhat speculative at this time, we have greatly improved its practical applicability by using the fluorour liquid effect.⁵⁵⁻⁵⁷ This has made dihydrogen splitting reactions readily available from frustrated Lewis pair combinations which cannot be kept active in other ways.³³ We have further found first indications that the resulting $[P]H^+/[B]H^-$ product **4a** can be usefully employed in imine reduction and we have shown that the solid state FLP reactions are not confined to the dihydrogen splitting reactions but can be developed beyond. The solid state approach can make FLPs available for small molecule activation beyond using the conventional methods leading to Lewis pair “frustration”. It needs to be explored if this will open new pathways of extending FLP chemistry beyond its existing scope, for example by opening FLP routes to the large field of heterogeneous catalysis, here to be performed without the aid of metals.²⁷⁻³⁰

Conflicts of Interest

There are no conflicts to declare.

Acknowledgements

Financial support from the European Research Council (G.E.) and the Deutsche Forschungsgemeinschaft (Leibniz award (S.G.)) is gratefully acknowledged. We thank Prof. Beat Meier and the ETH Zürich for supporting the high field solid-state NMR measurements.

Notes and references

L.W. did the synthetic work, C. G. D. did the X-ray crystal structure analyses, G.K. supervised the work and checked the experimental data, G.E. initiated and supervised the study, M.B., T.W., A.-L.W. and H.E. did the solid state NMR studies, L.L, J.G.B and S.G. did the theoretical study.

- 1 F. H. Stephens, V. Pons and R. Tom Baker, *Dalton Trans.*, 2007, 2613-2626.
- 2 A. Staubitz, A. P. M. Robertson and I. Manners, *Chem. Rev.*, 2010, **110**, 4079-4124.
- 3 A. Stute, G. Kehr, C. G. Daniliuc, R. Fröhlich and G. Erker, *Dalton Trans.*, 2013, 42, 4487-4499.
- 4 G. C. Welch, R. R. S. Juan, J. D. Masuda and D. W. Stephan, *Science*, 2006, **314**, 1124-1126.

- 5 G. C. Welch and D. W. Stephan, *J. Am. Chem. Soc.*, 2007, **129**, 1880-1881.
- 6 P. Spies, G. Erker, G. Kehr, K. Bergander, R. Fröhlich, S. Grimme and D. W. Stephan, *Chem. Commun.*, 2007, 5072-5074.
- 7 G. C. Welch, L. Cabrera, P. A. Chase, E. Hollink, J. D. Masuda, P. Wei and D. W. Stephan, *Dalton Trans.*, 2007, 3407-3414.
- 8 D. W. Stephan and G. Erker, *Angew. Chem. Int. Ed.*, 2015, **54**, 6400-6441.
- 9 D. W. Stephan and G. Erker, *Top. Curr. Chem.*, 2013, **332**, 1-350.
- 10 Y. Hasegawa, C. G. Daniliuc, G. Kehr and G. Erker, *Angew. Chem. Int. Ed.*, 2014, **53**, 12168-12171.
- 11 M. Sajid, L.-M. Elmer, C. Rosorius, C. G. Daniliuc, S. Grimme, G. Kehr and G. Erker, *Angew. Chem. Int. Ed.*, 2013, **52**, 2243-2246.
- 12 M. Sajid, G. Kehr, C. G. Daniliuc and G. Erker, *Angew. Chem. Int. Ed.*, 2014, **53**, 1118-1121.
- 13 M. Sajid, A. Lawzer, W. Dong, C. Rosorius, W. Sander, B. Schirmer, S. Grimme, C. G. Daniliuc, G. Kehr and G. Erker, *J. Am. Chem. Soc.*, 2013, **135**, 18567-18574.
- 14 A. J. P. Cardenas, B. J. Culotta, T. H. Warren, S. Grimme, A. Stute, R. Fröhlich, G. Kehr and G. Erker, *Angew. Chem. Int. Ed.*, 2011, **50**, 7567-7571.
- 15 M. Sajid, A. Stute, A. J. P. Cardenas, B. J. Culotta, J. A. M. Hepperle, T. H. Warren, B. Schirmer, S. Grimme, A. Studer, C. G. Daniliuc, R. Fröhlich, J. L. Petersen, G. Kehr and G. Erker, *J. Am. Chem. Soc.*, 2012, **134**, 10156-10168.
- 16 R. Liedtke, F. Scheidt, J. Ren, B. Schirmer, A. J. P. Cardenas, C. G. Daniliuc, H. Eckert, T. H. Warren, S. Grimme, G. Kehr and G. Erker, *J. Am. Chem. Soc.*, 2014, **136**, 9014-9027.
- 17 K. Chernichenko, Á. Madarász, I. Pápai, M. Nieger, M. Leskelä and T. Repo, *Nat. Chem.*, 2013, **5**, 718-723.
- 18 C. B. Caputo, L. J. Hounjet, R. Dobrovetsky and D. W. Stephan, *Science*, 2013, **341**, 1374-1377.
- 19 M.-A. Courtemanche, M.-A. Légaré, L. Maron and F.-G. Fontaine, *J. Am. Chem. Soc.*, 2013, **135**, 9326-9329.
- 20 T. Wang, G. Kehr, L. Liu, S. Grimme, C. G. Daniliuc and G. Erker, *J. Am. Chem. Soc.*, 2016, **138**, 4302-4305.
- 21 E. Theuergarten, J. Schlosser, D. Schluns, M. Freytag, C. G. Daniliuc, P. G. Jones and M. Tamm, *Dalton Trans.*, 2012, **41**, 9101-9110.
- 22 E. L. Kolychev, T. Bannenber, M. Freytag, C. G. Daniliuc, P. G. Jones and M. Tamm, *Chem. Eur. J.*, 2012, **18**, 16938-16946.
- 23 B. Inés, D. Palomas, S. Holle, S. Steinberg, J. A. Nicasio and M. Alcarazo, *Angew. Chem. Int. Ed.*, 2012, **51**, 12367-12369.
- 24 T. Mahdi and D. W. Stephan, *J. Am. Chem. Soc.*, 2014, **136**, 15809-15812.
- 25 T. Mahdi and D. W. Stephan, *Angew. Chem. Int. Ed.*, 2015, **54**, 8511-8514.
- 26 D. J. Scott, M. J. Fuchter and A. E. Ashley, *J. Am. Chem. Soc.*, 2014, **136**, 15813-15816.
- 27 K. K. Ghuman, L. B. Hoch, P. Szymanski, J. Y. Y. Loh, N. P. Kherani, M. A. El-Sayed, G. A. Ozin and C. V. Singh, *J. Am. Chem. Soc.*, 2016, **138**, 1206-1214.
- 28 K. K. Ghuman, T. E. Wood, L. B. Hoch, C. A. Mims, G. A. Ozin and C. V. Singh, *Phys. Chem. Chem. Phys.*, 2015, **17**, 14623-14635.
- 29 (a) S. Zhang, Z.-Q. Huang, Y. Ma, W. Gao, J. Li, F. Cao, L. Li, C.-R. Chang and Y. Qu, *Nat. Commun.*, 2017, **8**, 15266, and references cited therein; (b) Z.-Q. Huang, L.-P. Liu, S. Qi, S. Zhang, Y. Qu and C.-R. Chang, *ACS Catal.*, 2018, **8**, 546-554. (c) Review: N. J. Coville and L. Cheng, *J. Organomet. Chem.*, 1998, **571**, 149-169;
- 30 (a) J.-Y. Xing, J.-C. Buffet, N. H. Rees, P. Norby and D. O'Hare, *Chem. Commun.*, 2016, **52**, 10478-10481; see also: (b) Y.-J. Wanglee, J. Hu, R. E. White, M.-Y. Lee, S. M. Stewart, P.



- Perrotin, S. L. Scott, *J. Am. Chem. Soc.* 2012, **134**, 355-366; (c) K. C. Szeto, W. Sahyoun, N. Merle, J. L. Castlebou, N. Popoff, F. Lefebvre, J. Raynaud, C. Godard, C. Claver, L. Delevoye, R. M. Gauvin, M. Taoufik, *Catal. Sci. Technol.* 2016, **6**, 882-889.
- 31 "S. Döring, G. Erker, R. Fröhlich, O. Meyer and K. Bergander, *Organometallics*, 1998, **17**, 2183-2187.
- 32 M. Ullrich, A. J. Lough and D. W. Stephan, *J. Am. Chem. Soc.*, 2009, **131**, 52-53.
- 33 Y. Jiang, B. Schirmer, O. Blacque, T. Fox, S. Grimme and H. Berke, *J. Am. Chem. Soc.*, 2013, **135**, 4088-4102.
- 34 T. Wiegand, H. Eckert, O. Ekkert, R. Fröhlich, G. Kehr, G. Erker and S. Grimme, *J. Am. Chem. Soc.*, 2012, **134**, 4236-4249.
- 35 C. Jaeger and F. Hemmann, *Solid State Nucl. Magn. Reson.*, 2014, **57-58**, 22-28.
- 36 A. Böckmann, M. Ernst and B. H. Meier, *J. Magn. Reson.*, 2015, **253**, 71-79.
- 37 T. Wiegand, M. Siedow, H. Eckert, G. Kehr and G. Erker, *Isr. J. Chem.*, 2015, **55**, 150-178.
- 38 R. Sure and S. Grimme, *J. Comput. Chem.*, 2013, **34**, 1672-1685.
- 39 J. Tao, J. P. Perdew, V. N. Staroverov and G. E. Scuseria, *Phys. Rev. Lett.*, 2003, **91**, 146401.
- 40 Y. Zhao and D. G. Truhlar, *J. Phys. Chem. A*, 2005, **109**, 5656-5667.
- 41 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 42 S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.*, 2011, **32**, 1456-1465.
- 43 M. Elstner, D. Porezag, G. Jungnickel, J. Elsner, M. Haugk, T. Frauenheim, S. Suhai and G. Seifert, *Phys. Rev. B*, 1998, **58**, 7260-7268.
- 44 M. Gaus, A. Goez and M. Elstner, *J. Chem. Theory Comput.*, 2013, **9**, 338-354.
- 45 J. G. Brandenburg and S. Grimme, *J. Phys. Chem. Lett.*, 2014, **5**, 1785-1789.
- 46 L. Liu, J. G. Brandenburg and S. Grimme, *Philos. Trans. A Math. Phys. Eng. Sci.*, 2017, **375**, 20170006.
- 47 M. Sajid, A. Klose, B. Birkmann, L. Liang, B. Schirmer, T. Wiegand, H. Eckert, A. J. Lough, R. Fröhlich, C. G. Daniliuc, S. Grimme, D. W. Stephan, G. Kehr and G. Erker, *Chem. Sci.*, 2013, **4**, 213-219.
- 48 J. A. Gladysz, D. P. Curran and I. T. Horvath, *Handbook of Fluorous Chemistry*. Wiley-VCH: Weinheim, 2004.
- 49 I. Ryu, H. Matsubara, C. Emnet, J. A. Gladysz, S. Takeuchi, Y. Nakamura and D. P. Curran, in *Green Reaction Media in Organic Synthesis*, Blackwell Publishing Ltd, 2007, pp. 59-124.
- 50 H. Nakamura, T. Usui, H. Kuroda, I. Ryu, H. Matsubara, S. Yasuda and D. P. Curran, *Org. Lett.*, 2003, **5**, 1167-1169.
- 51 D. P. Curran, *Aldrichimica Acta*. 2006, **39**, 3-9.
- 52 D. P. Curran, in *Stimulating Concepts in Chemistry*, Wiley-VCH Verlag GmbH & Co. KGaA, 2005, pp. 25-37.
- 53 A. Studer, S. Hadida, R. Ferritto, S.-Y. Kim, P. Jeger, P. Wipf and D. P. Curran, *Science*, 1997, **275**, 823-826.
- 54 A. P. Dobbs and M. R. Kimberley, *J. Fluorine Chem.*, 2002, **118**, 3-17.
- 55 E. de Wolf, G. van Koten and B.-J. Deelman, *Chem. Soc. Rev.*, 1999, **28**, 37-41.
- 56 J. A. Gladysz, and M. Jurisch, *Top. Curr. Chem.*, 2012, **308**, 1-24.
- 57 G. Serratrice, J. J. Delpuech, and R. Diguët, *Nouv. J. Chem.*, 1982, **6**, 489.

